A DICTIONARY OF SCIENTIFIC TERMS

PRONUNCIATION, DERIVATION, AND DEFINITION OF TERMS IN BIOLOGY, BOTANY, ZOOLOGY, ANATOMY, CYTOLOGY, GENETICS, EMBRYOLOGY, PHYSIOLOGY

BY

I. F. HENDERSON, M.A.

AND

W. D. HENDERSON, M.A., B.Sc., Ph.D., F.R.S.E.

SIXTH EDITION

RV

J. H. KENNETH, M.A., PH.D., F.R.S.E., F.R.S.G.S.

OLIVER AND BOYD

EDINBURGH: TWEEDDALE COURT LONDON: 39A WELBECK STREET, W. 1

PREFACE

IT was thought desirable that there should be available to students and others interested in the Biological Sciences an up-to-date work somewhat on the lines of Stormonth's Manual of Scientific Terms (1879, republished 1903). The subjects selected for treatment are Biology and its allies: Botany, Zoology, Cytology, Genetics, Embryology, Anatomy, Physiology; some terms in Bacteriology and Palæontology are included. Specific, generic, ordinal, and other systematic names of plants and animals have necessarily been omitted, nor could references to the sources of terms be listed in a volume of moderate size.

After compilation of most of the primary lists and completion of a small number of terms, the work was handed over to I. F. and W. D. Henderson, who prepared the First and Second Editions, and added to the definitions of nearly ten thousand terms those of some fifteen hundred new terms. Further revision was undertaken in the preparation of the Third Edition, and a list of abbreviations occurring in biological literature was added. Many definitions were amplified or modified, and the number of terms and cross-references was very considerably augmented, in the Fourth Edition, and a list of equivalent units of measurement was introduced. The text was again expanded in the Fifth Edition. In the present, Sixth Edition, the text has been further revised and extended.

The method of spelling is in the main that used in Britain, but due attention is paid to American orthography, by means of cross-references or by reproducing in the original lettering terms culled from scientific literature published in the United States. Spelling, however, is not static, as may be illustrated by the tendency to substitute e for the diphthongs ae and oe, and by compound words which may be written as two separate words, or hyphenated, or integrated as one word.

In the statement of derivation of terms, Greek and Russian words have been transliterated, as science and medical students are seldom acquainted with those languages. On the advice

of one authority on Greek, the transliteration of certain combinations of letters represents the sound rather than the exact letters of the original, hence the occurrence of such words as brangchia, hydor, etc.

Pronunciation, upon which there can never be universal agreement, is the least satisfactory portion, and is likely to remain so. Quite apart from natural differences in Scottish, English, and American utterance, pronunciation is subject to different fashions in different centres of learning. Moreover, pronunciation is subject to continual fluctuations; in the case of Latin terms, for instance, there is a trend toward the Continental type of pronunciation. The accepted position of the accent also varies in different localities and from one generation to another. Some of the alternative styles of pronunciation given will doubtless be regarded by the purist as faulty or definitely wrong, but their inclusion seems justified by the law of common usage.

The accretion of additional terms might have been dealt with in the form of Supplements, but with the generous consent of the Publishers, the work has been reset for each edition in order to retain the facility of reference afforded by an unbroken alphabetical arrangement, and to allow of revision, compression, or amplification of the text of the previous edition.

Innumerable publications have been probed for new terms and for new meanings which have been attached to existing terms. Thanks are due to many individuals who, during the past thirty-six years, have furthered the work by suggestions and criticisms. The preparation of each new edition has also been essentially facilitated by the courtesy of the staff of various libraries, particularly of the Library of the University of Edinburgh; and of the Argyll County Library, during preparation of the present edition.

This Dictionary now deals with approximately fourteen thousand terms. However, as I. I. Henderson pointed out in the Preface to the Second Edition, "a work of this nature must inevitably remain subject to limitations of time and resources".

SOUND-SYMBOLS USED IN PRONUNCIATION

The sound-symbols have been made as simple as possible, only the broader differences in vowel-sounds being included. The phonetic spelling following each term represents a general indication of the prevailing varieties of pronunciation rather than a critically exact reproduction, as the key-words themselves have by no means a universal interpretation.

ā	as in	rate.	ō	as in	no.
ă â	,,	rat.	ŏ	٠,	not.
â	,,	far.	ô	.,	form.
ch	,,	church.	o	,,	anatomy.
	,,	he.	oi	,,	toy.
č č ć	,,	hen.	00	,,	good.
ċ	,,	her.	ow		cow.
g	,,	go.	S	,,	so.
	,,	guano.	sh	,,	fish.
gw î	,,	pine.	th	,,	thin.
ĭ	,,	pin.	ū	,,	pure.
j	,,	gem.	ŭ	,.	nut.
j k	,,	cat.	y	,,	yard.
kw		queen.	z	,,	maize.
ng	,,	sing.	zh	"	vision.

ABBREVIATIONS

A androecium	bot botanical, botany
A . argon	B.P British Pharmacopoeia
a anode	b.p boiling point
a anterior	Br bromine
a abundant (occurrence of	B.T.U. British thermal unit
species) `	C carbon
a. adjective	C centigrade; Celsius
Å . Ångström unit	(' (centum) hundred; cen-
aapm. amphiapomict	tury
A.C., a.c alternating current	C corolla
ACTH . adrenocorticotrophic hor-	c (circa) approximately
mone	C.A chronological age
adv adverb	Ca calcium
act (aetatis) age(d)	ca cathode
Ag . silver	ca (circa) approximately
alt alternate	Cal large calorie(s)
alt altitude	cal small calorie(s)
Al aluminium	c.c cubic centimetre
Am ammonium	Cd cadmium
amph amphimict	Ce cerium
anal analysis	Cel Celsius
anat anatomical, anatomy	cel cellulose
ant anterior	cent hundred; centigrade
APH . anterior pituitary hormone	c.e.s central excitatory state
APL . anterior-pituitary-like hor-	c.i.s central inhibitory state
mone	cf compare
apm. , apomict	cg., cgm. centigram(s)
appl applied to	Cl chlorine
aq. water; aqueous	c.l corpus luteum
Ar. Arabic	cm centimetre(s)
A.S. Anglo-Saxon	c.mm cubic millimetre(s)
As arsenic	cm/s centimetres per second
atm . atmosphere, atmospheric	CO cardiac output
Au . gold	Co cobalt
A-V . atrioventricular	col., cols. (bacterial) colony, colonies
av average	conc concentrated, concentra-
Az . azote (nitrogen)	tion
B boron	conch. conchology
B Bacillus	C.O.V. cross-over value
Ba . barium	cpd compound
Bact bact Bacterium, bacterial	cpl carpel
bar. barometric	Cr chromium
Bi . bismuth	Cs caesium
B.I. buffer index	c.s.f cerebrospinal fluid
biol biological, biology	Cu copper
B.M.R. basal metabolic rate	cu., cub cubic
B.N.A. Basle nomina anatomica	D . deuterium
Dane nomina anatomica	a doubtient

d.	(dexter), right	G	gynoecium
d. d.	dextrorotary		gram(s)
d.	dominant (appl. species)	gal	gallon(s)
Dan.	Danish	γ,	(gamma) microgram
	decibel(s)	gen	genus
	direct current	geog	. geographical
deg.	degree(s)	geol	geological
deg. •	decigram(s)	Ger	German
diam.	diameter	Gk	Greek
dil	dilute	gm	. gram(s)
dim	diminutive	gmol.	. gram-molecule
INA	deoxyribonucleic acid	G.M.T.	. Greenwich Mean Time
	dozen	gp	
dr	drachm(s), dram(s)	or	. grain(s), gram(s)
Dut	Dutch	gr. n.	Gram-negative Gram-positive
dwt	pennyweight	gr. p.	. Gram-positive
E	east	GSH .	glutathione
E D	effective dose	gt., gtt.	(gutta) drop, (guttac) drops
E.D	efferens, efferent	Н .	. hydrogen
еп	(exempli gratia) for	Н°	hydrogen ion concentra-
e.g	example		tion
1	embryological,	Нь .	. haemoglobin
embr			helium
EME	embryology erythrocyte maturing		. herbarium
E.M.F.	factor		high-frequency
	electromotive force		. mercury
			hectogram(s)
ena	endosperm		. hæmoglobin
entom	entomological, entomology		. horizontal
eos	eosinophil		. hour, hours
equiv.	equivalent	H W	. High Water
E.R.S	erythrocyte sedimentation	hyb	bybrid
	rate	hypoth	hypothetical
esp	especially		
est	estimated, estimation	Î.	incisor iodine
η . , · ·	(eta) viscosity	ih ihid	(<i>ibidem</i>) in the same place
	(et alii) and others	Isol	Icolondia
F		icei	. Icelandic
F	Fahrenheit	ichth.	. ichthyology . interstitial cell-stimulating
F	r rench	ICSH	
F ₁ , F ₂ , etc	. 1st, 2nd, etc. filial genera-		hormone
	tion		. (id cst) that is
	. female		. inch(es)
f	. frequent (occurrence of		. inferior
	species)		inflorescence
	. focal distance		. (idem quod) the same as
	. iron		. iridescent
пr(s) .	. flower(s)	It	. Italian
fm .	. Iainom fraccing point	1.U	. international unit(s)
I.p	freezing point	INT D	. isotopic weight
ron.	follicle - stimulating hor-	ງ.N.D. ຮ	. just noticeable difference
	mone	К . К .	
	. foot; feet		. potassium
fth			. kathode
fur	. furlong	KC .	. kilocycles

kg kilogram(s)	μ (mu) micron (0.001 mm.)
km kilometre(s)	μ g microgram
λ · . (lambda) wave length	$\mu\mu$ inicromicron (0.001 m μ)
L Latin	mV millivolt(s)
L ligament(um)	myc mycology
L., Linn. Linnaean, Linnaeus	N nitrogen
l litre	N normal solution
l laevorotary	N north
Lam. Lamarck	n nasal
lat latitude	
IAL IAILUUC	n nervus n neutral, normal
lb pound (weight)	n neutral, normal
1 f low frequency LH luteinising hormone	n haploid number of
LH luteinising hormone	chromosomes
Li lithium	2n . diploid number
liq liquid; liquor	n refractive index
L.L Late Latin	n noun
log logarithm	Na sodium
longlongitude	Ne . neon
l.p low pressure	n.g new genus
L.S longitudinal section	Ni nickel
l.t low tension	No., no. , number
L.W Low Water	norm normal
M Membrana, Musculus	n.p normal pressure
M (mille) thousand	n.sp new species
• • •	NTP normal temperature and
m metre(s)	pressure O . oxygen
m mile(s)	
m minim(s)	o ortho-
m musculus	o occasional (occurrence of
m meta-	species)
ma milliampère	obl oblique, oblong
Mal Malaysian	O.F Old French
max maximum	O.H.G Old High German
mb millibar(s)	Ω (Omega) ohm(s)
M.E Middle English	opp as opposed to; opposite
micr microscopic	opt optical
min minimum; minute(s)	opt optimal
Mg magnesium	org organic
mg., mgm. milligram(s)	orig original
ml millilitre(s); c.c.	orn., ornith. ornithology
MLD . minimum lethal dose	Os osmium
mm millimetre(s)	O.T old terminology
	ov ovary
$m\mu$. millimicron (0.001 μ)	Oz ozone
Mn manganese	
Mo molybdenum	oz ounce(s)
mo month	P perianth
mol gram-molecule	P phosphorus
mol. wt molecular weight	P. premolar
m.p melting point	p posterior
m.p.h miles per hour	p para-
m.p.h miles per hour m.p.s metres per second	p.a per annum
msec millisecond ; σ	pal palaeontology
msec millisecond; σ m.s.l mean sea-level M.U mouse unit	P ₁ , P ₂ , etc. 1st, 2nd, etc., parental
M.U mouse unit	generation
	_

PATH	. pituitary adrenotrophic	ret retarded
	hormone	Rh rhesus factor
path	. pathology	rh anti Rh agglutinin
Pb .		R.I refractive index
Pe .	. probable error	Rn radon
DF	. precipitation: evaporation	RNA . ribonucleic acid
ΙμΕ.	ratio	rot rotation, rotating
202		R.Q respiratory quotient
-	. perennial	R.U rat unit
perp	. perpendicular	Russ. Russian
peri.	. pertaining to	
pH .	. hydrogen ion concentra-	S smooth (bacterial colony)
_	tion	S south
phys.		S sulphur
	. physiology	s (sinister), left
	. (\$\rho_1\$) 3.14159265	Σ (Sigma) sum of
pigm.	. pigment	σ $(sigma)$ 0.001 second; msec.
Pl	. plasma, Plasmodium	σ (sigma) standard deviation
blu	. plural	S-A sino-atrial
P-M-C	. pollen mother-cell	Sb antimony
pois		S.D.A specific dynamic action
	. population	Se selenium
pot.		sec second, secondary
P-P	. pellagra-preventing	Si silicon
nn .	. post partum	sin sinus
p.p.	. precipitating	sing singular
ppg	. parts per million	sl slightly
p.p.m.	. precipitate	sm small
ppt	partly soluble	S-M-C . sperm or spore mother-cell
		Sn tin
Pt .	• •	sol soluble; solution
pt	· pint	
pt	. point . pregnancy urine	Sp Spanish
P.O	. pregnancy urine	sp., spp species
p.u	. plant unit(s)	sp. gr specific gravity
pulv.	. (pulvis) powder	sq square
	. temperature coefficient	Sr strontium
	. quadrilateral	s.s sensu stricto
	. quintal	sta stamen(s)
	. quarter	sta station
qt	. quantity	std standard
qt	. quart	sup superior
q.v	. (quod vide) which see	Sw Swedish
R	. Réaumur	sym symmetrical
R.	. rough (bacterial colony)	syn synonym
r	. Roentgen unit	syst system
r .	. correlation coefficient	syst systole
r	. rare (appl. species)	t ^o temperature
Ra.	. rare (<i>appl</i> . species) . radium	T_b , t_b . body temperature
	. radius	temp temperature
	. radix	tert tertiary
RE .		t.g type genus
	. rectangular	Ti titanium
	. reflex	Tl thallium
reg	. regular	tot total
	reticulo-endothelial system	trop tropical
	. Italiano chiaothenai system	uop tropicar

TSH.	. thyroid-stimulating hor-	vert vertical
	mone	ves vesica, vesicle
TSP.	. thyroid-stimulating hor-	vet veterinary
	mone of prepituitary	vic vicinal
U	. unit(s)	visc viscous
	. uranium	vol. , volume
uns.	. unsymmetrical	v.s (vide supra) see above
	. United States	vs versus
	Pharmacopoeia	W tungsten
U V	. ultra-violet	W west
	. vanadium	w watt(s)
	. vibrio	wh white
	. vision	wk(s) $week(s)$
v .	. vision	w.l wave length, λ
	. volt(s)	wt weight
	. velocity	x . haploid generation
v		2x diploid generation
	. variable, variety	yd(s) $yard(s)$
ven.	. vena, vein	yr(s) $year(s)$
Vert.	. Vertebrata	Zn zinc
	. vertebra, vertebrate	zool zoological, zoology
	•	_

EQUIVALENTS

```
One acre = 4840 square yards = 4046.873 square metres
One ångström = 0.0001 micron
One are = 119.6033 square yards = 0.02471 acre
One atmosphere = 14.7 pounds per square inch = 1033 grams per square
    centimetre
One bar = pressure of mercury column of 29.53 inches, one square centi-
    metre in area, in latitude 45°
One British thermal unit = 251.99 calories
One centigram - 0.15432 grain
One centimetre = 0.39370 inch
One chain = 22 yards = 20.116 metres
One cubic centimetre = 0.061 cubic inch
One cubic foot = 0.0283 cubic metre
One cubic inch = 16.387 cubic centimetres
One cubic metre == 1.308 cubic yards
One degree centigrade = 0.8 degrees Réaumur = 1.8 degrees Fahrenheit
One degree Fahrenheit = 0.5556 Centigrade
One degree of latitude at the equator = 68.704 statute miles
One degree of longitude at the equator = 69.65 statute miles
One degree Réaumur = 1.25 degrees Centigrade
One drachm = 60 minims
One fathom = 6 feet = 1.828767 metres
One foot = 12 inches = 30.4801 centimetres
One furlong -- 10 chains = 201.164 metres
One gallon (Imperial) = 1.2 gallons (U.S.A.) = 4.54596 litres
One gallon (U.S.A.) 0.8327 gallon (Imperial) = 3.785 litres
One grain (avoirdupois) = 0.0647989 gram
One gram = 0.035274 ounce = 15.432356 grains
One gram-calorie = 0.003968 British thermal units
One hectare = 2.47106 acres
One hundredweight = 112 pounds = 50.80 kilograms
One inch = 2.53995 centimetres
One inch (U.S.A.) = 2.54001 centimetres
One kilogram = 2.20462 pounds (avoirdupois)
One kilogram per square centimetre = 14.22 pounds per square inch
One kilometre = 0.62137 statute mile
One litre = 1.76077 pint (Imperial)
One metre = 3.28084 feet = 39.37011 inches
One microgram = 0.001 milligram = 0.00015 grain
One micron = 0.001 millimetre = 0.000039 inch
One mil = 0.001 inch
One mile (nautical) = 1.152 statute miles = 1.8532 kilometres
One mile (statute) = 1.6003 kilometres
One millibar = 0.001 bar
One milligram = 0.015432 grain
One millilitre = 0.001 litre
One millimetre = 0.03937 inch
```

One millimicron = 0.000001 millimetre One minim (British) = 0.05919 cubic centimetre One minim (U.S.A.) = 0.00376 cubic inch = 0.06161 cubic centimetre One ounce (avoirdupois) = 437.5 grains = 28.34954 grams One ounce (troy) = $31 \cdot 1035$ grams One pennyweight == 1.555 grams One pint (Imperial) = 568.2454 cubic centimetres One pound (avoirdupois) = 16 ounces = 453.59243 grams One pound (Troy) = 5760 grains = 373.24 gramsOne pound per square inch = 70.308 grams per square centimetre One quart (Imperial) = $1 \cdot 13649$ litre One quart (U.S.A.) = 0.94636 litre One quintal = 100 kilograms = 220.4621 pounds One rood = 40 poles = 10.1168 aresOne square centimetre = 0.15498 square inch One square foot = 929.034 square centimetres One square inch = 6.4516 square centimetres One square kilometre = 0.3861 square mile One square metre = 1550 square inches One square mile = 640 acres = 2.58998 square kilometresOne square millimetre = 0.00155 square inch One square yard = 0.8361 square metre One stere=35.3156 cubic feet One stone = 6.3303 kilograms One yard = 0.9144 metre

Sound velocity (air), mean = 331.7 metres per second Zero, absolute = -459.4° F. = -273° C. = -218.4° R. Zero, centigrade, and Réaumur = $+32^{\circ}$ F. Zero, Fahrenheit = -17.78° C

A DICTIONARY OF

SCIENTIFIC TERMS

- abactinal (ăbăk'tinăl, ăbăktī'năl) a. [L. ab, from; Gk. aktis, ray.] Appl. area of echinoderm body without tube-feet and in which madreporite is usually included; abambulacral; antambulacral.
- abambulacral (ăb'ămbūlā'krăl) a. [L. ab, from; ambulare, to walk.] Abactinal, q.v.
- abapical (ăbāp'īkāl) a. [L. ab, from; apex, summit.] Pert. or situated at lower pole.
- abaxial (ābāk'siāl) a. [L. ab, from; axis, axle.] Pert. that surface of any structure which is remote or turned away from the axis; excentric. Opp. adaxial.
- abaxile (ǎbāk'sīl) a. [L. ab, from; axis, axle.] Appl. embryo whose axis has not the same direction as axis of seed.
- **abbreviated** (ăbrē'viātēd) a. [L. ad, to; brevis, short.] Shortened; curtailed.
- abdomen (ăbdō'men) n. [L. abdomen, belly.] The belly; in vertebrates, part of body containing digestive organs; in Arthropoda and certain Polychaeta, posterior part of body; in Synascidiae, part of zooid below thorax.
- abdominal (ăbdŏm'ĭnăl) a. [L. abdomen, belly.] Pert. abdomen; appl. structures, organs, or parts of organs situated in, on, or closely related to, the abdomen.
- abdominal pores,—single or paired openings leading from coelom to exterior, in cyclostomes and certain fishes.
- abdominal reflex,—contraction of abdominal wall muscles when skin over side of abdomen is stimulated.
- abdominal regions,—nine areas into which the abdomen is divided by

- two horizontal and two vertical imaginary lines, viz., hypochondriac (2), lumbar (2), inguinal (2), epigastric, umbilical, hypogastric.
- abdominal ribs,—ossifications occurring in fibrous tissue between skin and muscles of certain reptiles.
- abdominal ring,—one of two openings in fasciae of abdominal muscles through which passes spermatic cord in male, round ligament in female; inguinal ring.
- abducens (abdu'senz) n. [L. abducere, to lead away.] The sixth cranial nerve, supplying the rectus externus muscle of the eyeball.
- abduct (ăbdūkt') v. [L. abductus, led away.] To draw away from median axis.
- abductor (ăbdūk'tŏr) n. [L. abductus, led away.] A muscle that draws a limb or part outwards.
- aberrant (ăběr'ánt) a. [L. aberrare, to stray.] With characteristics not in accordance with type; appl. species, etc.
- abience (ăb'iens) n. [L. abire, to depart.] Retraction from stimulus; avoiding reaction. Opp. adience.
- abient (äb'ient) a. [L. abire, to depart.] Avoiding the source of stimulation. Opp. adient.
- abiogenesis (ăbiöjen'esis) n. [Gk. a, not; bios, life; genesis, birth.] The production of living from non-living matter; spontaneous generation. Opp. biogenesis.
- abiology (abiol'oji) n. [Gk. a, not; bios, life; logos, discourse.] The study of non-living things.
- abjunction (ăbjungk'shun) n. [L. abjungere, to unyoke.] The delimitation of spores by septa at tip of hypha.

ablactation (ăb'lăktā'shūn) n. [L. ab, from; lactare, to give milk.] Cessation of milk secretion; weaning.

abomasum (ăbomā'sum) n. [L. ab, from; omasum, paunch.] The read or fourth chamber of stomach of

ruminants.

ABL-

aboral (abō'ral) a. [L. ab, from; os, oris, mouth.] Away from, or opposite to, the mouth.

abortion (ăbôr'shun) n. [L. abortus, premature birth.] Premature birth; arrest of development of an organ.

abranchiate (åbrang'kiāt) a. [Gk. a, without; brangchia, gills.] Without gills.

abrupt (abrupt') a. [L. abrumpere, to break off.] Appearing as if broken, or cut off, at extremity.

abruptly - acuminate, — having a broad extremity from which a point arises, appl. leaf.

abruptly-pinnate,—having the main axis of epipodium not winged, but bearing a number of secondary axes which are winged.

abscise (absiz') v. [L. abscidere, to cut off.] To become separated; to fall off, as leaves, fruit, etc.

- absciss (ăb'sīs) a. [L. abscindere, to cut off.] Appl. layer of meristematic cells just outside cork-layer, to whom fall of leaves, floral parts, fruits, and certain branches is due; abscissile.
- abscission (absish'un) n. [L. abscindere, to cut off.] The separation of
- absorption (ăbsôrp'shun) n. [L. absorbere, to suck in.] Intussusception of fluid by living cells or tissues; passage of nutritive material through living cells.
- abstriction (abstrik'shun) n. [L. abstringere, to cut off.] The process of detaching spores (conidia) by rounding off of tips of sporophores, as in mildews; abjunction and abscission.
- abterminal (ăbter'minăl) a. [L. ab, from; terminus, end.] Going from the end inwards.

abyssal (ăbĭs'ăl) a. [Gk. abyssos, unfathomed.] Pert. depths of ocean; appl. organisms or material usually found there; abysmal.

abyssobenthic (ābīs'öběn'thīk) a. Gk, abyssos, unfathomed; benthos, depths of sea.] Pert., or found on, bottom of ocean at depths exceeding

ca. 1000 metres.

abyssopelagic (ăbis'öpčlăj'ik) a. [Gk. abyssos, unfathomed; pelagos, sea.] Pert., or inhabiting, the ocean at depths exceeding ca. 1000 metres, i.e., below the bathypelagic zone.

acanaceous (ăkănā'seŭs) a. [Gk. akanos, thistle.] Prickly; bearing prickles, as leaves.

acantha (ǎkǎn'thǎ) n. [Gk. akantha, thorn.] Prickle; spinous process.

acanthaceous (ăkănthā'seŭs) a. [Gk. akantha, thorn.] Bearing thorns or prickles.

acanthin (ăkăn'thĭn) n. [Gk. akantha, thorn.] Substance forming skeleton of some Radiolaria.

- acanthion (äkän'thiŏn) n. [Gk. akanthion, small thorn.] The most prominent point on the anterior nasal spine.
- acanthocarpous (åkän'thökâr'pūs) a. [Gk. akantha, thorn; karpos, fruit.] Having fruit covered with spines or prickles.
- acanthocephalous (äkăn'thökĕf'ălŭs, -sĕf-) a. [Gk. akantha, thorn; kephale, head.] With hooked proboscis.
- acanthocladous (ǎkǎn'thöklād'ǔs) a. [Gk. akantha, thorn; klados, branch.] Having spiny branches.
- acanthocyst (ăkăn'thösist) n. [Gk. akantha, thorn; kystis, bladder.] A sac containing lateral or reserve stylets in Nemertea.
- acanthoid (ākān'thoid) a. [Gk. akantha, thorn; eidos, shape.] Resembling a spine or prickle; spiniform.
- acanthophore (ăkăn'thöför) n. [Gk. akantha, thorn; pherein, to bear.]
 A conical mass, the basis of median stylet in Nemertea.

- acanthopore (ākān'thöpōr) n. [Gk. akantha, thorn; poros, passage.]
 A tubular spine in certain Polyzoa.
- acanthosphenote (ākān'thösfē'nōt) a. [Gk. akantha, thorn; sphen, wedge.] Appl. echinoid spine made of solid wedges separated by porous tissue.
- acanthozooid (ἄkǎn'thözō'oid) n. [Gk. akantha, thorn; zoon, animal; eidos, form.] Tail part of proscolex of cestodes. Ορφ. cystozooid.

acapnia (ākāp'niā) n. [Gk. akapnos, without smoke.] Condition of low carbon dioxide content in blood

acaryote,—akaryote.

- acaulescent (ăkôlěs'ënt) a. [Gk. a, without; kaulos, stalk.] Having a shortened stem.
- acauline (åkô'līn) a. [Gk. a, without; kaulos, stalk.] Having no stem; appl. certain fungi.
- accelerator (ăksĕl'ĕrātŏr) n. [L. accelerare, to hasten.] Appl. muscle or nerve which increases rate of action.
- acceptor (ăksĕp'tŏr) n. [L. accipere, to accept.] Body or substance which receives and unites with another substance, as in oxidation-reduction processes where oxygen acceptor is the substance oxidised, hydrogen acceptor the substance reduced.
- accessorius (āksēsō'rīŭs) n. [L. accedere, to go to.] A muscle aiding in action of another; spinal accessory or eleventh cranial nerve.
- accessory bodies,—minute argyrophil particles originating from Golgi substance in spermatocytes; chromatoid bodies.
- accessory bud,—an additional axillary bud; a bud formed on a leaf. accessory cells,—see auxiliary cells. accessory chromosomes,—sex chromosomes.
- accessory food factors,—vitamins.
 accessory glands,—detached portions of glands; glands in relation with genital ducts.
- accessory pulsatory organs,—saclike structures of insects, variously situated, pulsating independently.
- acclimation (ăk'līmā'shun) n. [L. ad, to; clima, climate.] The

- habituation of an organism to a different climate or environment.
- acclimatisation (āklī'mātīzā'shūn) n. [L. ad. to; clima. climate.] Habituation of a species to a different climate or environment; acclimation under human management; acclimatation.
- accommodation (äk'ömödā'shūn) n. [L. ad, to; commodus, fitting.] Adjustment of eye to receive clear images of different objects; adaptation of receptors to a different stimulus.
- accrescent (ăkrēs'ēnt) a. [L. accrescere, to increase.] Appl. plants that continue to grow after flowering, or calyx growing after pollination.
- accretion (ăkrē'shūn) n. [L. accrescere, to increase.] Growth by external addition of new matter.
- accumbent (ākum'bent) a. [L. accumbere, to lie on.] Appl. embryo having cotyledons with edges turned towards radicle, as in Cruciferae.
- accumulators (ăkū'mūlātŏrz) n. plu. [L. ad, to; cumulus, heap.] Plants with a relatively high concentration of certain chemical elements in tissues.
- A-cells,—alpha cells of islets of Langerhans.
- acentric (ăsĕn'trĭk) a. [Gk. a, without; kentron, centre.] Having no centromere, appl. chromosomes and chromosome segments.
- acentrous (ăsen'trus) a. [L. a, without; centrum, centre.] With no vertebral centra, but persistent notochord, as certain fishes.
- acephalocyst (åkef'álösist, -sef-) n.
 [Gk. a, without; kephale, head; kystis, bladder.] Hydatid stage of certain tapeworms.
- acephalous (åkěf'ălūs, -sěf-) a. [Gk. a, not; kephale, head.] Having no structure comparable to head; appl. some molluscs; appl. larvae of certain Diptera; appl. ovary without terminal stigma.
- acerate (ăs'ērāt) a. [L. acer, sharp.]
 Needle-shaped; pointed at one
 end, appl. monaxon or oxeote
 spicules.

acerose (ăs'ërōs) a. [L. acer, sharp.] Narrow and slender, with sharp point, as leaf of pine.

acervuline (ăser'vulin) a. [L.L. dim. of acervus, heap.] Irregularly heaped together, appl. foraminiferal tests.

acervulus (ăsĕr'vūlŭs) n. [L.L. dim. of acervus, heap.] A small heap or cluster, especially of sporogenous mycelium.

acervulus cerebri,—brain sand, q.v. acetabular,—pert. acetabulum.

acetabulum (ăsĕtăb'ūlŭm) n. [L. acetabulum, vinegar-cup.] The cotyloid cavity or socket in pelvic girdle for head of femur; in insects, cavity of thorax in which leg is inserted; cavity in proximal end of spine, for articulation with mamelon, in echinoids; large posterior sucker in leeches; sucker on arm of cephalopod; one of the cotyledons of placenta in ruminants.

achaetous (ăkē'tŭs) a. [Gk. a, without; chaite, hair.] Without chaetae or bristles.

acheilary (ākī'lārī) a. [Gk. a, without; cheilos, lip.] Having labellum undeveloped, as some orchids.

achene (ākēn') n. [Gk. a, not; chainein, to gape.] A one-seeded, dry, indehiscent fruit; achenium.

Achillis tendo (ăkīl'īs tēn'dō) n. [Gk. Achilles; L. tendo, tendon.] The united strong tendon of gastrocnemius and solacus muscles, tendo calcaneus.

achlamydeous (åk'lămĭd'ëŭs) a. [Gk. a, without; chlamys, cloak.] Having neither calyx nor corolla.

achondroplasia (ākòn'dröplā'zĭā) n. [Gk. a, without; chondros, cartilage; plasis, a moulding.] Heritable dwarfism due to disturbance of ossification in the long bones of the limbs and of certain facial bones during development; cf. ateleosis.

achroacyte (ăkrō'ăsīt) n. [Gk. a. not; chros, colour; kytos, hollow.] Colourless or lymph cell; lymphocyte.

achroglobin (āk'rōglō'bĭn) n. [Gk. a, not; chros, colour; globos,

sphere.] A colourless respiratory pigment of some tunicates and molluscs

achromasie (ăkrō'māsī) n. [Gk. a, not; chroma, colour.] Emission of chromatin from nucleus; cf. chromasie.

achromatic (āk'rōmāt'ik) a. [Gk. a, without; chroma, colour.] Appl. threshold, the minimal stimulus inducing sensation of luminosity or brightness; cf. chromatic; appl. neutral colours; achromatinic, q.v.

achromatin (ākrō'mātīn) n. [Gk. a, without; chroma, colour.] The non-staining ground substance and linin of the nucleus.

achromatinic (ākrō'mātīn'īk) a. [Gk. a, without; chroma, colour.] Pert. achromatin, or resembling achromatin in properties.

achromite,—centromere, q.v.

achroous (ăkrō'ūs) a. [Gk. a, without; chros, complexion.] Unpigmented; colourless.

acicle (ăs'ikl) n. [L. acicula, small needle.] A thorn-shaped scaphocerite, as in Paguridae; acicula, a small spine.

acicular (ăsīk'ŭlăr) a. [L. acicula, small needle.] Like a needle in shape; sharp-pointed.

aciculum (ăsĭk'ūlŭm) n. [L. acicula, small needle.] A stiff basal seta in parapodium of Chaetopoda.

acid-fast,—remaining stained with aniline dyes on treatment with acids; appl. bacteria.

acidic (ăsid'ik) a. [L. acidus, sour.]

Appl. stains whose colour determinant plays the part of an acid, acting on protoplasm; cf. basic.

acidophil (as'idofil) a. [L. acidus, sour; Gk. philein, to love.] Oxyphil, q.v.; growing in acid media.

aciform (ăs'ifôrm) a. [L. acus, needle; forma, shape.] Needle-shaped.

acinaciform (ăsĭn'ăsĭfôrm) a. [L. acinaces, short sword; forma, shape.] Shaped like a sabre or scimitar; appl. leaf.

acinarious (ăsĭnā'rĭŭs) a. [L. acinus, berry.] Having globose vesicles, as some algae. aciniform (ăsĭn'ĭfôrm) a. [L. acinus, berry; forma, shape.] Grape- or berry-shaped; appl. a type of silk gland in spiders.

acinus (ăs'ĭnŭs) n. [L. acinus, berry.]
One of drupels composing fruit of bramble or raspberry; sac-like termination of branched gland.

acme (ăk'mē) n. [Gk. akme, prime.]
The highest point attained, or prime, in phylogeny and ontogeny;
cf. epacme, paracme.

tj. epacine, paracine.

acoelomate (ăsē'lömāt) a. [Gk. a, without; koilos, hollow.] Appl. animals not having a true coelom; acoelomatous; acoelous, q.v.

- acoelous (ăsē'lŭs) a. [Gk. a, without; koilos, hollow.] Appl. vertebrae with flattened centra; acoelomate, q.v.
- acondylous (ākŏn'dilŭs) a. [Gk. a, without; kondylos, knuckle.] Without nodes or joints.
- acone (ăkôn') a. [Gk. a, without; konos, cone.] Appl. insect compound eye without crystalline or liquid secretion in cone cells.
- acontia (ākŏn'tiă, ākŏn'shiă) n. plu. [Gk. akontion, small javelin.] Threadlike processes of mesenteric filaments armed with stinging cells, in actinians.
- acotyledon (ā'kŏtīlē'dŏn) n. [Gk. a, without; kotyledon, a cup-shaped hollow.] A plant without a cotyledon.
- acoustic (ăkoo'stĭk) a. [Gk. akouein, to hear.] Pert. organs or sense of hearing; appl. meatus, nerve, ctc.; pert. science of sound.
- acquired character,—a modification or permanent structural or functional change effected during the lifetime of the individual organism and induced by use or disuse of a particular organ, by disease, trauma, or other functional or environmental influences.
- acraspedote (ăkrăs'pēdōt) a. [Gk. a, without; kraspedon, border.]
 Having no velum.
- acroblast (ăk'röblăst) n. [Gk. akros, tip; blastos, bud.] A body in spermatid and which gives rise to acrosome; outer layer of mesoblast.

acrobryous (ākröb'riŭs) a. [Gk. akros, tip; bryein, to swell.] Growing at the tip only.

acrocarpic (åk'rökâr'pīk) a. [Gk. akros, tip; karpos, fruit.] Having the fructification terminating the axis; acrocarpous; appl. mosses.

- acrocentric (āk'rösĕn'trīk) a. [Gk. akros, tip; kentron, centre.] With centromere at end, appl. chromosome. n. A rod-shaped chromosome.
- acrochordal (äkrökôr'dăl) a. [Gk. akros, tip; chorde, cord.] Appl. a chondrocranial unpaired frontal cartilage in birds.
- acrocoracoid (ăk'rökŏr'ăkoid) n... [Gk. akros, tip; korax, crow; eidos, form.] A process at dorsal end of coracoid in birds.
- acrocyst (ăk'rösist) n. [Gk. akros, tip; kystis, bladder.] The spherical gelatinous cyst formed by gonophores at maturation of generative cells.
- acrodont (āk'rödönt) a. [Gk. akros, tip; odous, tooth.] Appl. teeth attached to the summit of a parapet of bone, as in lizards.
- acrodrome (ăk'rödrōm), acrodromous (ăkrŏd'römŭs) a. [Gk. akros, tip; dromein, to run.] Appl. leaf with veins converging at its point.
- acrogenous (ăkroj enus) a. [Gk. akros, tip; -genes, producing.] Increasing in growth at summit or apex.
- acrogynous (åkrŏj'inŭs) a. [Gk. akros, tip; gyne, female.] With archegonia arising from apical cell, appl. certain liverworts. Opp. anacrogynous.
- acromegaly (ăk'römëg'ăli) n. [Gk. akros, tip; megalon, great] Gigantism due to excessive activity of part of pituitary gland.
- acromial (ăkrō'miăl) a. [Gk. akros. summit; omos, shoulder.] Pert. acromion, appl. artery, process, ligament, etc.
- acromio-clavicular (ākrō'miöklāvīk'ūlār) a. [Gk. akromion, shouldersummit; L. claviculus, dim. of
 clavis, key.] Appl. ligaments covering joint between acromion and
 clavicle.

- acromion (ăkrō'miŏn) n. [Gk. akros, summit; omos, shoulder.] Ventral prolongation of scapular spine.
- acron (ăk'ron) n. [Gk. akros, tip.] Preoral region of insects; anterior, unsegmented part of young trilobite.
- acropetal (ākrōp'ētāl) a. [Gk. akros, summit; L. petere, to seek.] Ascending; appl. leaves, flowers, or roots, developing successively from an axis so that youngest arise at apex.
- acrophyte (ăk'röfit) n. [Gk. akron, peak; phyton, plant.] A plant growing at a high altitude; alpine plant.
- acropodium (āk'röpōdium) n. [Gk. akros, tip; pous, foot.] Digits,—fingers or toes; cf. metapodium.
- acrorhagus (ăkrörā'gūs) n. [Gk. akros, summit; rhax, grape.] A tubercle near the margin of certain Actiniaria, containing specialised nematocysts.
- acrosarc (āk'rösârk) n. [Gk. akros, summit; sarx, flesh.] A pulpy berry resulting from union of ovary and calyx.
- acroscopic (åk'rösköp'ik) a. [Gk. akros, tip; skopein, to view.] Facing towards the apex, opp. basiscopic.
- acrosome (åk'rosom) n. [Gk. akros, tip; soma, body.] Body at apex of spermatozoon; apical body; perforatorium.
- acrospire (ăk'röspīr) n. [Gk. akros, tip; speira, something twisted.] The first shoot or sprout, being spiral, at end of germinating seed.
- acrospore (åk'röspōr) n. [Gk. akros, tip; sporos. seed.] The spore at end of a sporophore.
- acroteric (ăk'rötĕ'rĭk) a. [Gk. akroterion. topmost point.] Pert. outermost points, as tips of digits, nose, ears, tail.
- acrotonic (ãk'rötŏn'īk) a. [Gk. akros, tip; tonos, brace.] Having anther united at its apex with rostellum; acrotonous. Opp. basitonic.
- acrotrophic (ăk rötrof'ik) a. [Gk. akros, tip; trophe, nourishment.]

 Appl. ovariole having nutritive cells at apex which are joined to

oocytes by nutritive cords; telotrophic.

actinal (ăk'tĭnăl, ăktī'năl) a. [Gk. aktis, ray.] Appl. area of echinoderm body with tube-feet; appl. oral area with tentacles in Actiniaria. actine (ăk'tĭn) n. [Gk. aktis, ray.] A

star-shaped spicule.

actinenchyma (āk'tiněng'kimā) n. [Gk. aktis, ray; en, in; chein, to pour.] Cellular tissue having a stellate appearance.

actiniform,—actinoid.

- actinobiology (āk'tīnöbīðl'öji) n.
 [Gk. aktis, ray; bios, life; logos, discourse.] The study of the effects of radiation upon living organisms.
- actinoblast (ăk'tinöblăst) n. [Gk. aktis, ray; blastos, bud.] The mother-cell from which a spicule is developed, as in Porifera.
- actinocarpous (āk'tīnökâr'pŭs) a. [Gk. aktis, ray; karpos, fruit.] Appl. plants with flowers and fruit radially arranged; actinocarpic.
- actinodrome (åktīn'odrōm) a. [Gk. aktis, ray; dromos, course.] Veined palmately; actinodromous.
- actinogonidial (ăk'tinögonid'iăl) a. [Gk. aktis, ray; gonos, offspring.] Having radiately arranged genital organs.
- actinoid (ăk'tinoid) a. [Gk. aktis, ray; eidos, shape.] Rayed; starshaped, stellate.
- actinomere (ăktin'ömēr) n. [Gk. aktis, ray; meros, part.] A radial segment.
- actinomorphic (ăk'tinömôr'fik) a. [Gk. aktis, ray; morphe, shape.] Radially symmetrical; actinomorphous.
- actinopharynx (ăk'tinöfār'ingks) n. [Gk. aktis, ray; pharynx, gullet.] The gullet of a sea-anemone.
- actinost (ăk'tīnost) n. [Gk. aktis, ray; osteon, bone.] Basal bone of fin-rays in teleosts.
- actinostele (ăk'tinöstē'lē) n. [Gk. aktis, ray; stele, pillar.] Stele with xylem radiating outwards and forming ridges, as in certain Pteridophyta.

actinostome (äktin'östöm) n. [Gk. aktis, ray; stoma, mouth.] The mouth of a sea-anemone; five-rayed oral aperture of starfish.

actinotrichia (ăk'tinötrik'iă) n. plu. [Gk. aktis, ray; thrix, hair.] Unjointed horny rays at edge of

fins in many fishes.

actinotrocha (åk'tinöt'rökå) n. [Gk. aktis, ray; trochos, wheel.] Free-swimming larval form of Phoronis.

actinula (ăktin'ūlă) n. [Gk. aktis, ray.] A larval stage in some Hydro-

medusae.

action system,—behaviour pattern. activator (āktīvā'tŏr) n. [L. activus, active.] A substance which promotes or protects enzyme action; a substance which stimulates development of any particular embryonic tissue or organ.

aculeate (ăkū'lēāt) a. [L. aculeus, prickle.] Having prickles, sharp

points, or a sting.

aculeiform (ăkū'leifôrm) a. [L. aculeus, prickle; forma, shape.] Formed like a prickle or thorn.

- aculeus (ǎkū'lēūs) n. [L. aculeus, prickle.] A prickle growing from bark, as in rose; a sting; a hairlike projection; a microtrichium.
- acuminate (äkū'mināt) a. [L. acumen, point.] Drawn out into long point; tapering; pointed.

acuminiferous (ăkū'mĭnĭf'ërŭs) a. [L. acumen, point; ferre, to carry.]

Having pointed tubercles.

acuminulate (åk'ūmĭn'ūlāt) a. [L. acuminulus, dim. of acumen, point.]
Having a very sharp tapering point.

acute (ăkūt) a. [L. acutus, sharpened.]
Ending in a sharp point; temporarily severe, not chronic.

- acyclic (åsik'lik) a. [Gk. a, without; kyklos, circle.] Appl. flowers with floral leaves arranged in a spiral.
- adamantoblast (ăd'ămăn'töblăst) n.
 [Gk. adamas, diamond; blastos,
 bud.] Enamel cell; ameloblast.
- adambulacral (ăd'ămbūlā'krăl) a.
 [L. ad, to; ambulare, to walk.]
 Appl. structures adjacent to ambulacral areas in echinoderms.

adaptation (ădăptā'shūn) n. [I. ad, to; aptare, to fit.] The process by which an organism becomes fitted to its environment; a structure or habit fitted for some special environment; the fitting of sensations to a point when discomfort ceases; adjustment of disturbance of nervous system without involving higher co-ordinating centres.

adaxial (ădăk'siăl) a. [L. ad, to; axis, axle.] Turned towards the

axis. Opp. abaxial.

ad-digital (ād'dīj'ītāl) n. [L. ad, to; digitus, finger.] A primary wingquill connected with phalanx of third digit.

adduction (ad'duk'shun) n. [L. ad, to; ducere, to lead.] Movement

towards the median axis.

adductor (åd'důk'tŏr) n. [L. ad, to; ducere, to lead.] A muscle which brings one part towards another.

adeciduate (å'dēsid'ūāt) a. [L. a, away from; decidere, to fall down.]
Not falling, or coming away; appl. evergreens; appl. placenta.

- adelocodonic (ádē'lökōdōn'ik) a.
 [Gk. adelos, concealed; kodon, bell.] Appl. undetached medusome of certain Gymnoblastea, which degenerates after discharging ripe sexual cells. Opp. phanerocodonic.
- adelomorphic (ădê'lömôr'fik) a. [Gk. adelos, concealed; morphe, shape.] Indefinite in form; appl. central cells of peptic glands; adelomorphous.
- adelophycean (ădel'löfisē'ăn) a. [Gk. adelos, concealed; phykion, seaweed.] Appl. stage or generation of many seaweeds when they appear as prostrate microthalli.

adelphogamy (ădēlfög'āmi) n. [Gk. adelphos, brother; gamos, marriage.] Brother-sister mating, as in certain ants.

adelphous (ăděl'fűs) a. [Gk. adelphos, brother.] Joined together in bundles, as filaments of stamens; cf. monadelphous, diadelphous.

adendritic (ă'děndrit'īk) a. [Gk. a, not; dendron, tree.] Adendric; without dendrites or branches; appl. cells.

- adendroglia (ăděndrógli'á) n. [Gk. a, not; dendron, tree; gloia, glue.] A type of neuroglia lacking processes.
- adenine (ăd'ënĭn) n. [Gk. aden, gland.] A compound occurring in many cells, hydrolysed by adenase to hypoxanthine; $C_bH_bN_b$.

adenoblast (ăd'enöblăst) n. [Gk. aden, gland; blastos, bud.] Em-

bryonic glandular cell.

- adenocheiri (ăd'enoki'rī) n. plu. [Gk. aden, gland; cheir, hand.] Elaborate accessory copulatory organs, outgrowths of atrial walls in Turbellaria.
- adenocyte (ad'enosit) n. [Gk. aden, gland; kytos, hollow.] Secretory cell of a gland.
- adenodactyli (åd'ënodåk'tili), n. plu.
 [Gk. aden, gland; daktylos, finger.]
 Adenocheiri, q.v.
- adenohypophysis (ăd'ënöhīpŏf'isis)
 n. [Gk. aden, gland; hypo, under;
 physis, growth.] The glandular lobe
 or portions of the pituitary body,
 derived from Rathke's pouch.
- adenoid (ăd'ënoid) a. [Gk. aden, gland; eidos, shape.] Pert. or resembling a gland or lymphoid tissue.
- adenophore (ăd'ënöfor') n. [Gk. aden, gland; pherein, to carry.]
 The stalk of a nectar gland.
- adenophyllous (ăd'ënöfil'ŭs) a. [Gk. aden, gland; phyllon, leaf.] Bearing glands on leaves.
- adenopodous (ădenop'odus) a. [Gk. aden, gland; pous, foot.] Bearing glands on peduncles or petioles.

adenose (ăd'ënōs) a. [Gk. aden, gland.] Glandular.

gland.] Glandular. adenostemonous (ăd'ë

- adenostemonous (ăd'ēnöstēm'onŭs)
 a. [Gk. aden, gland; stemon, spun thread.] Having glands on stamens.
- adequate,—appl. stimulus which normally acts on a given receptor, and induces the appropriate sensation.
- adermin, vitamin B₆, rat antidermatitis factor; pyridoxine.
- adesmic (ădēs'mik) a. [Gk. adesmos, unfettered.] Appl. cyclomorial scales made up of separate lepido-

- morial units; *cf.* monodesmic, polydesmic.
- adesmy (ăděs'mi) n. [Gk. adesmos, unfettered.] A break or division in an organ usually entire.
- adetopneustic (ād'ētŏnū'stĭk) a. [Gk. adetos, free; pnein, to breathe.] Having dermal gills occurring beyond abactinal surface, as in certain stelleroids.
- adfrontal (ădfrun'tăl) a. [L. ad, to; frons, forehead.] Appl. oblique plates beside frons of certain insect larvae.
- adherent (ădhē'rēnt) a. [L. ad, to; haerere, to stick.] Exhibiting adhesion, q.v.; attached to substratum, appl. zooecia of polyzoan colony.
- adhesion (ādhē'zhūn) n. [L. ad, to; haerere, to stick.] Condition of touching without growing together of parts normally separate, as between members of different series of floral leaves; cf. cohesion.
- adhesive cells,—various glandular or specialised cells for purposes of attachment; as on tentacles of Ctenophora, on epidermis of Turbellaria, on pedal disc of Hydra.
- adience (ăd'iëns) n. [L. adire, to approach.] Urge, or advance, towards stimulus; approaching reaction. Opp. abience.
- adient (ăd'iënt) a. [I.. adire, to approach.] Approaching the source of stimulation. Opp. abient.
- adipocellulose (ăd'ipösĕl'ūlōs) n. [L. adeps, fat; cellula, small cell.] Cellulose with a large amount of suberin, as in cork tissue.
- adipocyte (ăd'ípösīt) n. [L. adeps, fat; Gk. kytos, hollow.] One of the cells forming the fat-body in insects.
- adipoleucocyte (ăd'îpölū'kösīt, -loo-) n. [L. adeps, fat; Gk. leukos, white; kytos, hollow.] A leucocyte containing fat droplets or wax, in insects.
- adipose (ad'ipos) a. [L. adeps, fat.]

 Pert. animal fat; fatty.
- **A-disc,**—doubly refracting or anisotropic band in myofibrillae.

- aditus (ăd'itŭs) n. [L. aditus, entrance,] Anatomical structure forming approach or entrance to a part, e.g. to antrum, larynx, etc.
- adjustor (ădjūs'tor) n. [L.L. adjustare, to adjust, from L. ad, to; justus, just.] A muscle connecting stalk and valve in Brachiopoda; ganglionic part of a reflex arc, connecting receptor and effector.
- adlacrimal (ădlăk'rimăl) n. [L. ad, to; lacrima, tear.] Lacrimal bone of reptiles.
- adminiculum (ăd'minik'ūlum) n. [L. adminiculum, support.] A locomotory spine of certain pupae; posterior fibres of linea alba attached to os pubis.
- adnasal (ădnā'zăl) n. [L. ad, to; nasus, nose.] A small bone in front of each nasal in certain fishes.
- adnate (ădnāt') a. [L. ad, to; gnatus, born.] Pert. or designating the condition of being closely attached to side of petiole or stalk, as stipules or leaves; designating condition of anther with back attached throughout its length to filament, or to its continuation the connective; conjoined.
- adnexa (ădněk'sá) n. plu. [L. ad, to; nectere, to bind.] Structures or parts closely related to an organ; extra-embryonic structures, as foetal membranes, placenta.

adnexed (ădněksť) a. [L. ad, to; nectere, to bind.] Reaching to the stem only.

- adolescaría (ădöleskā'rīā) n. [L. adolescere, to grow up.] Encysted stage, between cercaria and marita, in trematodes; metacercaria.
- adoral (ădō'răl) a. [L. ad, to; os, mouth.] Near or pert. mouth.
- adpressed (ăd'prēst) a. [L. ad, to; pressus, pressed.] Closely applied to a surface.
- adradius (ădrā'dĭŭs) n. [L. ad, to; radius, radius.] In coelenterates, the radius midway between perradius and interradius, a radius of third order.
- adrectal (ădrěk'tăl) a. [L. ad, to;

- rectum, rectum.] Near to or closely connected with rectum.
- adrenal (ădrē'năl) a. [L. ad, to; renes, kidneys.] Situated near kidneys; suprarenal, appl. glands, the endocrines secreting hormones affecting sympathetic nervous system and blood pressure; appl. organs, the suprarenal and interrenal glands, and chromaphil bodies.
- adrenaline (ādrē'nālīn, ādrē'nālēn) n. [L. ad, to; renes, kidneys.] A hormone obtained from extract of suprarenal medulla; adrenalin, adrenin, suprarenin, epinephrin; C₉H₁₈O₈N.
- adrenergic (ădrēněr'jīk) a. [L. ad, to; renes, kidneys; Gk. ergon, work.] Appl. sympathetic nerves, which liberate sympathin or an adrenaline-like principle from their terminations; adrenalnergic; cf. cholinergic.
- adrenocortical (ădrē'nökôr'tĭkăl) a. [I.. ad, to; renes, kidneys; cortex, bark.] Pert., or secreted in, the adrenal cortex.
- adrenocorticotrophic (ădre'nökôr'-tikötrŏf'ik) a. [L. ad, to; renes, kidneys; cortex, bark; Gk. trophe, nourishment.] Appl. hormone secreted by anterior lobe of pituitary gland and which controls activity of adrenal cortex; ACTH.
- adrenotropic (ädrēnötröp'ik) a. [L. ad, to; renes, kidneys; Gk. trope, turn.] Adrenotrophic; appl. a pituitary hormone acting on the adrenal medulla.
- adrostral (ădrös'trăl) a. [L. ad, to; rostrum, beak.] Near to or closely connected with beak or rostrum.
- adsorption (ădsôrp'shun) n. [L. ad, to; sorbere, to suck in.] The adhesion of molecules to solid bodies; formation of unimolecular surface layer.
- adtidal (åd'tīdǎl) a. [L. ad, to; A.S. tid, time.] Appl. organisms living just below low-tide mark.
- adultoid,—appl. nymph having imaginal characters differentiated further than in normal nymph.

aduncate (ădung'kāt) a. [L. aduncus, hooked.] Crooked; bent in the form of a hook.

advehent (ad'vëhënt) a. [L. advehere, to carry to.] Afferent; carry-

ing to an organ.

ADU-

adventitia (ădventish'iă) n. [L. adventitius, extraordinary.] External connective tissue layer of blood vessels; tunica adventitia.

adventitious (ādvēntīsh'ūs) a. [L. adventitius, extraordinary.] Accidental; found in an unusual place; appl. tissues and organs arising in abnormal positions; secondary, appl. dentine.

aecial,-aecidial.

aecidia,-plu. of aecidium.

aecidial (ēsīd'iăl) a. |L. aecidium, cup.] Pert. aecidia, or aecidium; appl. primordium.

aecidiosorus (ēsĭd'iösō'rŭs) n. [L. aecidium, cup; Gk. soros, heap.]
A cluster or row of aecidiospores.

aecidiospores (ēsīd'iöspōrz') n. plu. [L. aecidium, cup; Gk. sporos, seed.] The spores produced in an aecidium; spring-spores.

aecidium (ēsīd'ſūm) n. [L. aecidium, cup.] A cup-shaped structure containing simple sporophores, bearing rows of spores; cluster-cup of rusts.

aeciospores,—aecidiospores, q.v.

aecium,—aecidium, q.v.

aedeagus (ēdē'agūs) n. [Gk. aidoia, genitals.] The male intromittent organ of insects; also aedoeagus.

aegithognathous (ē'jīthŏg'nāthūs) a. [Gk. aegithos, hedge-sparrow; gnathos, jaw.] With maxillo-palatines separate, vomers forming a wedge in front and diverging behind; appl. a type of palate found in Passeres.

aeolian (ēōl'ĭăn) a. [L. Aeolus, god of the winds.] Wind-borne; appl.

deposits.

aerenchyma (āërēng'kimă) n. [Gk. aer, air; engchyma, infusion.] Tissue between spore mass and capsulewall in mosses; cortex of submerged roots of certain swamp plants; aerating cortical tissue in floating portions of some aquatic plants.

aerial (āē'rlal) a. [L. aer, air.] In-

habiting the air; appl. roots growing above ground, e.g. from stems of ivy, for purposes of climbing; also to small bulbs appearing in leaf-axils.

aerobe (ā'érōb) n. [Gk. aer, air; bios, life.] An aerobic organism, capable of living in presence of oxygen. Opp. anaerobe.

aerobic (ā'ërôb'ik) a. [Gk. aer, air; bios, life.] Thriving only in pres-

ence of oxygen.

aerobiology (ā'ēröbīôl'öjī) n. [Gk. aer, air; bios, life; logos, discourse.] The study of airborne organisms and their distribution; biology of aeroplankton.

aerobiosis (â'ëröbiō'sis) n. [Gk. aer, air; biosis, manner of life.] Existence in presence of oxygen.

aerocyst (ā'ërösist) n. [Gk. aer, air; kystis, bladder.] An air vesicle of algae.

aerogenic (ā'ērojēn'ik) a. [Gk. aer, air; gennaein, to produce.] Gasproducing; appl. certain bacteria.

aeromorphosis (ā'erömôr'fōsis, -môrfō'sis) n. [Gk. aer, air; morphosis, form.] Modification of form or structure owing to exposure to air or wind.

aerophyte (ā'ëröfīt) n. [Gk. aer, air; phyton, plant.] A plant growing attached to an aerial portion of another plant; epiphyte.

aeroplankton (ā'ēröplăngk'tŏn) n.
[Gk. aer, air; plangktos. wandering.]
Living particles drifting in the air,
as spores, pollen, bacteria, etc.; also
applied to non-living particles.

aerostat (ā'ëröstăt) n. [L. aer, air; stare, to stand.] An air-sac in

insect body or in bird bone.

aerostatic (ā'ëröstāt'ik), a. [L. aer, air; stare, to stand.] Pneumatic; containing air-spaces.

aerotaxis (ā'ērötāk'sīs) n. [Gk. aer, air; taxis, arrangement.] The arrangement of bacteria and other micro-organisms towards or away from oxygen.

aerotropic (ă'ērötrŏp'ik) a. [Gk. aer, air; trope, turn.] Appl. curvature of a plant organ towards a higher

concentration of oxygen.

AGE-

aerotropism (āërŏt'röpizm) n. [Gk. aer, air; trope, turn.] Reaction to gases, generally to oxygen.

aesthacyte (ēs'thăsīt) n. [Gk. aisthesis, sensation; kytos, hollow.] A sensory cell of primitive animals.

aesthesis (ēsthē'sīs) n. [Gk. aisthesis, sensation.] Sensibility; sensepercept; aesthesia.

aesthetasc (ēsthē'tăsk) n. [Gk. aisthetes, perceiver; askein, to exercise.] An olfactory receptor on antennule of daphnids.

aesthetes (ēsthē'tēz) n. plu. [Gk. aisthetes, perceiver.] Sense organs.

aestival (ēstī'văl, ēs'tĭvăl) a. [L. aestivus, of summer.] Produced in, or pert. summer.

aestivation (ēs'tīvā'shŭn) n. [L. aestivus, of summer.] The mode in which different parts of flower are disposed in flower-bud; prefloration; torpor during summer, as in some animals; dormancy during heat and drought period, opp. hibernation.

aethalium (ēthā'lĭŭm) n. [Gk. aithalos, soot.] An aggregation of plasmodia or sporangia to form a compound fruit in Mycetozoa or Myxomycetes.

aetiology (ētiol'oji) n. [Gk. aitia, cause; logos, discourse.] The science of causation; or origin of causes; also etiology.

afferent (af'erent) a. [L. afferre, to bring.] Bringing towards; appl. nerves carrying impulses to nervous centres; appl. blood-vessels carrying blood to an organ or set of organs.

aflagellar (å'flajel'ar) a. [Gk. a, without; L. flagellum, whip.]

Without a flagellum.

afterbirth (âf'tërbërth) n. [A.S. aefter, behind; beran, to bring forth.] Placenta and foetal membranes expelled after offspring.

afterbrain,—myelencephalon, q.v. aftershaft (âf'tërshâft) n. [A.S. aefter, behind; O.E. sceaft, shaft.]

A small tuft of down near superior umbilicus of a feather.

agameon (ăgămē'ŏn) n. [Gk. a,

without; gamos, marriage; on, being.] A species comprising only apomictic individuals.

agamete (ăg'ămēt) n, [Gk. a, without; gametes, spouse.] An amoebula, or germ cell, which develops directly without syngamy into an adult.

agametoblast (ăgămē'toblāst) n. [Gk. a, not; gametes, spouse; blastos, bud.] A cytomere, q.v., of Caryotropha.

agamic (ăgăm'îk), a. [Gk. a, without; gamos, marriage.] Asexual; parthenogenetic; cryptogamic; agamous.

agamobium (ăg'ămöb'îŭm) n. [Gk. a, without; gamos, marriage; bios, life.] The asexual stage in metagenesis; the sporophyte.

agamogenesis (ăg'ămöjĕn'ësis) n. [Gk. a, without; gamos, marriage; genesis, descent.] Asexual reproduction; parthenogenesis.

agamogenetic (ăg'ămöjēnět'îk) a. [Gk. a, without; gamos, marriage; genesis, descent.] Asexual; produced asexually.

agamogony (ăg'ămŏg'öni) n. [Gk. a, without; gamos, marriage; gonos, generation.] Schizogony, or reproduction without sexual process.

agamont (ag'amont) n. [Gk. a, without; gamos, marriage; on, being.]
A schizont, or that stage which gives rise to agametes.

agamospecies (ăg'ămöspē'shēz) n.
[Gk. a, without; gamos, marriage;
L. species, particular kind.] Species
without sexual reproduction, as in
parthenogenetic aneuploids.

agamous,-agamic, q.v.

agar (âg'âr, ā'gâr) n. [Mal. agar-agar, a sea-weed.] A medium for bacterial and other cultures, prepared from agar-agar, a gelatinous substance, also of dietary utility, yielded by red algae.

age and area,—hypothesis of Willis that older species occur in a more extensive area than that occupied by more recent species.

ageotropic,—apogeotropic, q.v.

agglomerate (áglóm'erāt) a. [L. ad, to; glomus, ball.] Clustered, as a head of flowers; appl. adhering mass of protozoa, as in agglomeration of trypanosomes.

agglutination (ăg'lootinā'shun) n. [L. ad, to; glutinare, to glue.] The formation of clumps or floccules by pollen, bacteria, erythrocytes, spermatozoa, and some protozoa.

agglutinin (agloot'inin) n. [L. ad, to; glutinare, to glue.] A substance or specific antibody which

causes agglutination.

- agglutinogen (ăglootin'öjen) n. [L. ad, to; glutinare, to glue; Gk. gennaein, to produce.] Substance or antigen that produces agglutinin.
- aggregate (ăg'rëgāt) a. [L. ad, to; gregare, to collect into a flock.] Formed in a cluster; appl. a fruit or etaerio formed from apocarpous gynoecium of a single flower, as raspberry; appl. certain medullary rays; appl. a type of silk gland in certain spiders.
- aggregation (ăgregā'shun) n. [L. ad, to; gregare, to collect.] A grouping or crowding of separate organisms; movement of protoplasm in tentacle or tendril cells of sensitive plants, which causes tentacle or tendril to bend towards the point stimulated.
- aggressin (ăgres'in) n. [L. aggressus, Toxic substance proattacked.] duced by pathogenic organisms, inhibiting defensive reactions of host.
- aglomerular (ăglŏmĕr'ūlăr) a. [Gk. a, without; L. glomerare, to form into a ball.] Devoid of glomeruli, as kidney in certain fishes.
- aglossate (aglos'at) n. [Gk. a, without; glossa, tongue.] Having no tongue.
- agminated (ăg'mināted) a. ſL. agmen, herd.] Clustered; appl. glands, Peyer's patches.
- agnathostomatous (ăgnăth'östöm'ătus) a. [Gk. a, without; gnathos, stoma, mouth.] Having mouth unfurnished with jaws, as lamprey; agnathous.

agon (ăg'on) n. [Gk. agon, contest.] The active principle of an enzyme; cf. pheron, symplex.

agonist (ăg'onist) n. [Gk. agonistes, champion.] A prime mover or muscle directly responsible for change in position of a part.

- agranulocyte (ăgrăn'ūlösīt) n. [Gk. a, without; L. granulum, small grain; Gk. kytos, hollow.] non-granular or lymphoid leucocyte.
- agrestal (ăgres'tăl) a. [L. agrestis, rural.] Appl. plants growing on arable land.
- agriotype (ăg'riötīp) n. [Gk. agrios, wild; typos, image.] Wild or ancestral type.
- agrostology (àg'rőstől'öji) n. agrostis, grass; logos, discourse.] That part of botany dealing with grasses.
- aheliotropism,—apheliotropism, q.v. A-horizon,—the upper, or leached, soil layers.
- air-bladder (ār'-blăd'er) n. [L. aer, air; A.S. bladder.] The swim-bladder in fishes; hollow dilatation of thallus in bladderwrack
- air-cells,-thin-walled cavities ethmoidal labyrinth; numerous cavities in mastoid; alveoli of lungs; air spaces in plant tissue.

air-pore,—stoma, q.v., of plants.

- air-sacs,—spaces filled with air and connected with lungs in birds; dilatations of tracheae in many insects; sacs representing tracheal system and having hydrostatic function in certain insect larvae.
- air sinuses,—cavities in frontal ethmoid, sphenoid and maxillary bones, with passages to cavities.
- aitionastic (ī'tiönăs'tīk) a. [Gk. aitios, causing; nastos, pressed.] Appl. curvature of part of a plant and induced by a diffuse stimulus.
- akanth-,-see acanth-.
- akaryocyte,—akaryote; an erythrocyte.

- akaryote (ākār'iōt) n. [Gk. a, without; karyon, nut.] A cell in which nucleoplasm has not collected together to form a nucleus; a non-nucleated cell, condition present in many protista. a. Non-nucleated.
- akene,-achene.
- akinesis (ăkinē'sis) n. [Gk. a, not; kinesis, movement.] Absence or arrest of motion.
- akinete (āk'īnēt) n. [Gk. a, not; kinein, to move.] A resting cell in certain green algae, which will later reproduce.
- ala (ā'lā, àlā) n. [L. ala, wing.] Any winglike projection or structure; lateral petal of papilionaceous flower; membranous expansion on some seeds. Plu. alae.
- alar (ā'lār) a. [L. ala, wing.] Wing-like; pert. wings or alae; axillary; appl. ligaments, cartilages, etc.

alary (ā'lări) a. [L. ala, wing.] Wing-like; pert. wings.

- alate (ā'lāt) a. [L. alatus, winged.] Having a wing-like expansion, as of petiole or stem; broad-lipped, appl. shells; appl. a spicular system in Calcarea which is sagittal because of inequality of angles; winged.
- albedo (ālbē'dö) n. [L. albus, white.] Diffused reflection, the ratio of the amount of light reflected by a surface to the amount of incident light; mesocarp of hesperidium, cf. flavedo.
- albescent (ălbes'ent) a. [L. albescere, to grow white.] Growing whitish.
- albicant (ăl'bīkănt) a. [L. albicare, to grow white.] Tending to become white.
- albinism (ăl'binizm) n. [L. albus, white.] Absence of pigmentation in animals normally pigmented; state of having colourless chromatophores.
- albino (ălbē'nö, ălbī'nö) n. [Sp. albino, white, from L. albus.] Any animal with congenital deficiency of pigment in skin, hair, eyes, etc.; a plant with colourless chromator phores, due to absence of chloroplasts or undeveloped chromoplasts.

- albuginea (ălbūjīn'ēă) n. [L. albus, white; gignere, to beget.]
 Tunica albuginea: white, dense connective tissue surrounding testis, ovary, corpora cavernosa, spleen, or eye.
- albumen (ălbū'men) n. [L. albumen, white of egg.] White of egg; nutritive material stored in seed.
- albumin (ălbū'mīn) n. [I.. albumen, white of egg.] One of a group of heat-coagulable, water-soluble proteins occurring in egg-white, blood serum, milk, and many animal and vegetable tissues.
- albuminoids (ălbū'minoidz) n. plu. [L. albumen, white of egg; Gk. eidos, form.] Scleroproteins, g.v.
- albuminous (ălbū'minus) a. [L. albumen, white of egg.] Pert., containing, or of nature of, albumen or an albumin.
- **albuminous cells,** parenchyma cells associated with sieve cells, as in pteridophytes and gymnosperms.
- alburnum (ălbūr'nūm) n. [L. albus, white.] Sap-wood or splint-wood, soft white substance between inner bark and true wood; outer young wood of dicotyledon.
- alecithal (ălĕs'ithăl) a. [Gk. a, without; lekithos, yolk.] With little or no yolk; appl. ova; alecithic.
- aletocyte (ale'tösit) n. [Gk. aletes, wanderer; kytos, hollow.] Wandering cell.
- aleuron (ălū'rŏn) n. [Gk. aleuron, flour.] Appl. protein grains found in general protoplasm and used as reserve food-material; appl. layer containing protein, of endosperm in monocotyledons; aleurone.
- aleurospore (ălū'röspōr) n. [Gk. aleuron, flour.] A lateral conidium of certain fungous parasites of skin; spore or tip early separated from hypha by a septum.
- alexine (ălĕk'sin) n. [Gk. alexein, to ward off.] A substance in blood serum which combines with an amboceptor to produce lysis; alexin; complement.
- algesis (ăljē'sis) n. [Gk. algesis, sense of pain.] The sense of pain.

- algin (ăl'jin) n. [L. alga, seaweed.] A mucilaginous substance, alginic acid, obtained from certain algae.
- algoid (ăl'goid) a. [L. alga, seaweed; Gk. eidos, shape.] Pert., resembling, or of the nature of an alga.

algology (ălgŏl'öjĭ) n. [L. alga, seaweed; Gk. logos, discourse.] The study of algae; phycology.

Algonkian (ălgŏng'kĭăn) a. [Algonquian tribe of Indians.] Pert. late Proterozoic era.

alima (ăl'îmă) n. [Gk. halimos, pert. sea.] A larval stage of certain Crustacea.

- alimentary (ăl'iměn'tări) a. [L. alere, to nourish.] Pert. nutritive functions; appl. system, canal, tract, etc.
- alisphenoid (ăl'îsfē'noid) n. [L. ala, wing; Gk. sphen, wedge; eidos, form.] Wing-like portion of sphenoid forming part of cranium; ala temporalis.
- alitrunk (äl'itrungk) n. [L. ala, wing; truncus, trunk.] Thorax of insect when fused with first segment of abdomen.
- alkaloid (ăl'kăloid) n. [Ar. al, the; qali, ash; Gk. eidos, form.] Basic nitrogenous organic substance with poisonous or medicinal properties, as caffeine, morphine, nicotine, strychnine, ptomaine, etc.

allaesthetic (äl'ēsthēt'īk) a. [Gk. allos, other; aisthetes, perceiver.]

Appl. characters effective when perceived by other organisms.

allantochorion (ălăn'tökō'rĭŏn) n. [Gk. allas, sausage; chorion, skin.] Foetal membrane formed of outer wall of allantois and the primitive chorion; true chorion.

allantoid (alan'toid) a. [Gk. allas, sausage; eidos, form.] Sausage-shaped; botuliform.

allantoin (ălăn'töin) n. [Gk. allas, sausage.] The end-product of purine metabolism, occurring in allantoic fluid and urine of certain mammals; C₄H₆O₃N₄.

allantois (ălăn'töis) n. [Gk. allas, sausage.] An embryonic organ, a

membranous sac arising from posterior part of alimentary canal in higher vertebrates, and acting as an organ of respiration or nutrition or both.

allassotonic (ălăs'ötŏn'īk) a. [Gk. allassein, to change; tonos, strain.] Induced by stimulus, appl. movements of grown plants; cf. auxotonic.

allele (alēl') n. [Gk. allelon, one another.] Allelomorph.

allelism (ale'lizm) n. [Gk. allelon, one another.] The relationship between two alleles; allelomorphism.

- allelocatalysis (ălē'lokātāl'īsīs) n. [Gk. allelon, one another; katalysis, dissolution.] Allelocatalytic or mutually accelerating or retarding effect of contiguous cells; e.g. acceleration of rate of fission with increase in number of individual protozoa present.
- allelomorph (ălē'lömôrf) n. [Gk. allelon, one another; morphe, form.] One of any pair of alternative hereditary characters; gene which can occupy the same locus as another gene in a particular chromosome.
- allelopathy (ălēlŏp'āthĭ) n. [Gk. allelon, one another; pathos, suffering.] The influence or effect of one living plant upon another.

allergen (ăl'ërjen) n. [Gk. allos, other; ergon, activity; -genes, producing.] A substance which induces allergy; atopen.

allergy (ăl'ērjī) n. [Gk. allos, other; ergon, activity.] Changed reactivity on second or subsequent infection or poisoning; exaggerated or unusual susceptibility; anaphylaxis; atopy, q.v.

allesthetic,—allaesthetic, q.v.

alliaceous (ăliā'shūs) a. [L. allium, garlic.] Pert. or like garlic; appl. a class of odours.

allobiosis (al'öbiō'sis) n. [Gk. allos, other; biosis, manner of life.] Changed reactivity of an organism in a changed internal or external environment.

- allocarpy (ăl'ökârpi) n. [Gk. allos, other; karpos, fruit.] The production of fruit after cross-fertilisation.
- allochroic (ăl'ökrō'ĭk) a. [Gk. allos, other; chros, colour.] Able to change colour; with colour variation.
- allochronic (al'okron'ik) a. [Gk. allos, other; chronos, time.] Not contemporary; appl. species, etc. Opp. synchronic.

allochthonous (ălŏk'thönūs) a. [Gk. allos, other; chthon, the ground.] Exotic; not aboriginal; acquired.

Opp. autochthonous.

- allocortex (ăl'ökôr'těks) n. [Gk. allos, other; L. cortex, bark.] The primitive cortical areas or cortex of olfactory brain, opp. isocortex.
- allogamous (ălŏg'ămŭs) a. [Gk. allos, other; gamos, marriage.] Reproducing by cross-fertilisation, opp. autogamous.
- allogamy (alog'ami) n. [Gk. allos, other; gamos, marriage.] Crossfertilisation, opp. autogamy.

allogene (ăl'ojēn) n. [Gk. allos, other; genos, descent.] A recessive allelomorph; opp. protogene.

allogenic (ălöjěn'îk) a. [Gk. allos, other; genos, descent.] Caused by external factors; appl. plant successions; pert. allogenes; derived from elsewhere, opp. autogenic; allogenous or exogenous, opp. endogenous; allochronic, q.v.

alloheteroploid (ăl'öhĕt'ëröploid) n.
[Gk. allos, other; heteros, other; aploos, onefold; eidos, form.]
Heteroploid derived from specific-

ally distinct genomes.

alloiogenesis (ăl'oiöjēn'ēsīs) n. [Gk. alloios, different; genesis, descent.]
The alternation, in a life-history, of a sexual and a non-sexual form; alternation of generations.

alloiometron (ăl'oiöm'ētron) n. [Gk. alloios, different; metron, measure.] Measurable change of proportion or intensity of development within species or races, e.g. head, limb, tooth, etc. proportions.

allokinesis (ăl'ökinē'sis) n. [Gk. allos, other; kinesis, movement.] Reflex, or passive, movement; involuntary movement.

allometry (alom'etri) n. [Gk. allos, other; metron, measure.] Study of relative growth; change of proportions with increase of size; growth rate of a part differing from a standard growth rate or from the growth rate of the whole.

alloparalectotype (ål'opārālēk'tötīp)
n. [Gk. allos, other; para, beside;
lektos, chosen; typos, pattern.]
Specimen, from the original collection, of the sex opposite to that of
the holotype, and described sub-

sequently.

allopatric (ălöpăt'rīk) a. [Gk. allos, other; patra, native land.] Having separate and mutually exclusive areas of geographical distribution. Opp. sympatric.

allopelagic (ăl'öpĕlāj'īk) a. [Gk. allos, other; pelagos, sea.] Pert. organisms found at any depth of

the sea.

alloplasm (ăl'öplăzm) n. [Gk. allos, other; plasma, mould.] The differentiated portion of cell-substance not forming independent organelles.

alloplasmatic (āl'öplāzmāt'īk) a. [Gk. allos, other; plasma, mould.] Appl. differentiated portion of cell

protoplasm; alloplasmic.

alioplast (ăl'öplăst) n. [Gk. allos, other; plastos, formed.] A morphological cell-unit of more than one kind of tissue; opp. homoplast.

- allopolyploid (āl'öpŏl'iploid) n. [Gk. allos, other; polys, many; aploos, onefold; eidos, form.] An organism with more than two sets of chromosomes derived from different species by hybridisation.
- allorhizal (äl'örī'zăl) a. [Gk. allos, other; rhiza, root.] Having opposed root and shoot poles. Opp. homorhizal.
- all-or-none,—principle that response to a stimulus is either completely effected or is absent, first observed in heart muscle (Bowditch's law).

allosematic (ăl'osēmăt'ik) a. [Gk. allos, other; sema, sign.] Having markings or coloration imitating warning signs in other, usually

dangerous, species.

allosomal (ăl'oso'măl) a. [Gk. allos, other; soma, body.] Pert. allosome; appl. inheritance of characters controlled by genes located in an allosome.

allosome (ăl'ösōm) n. [Gk. allos, other; soma, body.] A chromosome other than an ordinary or typical one; heterochromosome, cf. autosome.

allostoses (ăl'osto'sēs) n. plu. [Gk. allos, other; osteon, bone.] Bones formed in membrane; cf. autostoses. allosynapsis,—allosyndesis, q.v.

- allosyndesis (ăl'ösĭn'desis) n. [Gk. allos, other; syndesis, a binding together.] Pairing of homologous chromosomes from opposite parents, in a polyploid; cf. autosyndesis.
- allotetraploid,—amphidiploid, q.v. allotherm (ăl'otherm) n. [Gk. allos, other; therme, heat.] An organism with body temperature dependent on environmental temperature.
- allotrophic (ăl'ötrŏf'ĭk) a. allos, other; trophe, nourishment.] Obtaining nourishment from other organisms; saprophytic, or saprozoic; heterotrophic.
- allotropic (ălötrop'ik) a. [Gk. allos, other; trope, turn.] Exhibiting mutual tropism, as between gametes.
- allotype (ăl'ötīp) n. [Gk. allos, other; typos, pattern.] Paratype of the sex opposite to that of the holotype.
- allozygote (ăl'özī'gōt) n. [Gk. allos, other; zygon, yoke.] A homozygote having recessive characters; opp. protozygote.
- alluvial (ălū'vĭāl, ăloo-) a. alluere, to wash to.] Pert. deposits formed by finely divided material laid down by running water.
- alpha(α) cells,—oxyphilic cells in pars glandularis of pituitary gland; cells with granules insoluble in alcohol, in islets of Langerhans; A-cells.

- alpha (a) granules,—metachromatic granules in central region of protoplast. as in blue-green algae.
- alpha (a) tocopherol,—vitamin E.
- alsinaceous (ălsĭnā'shŭs) a. [Gk. alsine, chickweed.] Appl. polypetalous corolla where intervals occur between petals, as in chickweed.
- alteration theory,—explains electromotive forces of nerve and muscle by alterations in chemical composition of tissue at cross-section.
- alternate (ăl-, ôltěr'nāt) a. alternus, one after another.] Not opposite; appl. leaves, branches, etc., occurring at different levels successively on opposite sides of stem; every other; taking turns.

alternating cleavage, spiral cleavage, q.v.

alternation of generations,—the occurrence in one life-history of two or more different forms differently produced, usually an alternation of a sexual with an asexual form; alloiogenesis; metagenesis; digenesis; heterogamy; genesis; heterogony.

alternation of parts,—general rule that leaves of different whorls alternate in position with each other, sepals with petals, stamens

with petals.

alternative inheritance,-allelism, allelomorphism.

- alterne (ăltěrn', ôl-) n. [L. alternus, one after another.] Vegetation exhibiting disturbed zonation due to abrupt change in environment, or to interference with normal plant succession.
- alternipinnate (ăltěr'nipin'āt, ôl-) a. [L. alternus, one after another; pinna, wing.] Appl. leaflets or pinnae arising alternately on each side of mid-rib.
- altrices (ăltrī'sēz, ăl'trīsēz) n. plu. [L. altrix, nourisher.] Birds whose young are hatched in a very immature condition; cf. praecoces.
- altricial (ăltris'iăl) a. [L. altrix, Requiring care or nourisher. nursing after hatching or birth.

alula (ăl'ūlă) n. [L. alula, dim. of ala, wing.] A small lobe separated off from wing-base on its posterior edge in certain insects; spurious or bastard wing of birds.

alutaceous (ălūtā'shūs) a. [L. aluta, alum-dressed leather.] Tancoloured; leathery; having appearance of minute cracks, appl. markings on elytra of certain beetles.

alveola (ăl'veölă) n. [L. alveolus, small cavity.] A pit on the surface

of an organ; alveolus, q.v.

alveolar (ăl'vrolăr) a. [L. alveolus, small pit.] Pert. an alveolus; pert. tooth socket; appl. artery, nerve, process, canal, in connection with the jaw-bone; appl. small cavities in lungs, glands, etc.; appl. a theory of structure of protoplasm.

alveolate (ăl'veolat, ălve'olat) a. [L. alveolatus, pitted.] Deeply pitted

or honey-combed.

alveolation (ăl'veólā'shŭn) n. [L. alveolatus, pitted.] The formation of alveoli; alveolate appearance.

- alveolus (ălvē'olūs) n. [L. alveolus, small pit.] A small pit or depression; tooth socket; pyramidal ossicle, supporting tooth in seaurchin; air-cell of lung; a cavity in glands; cavity in tarsus of spiders, receptacle for haematodocha; pit for articulation of macrotrichia.
- alveus (ăl'veŭs) n. [L. alveus, cavity.] A white layer of fibres on ventricular surface of hippocampus; utricle of ear; dilatation of thoracic duct.
- amacrine (ăm'ăkrin) a. [Gk. a, not; makros, long; is, fibre.] Having no axon; appl. cells in inner nuclear layer of retina, with dendrites in inner plexiform layer.
- amb (amb) n. [L. ambulare, to walk.]
 Ambulacral area.
- ambiens (ăm'biĕnz) n. [L. ambire, to go round.] A thigh muscle in certain birds, the action of which causes the toes to maintain grasp on perch.
- ambient (ăm'biënt) a. [L. ambire, to go round.] Surrounding; appl.

vein, the costal nervure when encircling insect wing.

ambiparous (ămbip'ārūs) a. [L. ambo, both; parere, to produce.] Containing the beginnings of both flowers and leaves; appl. buds.

ambisexual (ăm'bisek'suăl) a. [L. ambo, both; sexus, sex.] Pert. both sexes; ambosexual; monoecious an

cious, q.v.

ambisporangiate (ăm'bisporăn'jiāt)
a. [L. ambo, both; Gk. sporos, seed; anggeion, box.] Amphisporangiate, q.v.

ambital (ām'bītāl) a. [L. ambire, to go round.] Appl. interambulacral and antambulacral plates of asteroids; outer skeleton of ophiuroid arm.

ambitus (ăm'bitus) n. [L. ambire, to go around.] The outer edge or margin; outline of echinoid shell viewed from apical pole.

amboceptor (ăm'bösĕp'tŏr) n. [L. ambo, both; capere, to take.] A specific antibody or immune body necessary for ferment-like action of complement on a toxin or a red blood corpuscle; a lysin.

ambon (am'bon) n. [Gk. ambon, raised platform.] Fibrocartilaginous ring surrounding an articular socket, as around acetabulum; circumferential fibrocartilage; labrum.

ambosexual (ăm'bösĕk'sūăl) a. [L. ambo, both; sexus, sex.] Common to, or pert., both sexes; activated by both male and female hormones.

ambulacra (ăm'būlā'kră) n. plu. [L. ambulare, to walk.] Locomotor tube-feet of echinoderms.

ambulacralia (ăm'būlăkrā'lĭā) n. plu. [L. ambulare, to walk.] Ambulacral plates, i.e. plates through which tube-feet protrude.

ambulacriform (ămbūlāk'rīfôrm) a. [L. ambulare, to walk; forma, shape.] Having the form or appearance of ambulacra.

ameba,—amoeba.

ameiosis (āmīō'sis) n. [Gk. a, without; meiosis, diminution.] Occurrence of only one division in meiosis instead of two.

18 AME-

ameiotic (amīot'ik) a. [Gk. a, without; meion, smaller.] Appl. parthenogenesis in which meiosis is suppressed.

(ăměl'ĭfĭkā'shŭn) n. amelification [M.E. amell, enamel; L. facere, to make.] Formation of toothenamel.

ameloblast (ăměl'öblast) n. [M.E. amell, enamel; Gk. blastos, bud.] A columnar or hexagonal cell of epithelium of enamel internal organ; enamel cell, adamantoblast,

ganoblast.

amentaceous (ămenta'shus), amen-(ăměntĭf'ërŭs) a. tiferous amentum, thong; ferre, to carry.] Appl. plants bearing amenta or

- amentum (ăměn'tům) n. [L. amentum, thong.] A catkin, consisting of bracted axis bearing unisexual flowers, as in poplar and willow; ament.
- ameristic (ăměris'tik) a. [Gk. a, without; meristos, divided.] Not divided into parts; unsegmented.

ametabolic (āměť abol'ík) a. [Gk. a, without; metabole, change.] Appl. insects that do not pass through marked metamorphosis.

amicron (amik'ron) n. [Gk. a, without; mikros, small.] An element so small that even the ultramicroscope can only indicate it as a diffuse illumination in the track of the beam; cf. submicron.

amicronucleate (ămī'krönū'klēāt) a. [Gk. a, without; mikros, small; L. nucleus, kernel.] Appl. fragments of certain Protozoa in which there

is no micronucleus.

amine (ăm'în) n. [Gk. ammoniakon, resinous gum.] A nitrogen compound formed in plants, also produced by bacterial action on amino acids, a derivative from ammonia by hydrogen replacement.

amino acids, compounds containing amino (NH2) and carboxyl (COOH) groups, and produced from proteins by hydrolysis.

amitosis (ămītō'sīs) n. [Gk. a, without; mitos, thread.] Direct celldivision and cleavage of nucleus without thread-like formation of nuclear material. Opp. mitosis.

ammonitiferous (ăm'onitif'ërus) a. [Gk. Ammon, Jupiter; L. ferre, to carry.] Containing fossil remains of ammonites.

- amnion (ăm'nion) n. [Gk. amnion, foetal membrane.] A foetal membrane of reptiles, birds, and mammals; inner embryonic membrane of insects; viscous envelope of certain ovules.
- amnionic, -- amniotic.
- amniote (ăm'niot) n. [Gk. amnion, foetal membrane.] An animal characterised by possession amnion in foetal life.
- amniotic (ămniŏt'ik) a. [Gk. amnion, foetal membrane.] Pert. amnion; appl. folds, sac, cavity, fluid; amnionic.
- amoeba (ămē'bă) n. [Gk. amoibe, change.] A protozoon in which the shape is subject to constant alterations due to formation and retraction of pseudopodia; generally used to typify most primitive animal commonly known.
- amoebadiastase (ămē'bădī'āstās) n. [Gk. amoibe, change; dia, through; histanai, to set.] The digestive ferment secreted by amoebae.
- amoebiform (ămē'bĭfôrm) a. [Gk. amoibe, change; L. forma, shape.] Shaped like or resembling an amoeba.
- amoebism (ămē'bĭzm) n. [Gk. amoibe, change.] Amoeboid form or behaviour, as of leucocytes.
- amoebocyte (ămē'bösīt) n. amoibe, change; kytos, hollow.] Any cell having the shape or properties of an amoeba; one of certain cells in coelom of echinoderms; a leucocyte, q.v.
- amoeboid (ămē'boid) a. [Gk. amoibe, change; eidos, shape.] Resembling an amoeba in shape, in properties, or in locomotion.
- amoebula (ămē'būlă) n. [Gk. amoibe, The swarm-spore of a change. protist when furnished with pseudopodia; pseudopodiospore.

amorphous (ămôr'fus) a. [Gk. a, without; morphe, shape.] Of indeterminate or irregular form; with no visible differentiation in structure.

amphiapomict (ăm'fiăp'ömĭkt) n. [Gk. amphi, both; apo, away; miktos, mixed.] A biotype reproduced from facultative sexual forms.

amphiarthrosis (ăm'fiârthrō'sīs) n. [Gk. amphi, both; arthron, joint.] A slightly movable articulation, as a symphysis or a syndesmosis.

amphiaster (ăm'fiăs'tër) n. [Gk. amphi, both; aster, star.] The two asters connected by the achromatic spindle formed in mitotic cell division.

amphiastral (ăm'fiăs'trăl) a. [Gk. amphi, both; aster, star.] Appl. a type of mitosis in which true asters are present at the spindle-poles.

amphibious (ămfīb'īūs), amphibian (ămfīb'īān) a. [Gk. amphi, both; bios, life.] Adapted for life either on land or in water; emersed, q.v.

amphiblastic (ăm'fiblăs'tik) a. [Gk. amphi, both; blastos, bud.] Appl. telolecithal ova with complete but unequal segmentation.

amphiblastula (ăm'fīblăs'tūlā) n. [Gk. amphi, both; blastos, bud.] Stage in development of certain sponges, in which posterior end of embryo is composed of granular archaeocytes, and anterior end of flagellate cells.

amphibolic (ăm'fibòl'îk) a. [Gk. amphi, both; bole, throw.] Capable of turning backwards or forwards, as outer toe of certain birds.

amphicarpous (ăm'fikâr'pŭs) a. [Gk. amphi, both; karpos, fruit.] Producing fruit of two kinds.

amphicoelous (ăm'fisē'lūs) a. [Gk. amphi, both; koilos, hollow.] Concave on both surfaces; appl. biconcave vertebral centra; amphicelous.

amphicondylous (ām'fīkŏn'dĭlŭs) a. [Gk.amphi, both; kondylos, knuckle.] Having two occipital condyles.

amphicone (ăm'fikōn) n. [Gk. amphi, both; konos, cone.] Cusp of molar of extinct mammals, believed to have evolved into metacone and paracone. amphicribral,-amphiphloic.

amphicytes (ăm'fisits) n. plu. [Gk. amphi, both; kytos, hollow.] Endothelial cells surrounding, or forming, capsules of cells of a dorsal root ganglion; capsule cells.

amphidelphic (ăm'fiděl'fik) a. [Gk. amphi, both; delphys, womb.] Having a paired uterus, as in certain

nematodes; didelphic.

amphidetic (ăm'fidět'îk) a. [Gk. amphi, both; detos, bound.] Extending behind and in front of umbo; appl. hinge ligaments of some bivalve shells; cf. opisthodetic.

amphidiploid (ăm'fidip'loid) a. [Gk. amphi, both; diploos, double.]
Double diploid; allotetraploid. n.
A hybrid having diploid genomes of both parental species.

amphidisc (ām'fidisk) n. [Gk. amphi, both; diskos, round plate.] A grapnel-shaped spicule of some

freshwater sponges.

amphids (ăm'fidz) n. plu. [Gk. amphi, both.] Two anterior lateral sensory organs in nematodes.

amphigastria (ăm'figăs'triă) n. plu. [Gk. amphi, both; gaster, stomach.] Rudimentary leaves, or scales, on under surface of foliose liverworts.

amphigenesis (ăm'fijēn'ēsīs) n. [Gk. amphi, both; genesis, descent.]
Amphigony; sexual reproduction.

amphigonic (ăm'figŏn'ik) a. [Gk. amphi, both; gone, seed.] Producing male and female gametes in separate gones in different individuals; bisexual; pert. amphigony; cf. digonic, syngonic.

amphigony (\(\text{amfig'oni}\)) n. [Gk. amphi, both; gonos, offspring.]
Reproduction involving two individuals; sporogony; amphi-

genesis.

amphigynous (ămfij'înŭs) a. [Gk. amphi, both; gyne, female.] Appl. antheridium surrounding the base of the oogonium, as in some Peronosporales.

amphikaryon (ăm'fikăr'iŏn) n. [Gk. amphi, both; karyon, nut.] An amphinucleus or nucleus with large

karyosome (in reference to supposed encapsuling of kinetic nucleus by trophic nucleus); nucleus with two haploid sets of chromosomes.

amphimict (ăm'fimikt) n. [Gk. amphi, both; miktos, mixed.] A biotype resulting from sexual reproduction; an obligate sexual

organism.

AMP-

amphimixis (ām'fīmīk'sīs) n. [Gk. amphi, both; mixis, mingling.] The mingling of paternal and maternal characteristics by union of male and female pronuclei in fertilisation. Opp. apomixis.

amphinucleolus (ăm'finūklē'ölūs) n. [Gk. amphi, both; L. nucleolus, a small kernel.] A double nucleolus comprising basiphil and oxyphil

components.

amphinucleus,—amphikaryon, q.v. amphiodont (ăm'fiödŏnt) a. [Gk. amphi, both; odous, tooth.] Appl. an intermediate state of mandible development in stag-beetles.

amphiont (am'fiont) n. [Gk. amphi, both; on, being.] Zygote or sporont formed by coming together

of two individuals.

amphiphloic,—periphloic, q.v.

- amphiplatyan (ăm'fiplătiăn) a. [Gk. amphi, both; platys, flat.] Flat on both ends; appl. vertebral centra.
- amphipneustic (ăm'finū'stĭk, -pn-) a. [Gk. amphi, both; pnein, to breathe.] Having both gills and lungs throughout life-history; with only anterior and posterior pairs of spiracles functioning, as in most dipterous larvae; amphipneustous.

amphipodous (ămfīp'ödūs) a. [Gk. amphi, both; pous, foot.] Having feet for walking and feet for

swimming.

amphipyrenin (ăm'fipīrē'nin) n. [Gk. amphi, both; pyren, fruit-stone.]
Substance of which nuclear membrane is composed.

amphirhinal (ăm'firi'năl) a. [Gk. amphi, both; rhis, nose.] Having, or pert., two nostrils.

amphisarca (ăm'fīsâr'kă) n. [Gk. amphi, both; sarx, flesh.] A

superior indehiscent many-seeded fruit with pulpy interior and woody exterior.

amphispermous (ăm'fīspēr'mŭs) a. [Gk. amphi, both; sperma, seed.] Having seed closely surrounded by

pericarp.

- amphisporangiate (ăm'fispörăn'jiāt)
 a. [Gk. amphi, both; sporos, seed; anggeion, vessel.] Having sporophylls bearing both megasporangia and microsporangia; hermaphrodite, appl. flowers.
- amphispore (ăm'fispor') n. [Gk. amphi, both; sporos, seed.] A reproductive spore which functions as a resting spore in certain algae mesospore; a uredospore modified to withstand dry environment.
- amphisternous (ăm'fīstēr'nūs) a. [Gk. amphi, both; sternon, breastbone.] Appl. type of sternum structure in Atelostomata.
- amphistomatic (ăm'fistömăt'îk) a. [Gk. amphi, both; stoma, mouth.] Having stomata on both surfaces, appl. certain types of leaves.

amphistomous (ămfīs'tömŭs) a.
[Gk. amphi, both; stoma, mouth.]
Having a sucker at each end of body, as certain worms.

- amphistylic (äm'fistil'ik) a. [Gk. amphi, both; stylos, pillar.] Having jaw arch connected with skull by both hyoid and quadrate, or by both hyoid and palato-quadrate; exhibiting condition of amphistyly.
- amphitene (ăm'fitēn) a. [Gk. amphi, both; tainia, band.] Stage of meiosis in which spireme threads are uniting in pairs; zygotene.

amphithecium (ămfĭthē'sĭūm) n. [Gk. amphi, both; thekion, box.] Peripheral layer of cells in sporangia of liverworts and mosses.

- amphitoky (ămfit'öki) n. [Gk. amphi, both; tokos, birth.] Parthenogenetic reproduction of both males and females.
- amphitriaene (ăm'fitrī'ēn) n. [Gk. amphi, both; triaina, trident.] A double trident-shaped spicule.

- amphitrichous (ămfit'rikŭs) a. [Gk. amphi, both; thrix, hair.] With flagellum at each pole; appl. bacteria; amphitrichate, amphitrichic.
- amphitrocha (ămfit'rökă) n. [Gk. amphi, both; trochos, wheel.] A free-swimming annelid larva with two rings of cilia.
- amphitropous (āmfīt'röpūs) a. [Gk. amphi, both; trope, turning.] Having the ovule inverted, with hilum in middle of one side.

amphixylic,—perixylic, q.v.

- amphocyte (ăm'fosīt) n. [Gk. ampho, both of two; kytos, hollow.]
 An amphophil cell.
- amphogenic (ăm'föjen'ik) a. [Gk. ampho, both of two; -genes, producing.] Producing offspring consisting of both males and females.
- amphophil (ăm'föfil) a. [Gk. ampho, both of two; philein, to love.] Appl. cells staining with basic and acid dyes; amphochromatophil; neutrophil. n. Amphocyte.
- amphoteric (am'foter'ik) a. [Gk. amphotere, in both ways.] With opposite characters; acidic and also basic.
- amplectant (ămplěk'tănt) a. [L. amplecti, to embrace.] Clasping or winding tightly round some support, as tendrils.
- amplexicaul (ămplēk'sĭkôl) a. [L. amplecti, to embrace; caults, stem.] Clasping or surrounding the stem, as base of leaf.
- amplexus (ămplěk'sŭs) n. [L. amplexus, embrace.] Sexual embrace, in batrachians.
- ampliate (ăm'pliāt) a. [L. ampliatus, made wider.] Having outer edge of wing prominent, as in certain insects.
- amplification (ăm'plifikā'shūn) n. [L. amplificatio, enlargement.] Changes towards increased structural or functional complexity in ontogeny or phylogeny. Opp. reduction.
- ampulla (ămpool'ă, -pŭl'-) n. [L. ampulla, flask.] A membranous vesicle; dilatation of a lactiferous

- tubule beneath areola; dilated portion at one end of each semicircular canal of ear; dilatation of united common bile-duct and pancreatic duct; part of oviduct between infundibulum and isthmus; dilated portion of vas deferens at fundus of urinary bladder; terminal dilatation of rectum; internal reservoir on ring canal of watervascular system in echinoderms; terminal vesicle of sensory canals of elasmobranchs; submerged bladder of Utricularia.
- ampullaceal (ămpŭlā'seăl) a. [L. ampulla, flask.] Flask-shaped, appl. arachnid spinning glands which furnish silk for foundations, lines, and radii; appl. sensillae.
- ampullaceous (ămpūlā'sēŭs) a. [L. ampulla, flask.] Flask-shaped; appl. sensillae.
- ampullary (ămpool'ărĭ, -pŭl'-) a. [L. ampulla, flask.] Pert. or resembling an ampulla.
- ampullula (ămpool'ūlă, -pŭl-) n. [Dim. of L. ampulla, flask.] A small ampulla, as of some lymphatic vessels.
- ampyx (ăm'piks) n. [Gk. ampyx, fillet.] A transverse bar connecting the rostralia of Palaeospondylus.
- amyelinic (āmīēlin'īk) a. [Ġk. a, without; myelos, marrow.] Without myelin; appl. non-medullated or grey nerve-fibres; amyelinate.
- amygdala (ămīg'dālā) n. [L. from Gk. amygdale, almond.] Almond; one of palatal tonsils; rounded lobe at side of vallecula of cerebellum.
- amygdalin (āmig'dālin) n. [Gk. amygdale, almond.] A compound occurring in fruit kernels of bitter almonds and other Rosaceae, and producing hydrocyanic acid, glucose and benzaldehyde upon hydrolysis; C₂₀H₂₇O₁₁N.
- amylase (am'ilas) n. [L. amylum, starch.] An enzyme which converts (a) starch into dextrin, or (β) dextrin into maltose; amylolytic enzyme.
- amyloclastic,—amylolytic, q.v.

amyloid (ăm'iloid) a. [Gk. amylon, starch; eidos, form.] Starch-like.
n. Starch-like substance.

22

- amyloid bodies,—concretions found in alveoli of adult prostate gland.
- amylolytic (ăm'ilölit'ik) a. [Gk. amylon, starch; lysis, loosing.] Starch-digesting, appl. enzymes.
- amylome (am'ilom) n. [Gk. amylon, starch.] Starch-containing wood-parenchyma; layer of starch-containing cells between central cylinder and leptoids of certain moss rhizomes.
- amyloplast (ăm'ilöplăst') n. [Gk. amylon, starch; plastos, formed.] A leucoplast or colourless starch-forming granule in plants; amyloplastid.
- amylopsin (ăm'ilöp'sin) n. [Gk. amylon, starch; opson, seasoning.]
 Pancreatic amylase.
- amylose (ām'īlōs) n. [L. amylum, starch.] The substance forming starch.
- amylostatolith (ăm'îlöstăt'ölith) n. [Gk. amylon, starch; statos, stationary; lithos, stone.] A starch grain which moves under the influence of gravity in a statocyte; cf. statolith.
- amylum (ăm'ilūm) n. [L. amylum, starch.] Vegetable starch; $(C_6H_{10}O_5)_x$.
- anabiosis (an'abiō'sis) n. [Gk. ana, up; bios, life.] Resuscitation after apparent death; power of revivification, as seen in certain Tardigrada.
- anabolism (ănăb'ölizm) n. [Gk. ana, up; bole, throw.] The constructive chemical processes in living organisms, opp. katabolism.
- anabolite (ănăb'ölīt) n. [Gk. ana, up; bole, throw.] A substance participating in anabolism.
- anacrogynous (ănăkrōj'inŭs) a. [Gk. an, not; akros, apex; gyne, female.] Appl. certain liverworts in which female reproductive bodies do not arise at or near apex of shoot. Opp. acrogynous.
- anacromyoidian (anak'römioid'ian)
 a. [Gk. ana, up; akros, apex;
 mys, muscle; eidos, form.] With

- syringeal muscles attached at dorsal ends of bronchial semi-rings.
- anadromous (ănăd'römŭs) a. [Gk. ana, up; dromein, to run.] Appl. fishes which migrate from salt to fresh water annually. Opp. catadromous.
- anaerobe (ănā'erōb) n. [Gk. an, without; aer, air; bios, life.] An anaerobic organism, capable of living in absence of free oxygen. Opp. aerobe. a. Anaerobic.
- anaerobiosis (ănā'ēröbīō'sĭs) n. [Gk. an, without; aer, air; biosis, manner of life.] Existence in absence of free oxygen.
- anaesthesia (ănesthe'siă) n. [Gk. an, without; aisthesis, feeling.]
 Local or general insensibility.
- anagenesis (ăn'ăjēn'ēsis) n. [Gk. ana, again; genesis, descent.] Regeneration of tissues; progressive evolution.
- anakinetic (ăn'ăkinĕt'îk) a. [Gk. ana, up; kinein, to move.] Appl. process which restores energy; cf. katakinetic.
- anakinetomeres (ăn'ăkinē'tömērz)
 n. plu. [Gk. ana, up; kinein, to move; meros, part.] Energy-rich reactive atoms or molecules.
- anal (ā'năl) a. [L. anus, anus.]

 Pert., or situated at or near, the anus; appl. posterior median ventral fin of fishes, margin and vein of insect wing, posterior ventral scute of reptiles, etc.
- analogues (ăn'ălögz) n. plu. [Gk. analogia, proportion.] Organs of different plants or animals with like function but of unlike origin.
- analogy (ănăl'öji) n. [G. analogia, proportion.] Resemblance in function though not in structure or development.
- anamestic (ănăměs'tik) a. [Gk. ana, up; mestos, filled.] Appl. small variable bones filling spaces between larger bones of more fixed position, as in fish skulls.
- anamniote (ănăm'niōt) n. [Gk. a, not; amnion, foetal membrane.] An animal which has no amnion in embryonic life.

- anamorpha (ănămôr'fă) n. plu. [Gk. ana, backwards; morphe, form.] Larvae hatched with incomplete number of segments; cf. epimorpha.
- anamorphosis (ăn'ămôr'fōsis) n. [Gk. ana, throughout; morphosis, shaping.] Evolution from one type to another through a series of gradual changes.
- anandrous (ănăn'drŭs) a. [Gk. a, without; aner, male.] Without stamens.
- anangian (ănăn'jiăn) a. [Gk. a, without; anggeion, vessel.] Appl. worms without a vascular system.
- anantherous (ăn'ănthërus) a. [Gk. a, without; antheros, flowering.] Without anthers.
- ananthous (ănăn'thūs) a. [Gk. a, without; anthos, flower.] Not flowering; without inflorescence.
- anaphase (an'afaz) n. [Gk. ana, up; phasis, appearance.] A stage in mitosis during divergence of daughter chromosomes; the stages of mitosis up to division of chromatin into chromosomes; cf. kataphase.
- anaphylaxis (ăn'ăfilăk'sis) n. [Gk. ana, up; phylax, guard.] Condition of being hypersensitive to a serum or foreign protein, caused by first or sensitising dose.
- anaphyte (ăn'ăfit) n. [Gk. ana, up; phyton, plant.] Transverse segment of a shoot; an internode.
- anaplasia (ănăplā'ziă) n. [Gk. ana, again; plassein, to form.] Undifferentiation; reversion to a less differentiated structure.
- anaplast (ăn'ăplăst) n. [Gk. ana, up; plastos, formed.] A leucoplastid; anaplastid.
- anapleurite (an'aploor'it) a. [Gk. ana, up; pleura, side.] Upper thoracic pleurite, as in certain Thysanura.
- anapophysis (ăn'ăpŏf'īsīs) n. [Gk. ana, up; apo, from; physis, origin.]
 A small dorsal projection rising near transverse process in lumbar vertebrae.
- anapsid (ănăp'sid) a. [Gk. ana, up;

- apsis, arch.] With skull wholly imperforate or completely roofed over; stegocrotaphic.
- anaptychus (ănăp'tikūs) n. [Gk. ana, throughout; ptyche, plate.]
 Aptychus or operculum consisting of a single plate, as in certain ammonites; cf. synaptychus.
- anarthrous (ănâr'thrus) a. [Gk. a, without; arthron, joint.] Having no distinct joints.
- anaschistic (ăn'ăskis'tik) a. [Gk. ana, up to; schistein, to split.] Appl. type of tetrads which divide twice longitudinally in meiosis; cf. diaschistic.
- anastates (ăn'ăstāts) n. plu. [Gk. ana, up to; statos, standing.] Various materials that arise owing to metabolism in a cell, in formation of complex from simple substances. Opp. katastates.
- anastomosis (ănăs'tömō'sis) n. [Gk. ana, up to; stoma, mouth.] Union of ramifications of leaf-veins; union of blood-vessels arising from a common trunk; union of nerves; fine threads joining chromonemata in resting nucleus; formation of a network or anastomotic meshwork.
- anastral (ănăs'trăl) a. [Gk. an, not; aster, star.] Appl. type of mitosis without aster-formation.
- anatomy (ănăt'ömi) n. [Gk. ana, up; tome, cutting.] The science which treats of the structure of plants and of animals, as determined by dissection; usually, human anatomy.
- anatoxin,—toxoid, q.v.
 anatrepsis (ănătrēp'sīs) n. [Gk.
 anatrepein, to turn over.] Stage
 of increasing movement in blasto-
- of increasing movement in blastokinesis.

 anatriaene (ăn'ătrī'ēn) n. [Gk. ana,
- up; triaina, trident.] Triaene with backwardly directed branches. anatropous (ănăt'röpūs) a. [Gk. anatrope, overturning.] Inverted, appl. ovules with hilum and micropyle close together and chalaza at other end; anatropal.
- anaxial (ănăk'sĭāl) a. [Gk. a, without; axis, axle.] Having no distinct axis; asymmetrical.

24

ancestrula (ănsĕs'troolă) n. [L. antecedere, to go before.] First zooecium of polyzoan colony.

anchor (ăng'kör) n. [L. ancora, anchor.] Anchor-shaped spicule found in skin of Holothuria.

anchylosis (ăng'kilō'sis) n. [Gk. angchein, to press tight.] Union of two or more bones or hard parts to form one part, e.g. of bone to bone, or tooth to bone; ankylosis.

anconeal (ăngkō'neal) a. [Gk. angkon, elbow.] Pert. the elbow.

anconeus (ăng'konē'ŭs) n. [Gk. angkon, elbow.] Small extensor muscle situated over elbow; anconaeus.

andrase (ăn'drās) n. [Gk. aner, male.] A male-determining factor in form of an enzyme or hormone.

andrin (ăn'drin) n. [Gk. aner, male.] The testicular androgens.

androconia (ăn'drökō'niă) n. plu.
[Gk. aner, male; konia, dust.]
Modified wing-scales producing a sexually attractive scent in certain male butterflies.

androcyte (ăn'drosīt) n. [Gk. aner, male; kytos, hollow.] A cell arising by growth from an androgonium and giving rise to antherozoid.

androdioecious (ān'drödīē'siŭs) a. [Gk. aner, male; dis, two; oikos, house.] Having male and hermaphrodite flowers on different plants.

androecium (ăndrē'siŭm) n. [Gk. aner, male; oikos, house.] Male reproductive organs of a plant; stamens taken collectively.

androgametangium (ăn'drögăm'ētăn'jiŭm) n. [Gk. aner, male;
gametes, spouse; anggeion, vessel.]
A structure producing male sexual
cells; antheridium.

androgen (ăn'dröjēn) n. [Gk. aner, male; genos, descent.] A male hormone; a masculinising substance.

androgenesis (ăn'dröjen'esis) n. [Gk. aner, male; genesis, descent.]
Development of egg furnished with paternal chromosomes only; male parthenogenesis.

androgenetic (ăn'dröjenet'ik)

[Gk. aner. male; genesis, descent.] Having paternal chromosomes only.

androgenic (ăn'dröjěn'ik) a. [Gk. aner, male; gennaein, to produce.] Stimulating male characters; masculinising; appl. hormones; appl. tissue capable of elaborating an androgenic hormone; androgenous, q.v.

androgenous (ăndroj'enus) a. [Gk. aner, male; genos, descent.] Producing only male offspring.

androgonidia (ăn'drogonid'iă) n. plu. [Gk. aner, male; gonos, offspring; idion, dim.] Male sexual elements formed after repeated divisions of parthenogonidia of Volvox.

androgonium (ăn'drögō'nĭŭm) n.
[Gk. aner, male; gonos, offspring.]
An early stage in formation of

sperm-cells of plants.

androgynal (androj'inăl) a. [Gk. aner, male; gyne, female.] Hermaphrodite; bearing both staminate and pistillate flowers in the same infloresence; with antheridium and oogonium on the same hypha; androgynous.

androgynary (åndröj'inări) a. [Gk. aner, male; gyne, female.] Having flowers with stamens and pistils

developed into petals.

androgyne (ăn'dröjīn, ăn'drögīn) a.,
n. Hermaphrodite.

androgynism (ăndrŏj'ĭnĭzm) n. [Gk. aner, male; gyne, female.] The condition of bearing both stamens and pistils; hermaphroditism.

andromerogony (ăn'dröměrŏg'öni)
 n. [Gk. aner, male; meros, part; gone, generation.] The development of an egg fragment with only paternal chromosomes.

andromonoecious (ăn'drömönē'sĭŭs)
a. [Gk. aner, male; monos, alone; oikos, house.] Having male and hermaphrodite flowers on the same plant.

andropetalous (ăn'dröpět'ălŭs) a. [Gk. aner, male; petalon, leaf.]

Having petaloid stamens.

androphore (ăn'dröfōr) n. [Gk. aner, male; phora, carrying.] Stalk supporting androecium or stamens; stalk carrying male gonophores in Siphonophora.

- androphyll (ăn'dröfil) n. [Gk. ancr, male; phyllon, leaf.] The leaf bearing microspores; the microsporophyll.
- androsome (ăn'drösōm) n. [Gk. aner, male; soma, body.] A male-limited chromosome.
- androsporangium (ăn'dröspörăn'jium) n. [Gk. aner, male; sporos, seed; anggeion, vessel.] A sporangium containing androspores.
- androspore (ăn'dröspōr) n. [Gk. aner, male; sporos, seed.] An asexual zoospore which gives rise to a male dwarf plant; male spore; microspore; pollen grain.
- androsterone (ăn'dröstē'rōn) n. [Gk. aner, male; stear, suet.] Male hormone, present in adrenal cortex, obtained from urine; C₁₉H₃₀O₂.
- androtype (ăn'drotīp) n. [Gk. aner, male; typos, pattern.] Type specimen of the male of a species.
- anelectrotonus (ăn'ēlēktrötō'nŭs, ăn'ēlēktrōt'önŭs) n. [Gk. ana, up; elektron, amber; tonos, tension.] Decrease in irritability of a nerve under influence of a non-polarising electric current.
- anellus (ăněl'ús) n. [L. anellus, little ring.] A small ring-shaped or triangular plate supported by valves and vinculum, in Lepidoptera.
- anemochorous (ǎněmókō'rǔs) a. [Gk. anemos, wind; chore, place.] Dispersed by wind; with seeds so dispersed.
- anemophilous (ăněmôf'īlūs) a. [Gk. anemos, wind; philein, to love.] Wind-pollinated.
- anemophily (anemof'ili) n. [Gk. anemos, wind; philein, to love.] Plant-fertilisation by agency of wind.
- anemoplankton (ănĕm'oplăngk'tŏn)
 n. [Gk. anemos, wind; plangktos, wandering.] Wind-borne organisms and living particles; aeroplankton, q.v.
- anemosporic (ăněm'öspŏr'ik) a. [Gk. anemos, wind; sporos, seed.] Having spores or seeds disseminated by air currents.
- anemotaxis (ăněm'ötăk'sis) n. [Gk. anemos, wind; taxis, arrange-

- ment.] Directed movement in response to air currents.
- anemotropism (ăněmot'röpĭzm) n.
 [Gk. anemos, wind; trope, turn.]
 Orientation of body, or plant
 curvature, in response to air
 currents.
- anencephaly (ăněnkěťálĭ, -sěť-) n. [Gk. an, not; engkephalon, brain.] Condition of having no brain.
- anenterous (ăněn'tërus) a. [Gk. an, without; enteron, gut.] Having no alimentary tract; anenteric.
- aner (ān'ér, ănār) n. [Gk. aner, male.] The male of insects, especially of ants.
- anestrum,—anoestrus, q.v.
- aneucentric (ānūsēn'trīk) a. [Gk. a. without; eu, well; kentron, centre.] Acentric and dicentric, resulting from translocation involving centromere of a chromosome.
- aneuploid (ăn'ūploid) a. [Gk. a, without; eu, well; aploos, onefold.] Having fewer or more chromosomes than an exact multiple of the haploid number, opp. euploid.
- aneurine (ănū'rīn) n. [Gk. a, without; neuron, nerve.] Vitamin B₁, the anti-beri-beri factor in yeast, legumes, cereals, and other foods; aneurin; thiamine (U.S.A.); C₁₂H₁₈ON₄SCl₂.
- aneuronic (anūron'ik) a. [Gk. a, without; neuron, nerve.] Without innervation; appl. chromatophores controlled by hormones.
- anfractuose (ănfrăk'tūōs) a. [L. anfractus, bending.] Wavy, sinuous.
- angienchyma (ăn'jičng'kimă) n. [Gk. anggeion, vessel; engchein, to pour.] Vascular tissue.
- angioblast (ăn'jiöbläst) n. [Gk. anggeion, vessel; blastos, bud.]
 One of cells from which lining of blood-vessels is derived; vasoformative cell.
- angiocarpic (ăn'jiökâr'pik) a. [Gk. anggeion, vessel; karpos, fruit.] Having fruit enclosed; angiocarpous; appl. fungi. Opp. gymnocarpic.

(ăn'jiŏl'öji) n. [Gk. anggeion, vessel; logos, discourse.] Anatomy of blood and lymph vascular systems.

angiospermous (ănjiöspēr'mus) a. [Gk. anggeion, vessel; sperma, seed.] Having seeds in a closed

case, the ovary.

angiosporous (ănjiŏs'pörŭs) a. [Gk. anggeion, vessel; sporos, seed.] Having spores contained in a theca or spore capsule.

angiostomatous (ăn'jiöstom'ătus) a. [Gk. anggeion, vessel; mouth.] Narrow-mouthed, appl. an order of molluscs, and to a sub-order of snakes, with nondistensible mouth.

angiotonin (anjiot'onin) n. anggeion, vessel; tonos, tension.]

Substance in circulating blood, formed by reaction between hypertensinogen elaborated in the liver, and renin, causing constriction of

arterioles; hypertensin.

ångström (ông'strëm) n. $[A.\ J.$ Angström, Swedish physicist.] One ten millionth part of a millimetre, symbol A.

angular (ăng'gūlăr) n. [L. angulus, corner.] A membrane bone of lower jaw in most vertebrates. a. Having, or *pert*., an angle; *appl*. leaf originating at forking of stem, as in many ferns; appl. collenchyma with cell-walls thickened in the angles of the cells.

angulosplenial (ăng'gūlösplē'nĭāl) n. [L. angulus, corner; splenium, patch.] Bone forming most of lower and inner part of mandible

in Amphibia.

angulus (ăng'gulus) n. [L. angulus, angle.] An angle, as that formed by junction of manubrium and body of sternum (angle of Louis).

angustifoliate (ănggūs'tĭfō'lĭāt) a. [L. angustus, narrow; folium, leaf.]

With narrow leaves.

angustirostrate (ănggŭs'tĭrŏs'trāt) a. [L. angustus, narrow; rostrum, beak.] With narrow beak or snout.

anholocyclic (ăn'hŏlösīk'lĭk) a. [Gk.

an, not; holos, whole; kyklos, circle.] Pert. alternation of generations with suppression of sexual part of cycle; permanently parthenogenetic.

anidian (ănid'iăn) a. [Gk. an, not; eidos, form. Formless; blastoderm without apparent em-

bryonic axis.

pole,—the animal upper, rapidly segmenting, portion of a telolecithal egg. Opp. vegetal pole.

ſGk.

anion (ăn'iŏn, ăn'iŏn) n. [Gk. ana, up; ienai, to go.] A negativelycharged particle or ion which moves up towards the anode or positive pole.

anisocarpous (ănīsökâr'pus) a. [Gk. anisos, unequal; karpos, fruit.] Having number of carpels less than that of other floral whorls.

anisocercal (ănīsösēr'kāl) a. anisos, unequal; kerkos, tail.] With lobes of tail-fin unequal.

anisochela (ănīsökē'lă) n. anisos, unequal; chele, claw.] A chela with the two parts unequally developed.

anisodactylous (ănisödăk'tilus) a. [Gk. anisos, unequal; daktylos, finger.] Having unequal toes, three toes forward, one backward.

anisogamete (ăn'isögămēt) n. [Gk. anisos, unequal; gametes, spouse.] One of two conjugating gametes differing in form or size.

anisogamous (ănisŏg'ămŭs) a. [Gk. anisos, unequal; gamos, marriage.] Appl. differentiated gametes or

conjugating bodies.

anisogamy (ănisŏg'ămi) n. anisos, unequal; gametes, spouse.] Conjugation between sharply differentiated gametes; heterogamy.

anisognathous (ăn'īsŏg'nāthūs) a. [Gk. anisos, unequal; gnathos, jaw.] With jaws of unequal width; having teeth in upper and lower jaws unlike.

anisomerous (ănisom'erus) a. [Gk. anisos, unequal; meros, part.] Having unequal numbers of parts in floral whorls.

anisophylly (ăn'fisöfil'i) n. [Gk. anisos, unequal; phyllon, leaf.]
Condition of having leaves of two or more sizes or shapes, as in some conifers and aquatic plants.

anisopleural (ănisoploo'răl) a. [Gk. anisos, unequal; pleura, side.]

Asymmetrical bilaterally.

anisoploid (ăn'isöploid) a. [Gk. anisos, unequal; aploos, onefold; eidos, form.] With an odd number of chromosome sets in somatic cells. n. An anisoploid individual.

anisopogonous (ănisöpōg'önŭs) a. [Gk. anisos, unequal; pogon, beard.] Unequally webbed, with reference to feathers.

anisopterus (ănisop'terus) a. [Gk. anisos, unequal; pteron, wing.] Unequally winged; appl. seeds.

anisospore (ănīsöspōr) n. [Gk. anisos, unequal; sporos, seed.] A dimorphic spore, the sexes differing in size.

anisostemonous (ănisostem'onus) a. [Gk. anisos, unequal; stemon, spun thread.] Having the number of stamens unequal to the number of parts in other floral whorls; having stamens of unequal size.

anisotropic (ănisotrop'ik) a. [Gk. anisos, unequal; trope, turn.] Appl. eggs with predetermined axis or axes; exhibiting anisotropy; doubly refracting, appl. dark bands of voluntary muscle fibre. Opp. isotropic.

isotropic

ankylosis,—anchylosis, q.v.

ankyroid (ăng'kiroid) a. [Gk. agkyra, hook; eidos, form.] Hook-shaped. anlage (ân'lâgë) n. [Ger. Anlage, predisposition.] The first structure or cell group indicating development of a part or organ; inception; primordium; ébauche.

annectent (ăněk'tënt) a. [L. annectere, to bind together.] Linking, appl. intermediate species or genera.

annelid (ăn'ēlĭd) a. [L. annulus, ring; Gk. eidos, form.] Constructed of ring-like segments, as ringed worms; pert. Annelida.

annotinous (ănnotinus) a. [L. annus,

year.] A year old; appl. growth during the previous year.

annual (ăn'ūāl) a. [L. annus, year.] Appl. structures or features that are marked off or completed yearly; living for a year only.

annual ring,—one of the rings, seen in transverse sections of dicoty-

in transverse sections of dicotyledons, indicating the secondary growth during a year; growth ring

of bivalve shells.

annular (ăn'ūlăr) a. [L. annulus, ring.] Ring-like; appl. certain ligaments of wrist and ankle; appl. (orbicular) ligament encircling head of radius and attached to radial notch of ulna; appl. certain lamina or sternal plates in ants; appl. certain vessels in xylem, owing to ring-like thickenings in their interior; appl. bands formed on inner surface of cell-wall.

annulate (ăn'ūlāt) a. [L. annulus, ring.] Ring-shaped; composed of ring-like segments; having colour arranged in ring-like bands or annuli.

annulus (ăn'ūlŭs) n. [L. annulus, ring.] Any ring-like structure; special ring in fern sporangium, by action of which sporangium bursts; remains of veil in mushrooms; ring of cells in moss capsule whose rupture causes opening; circular groove for transverse flagellum in Dinoflagellata; ring of annelid; fourth digit of hand.

anococcygeal (ā'nököksīj'ēăl) a. [L. anus, anus; coccyx; Gk. kokkyx, cuckoo.] Pert. region between coccyx and anus; appl. body of fibrous and muscular tissue, nerves,

anoestrus (ănē'strūs) n. [Gk. an, not; oistros, gad-fly.] The non-breeding period; period of absence of sexual urge; anoestrum; cf. dioestrus.

anomaly (ănom'ăli) n. [Gk. anomalos, uneven.] Any departure from type characteristics.

anomophyllous (ăn'ŏmöfĭl'ŭs) a. [Gk. anomos, lawless; phyllon, leaf.] With irregularly placed leaves.

anorganology (ăn'ôrgănôl'öji) n.
|Gk. a, not: organon, instrument;
logos, discourse.] Study of nonliving things; abiology.

anorthogenesis (ān'ôrthöjĕn'ēsĭs) n. [Gk. an, not; orthos, straight; genesis, descent.] Evolution manifesting changes in direction of adaptations, owing to preadaptation; 'zigzag' evolution.

anorthospiral (ăn'ôrthöspī'răl) a. [Gk. an, not; orthos, straight; speira, coil.] Relationally coiled, spirals not interlocking; paranemic. Opp. orthospiral, plectonemic.

anosmatic (ănŏsmăt'ĭk) a. [Gk. a, without; osme, smell.] Having no sense of smell; anosmic.

anosmia (anos'mia) n. [Gk. a, without; osme, smell.] Absence or loss of sense of smell.

anoxybiotic (ănŏk'sibīot'ik) a. [Gk. a. not; oxys, sharp; biotos, means of life.] Capable of living in absence of oxygen; anaerobic.

ansa (ăn'să) n. [L. ansa, handle.] Loop, as of certain nerves.

anserine (ăn'sērĭn) n. [L. anser, goose.] A constituent of bird's muscle; C₁₀H₁₆O₃N₄.

ansiform (ăn'sifôrm) a. [L. ansa, handle; forma, shape.] Loopshaped; looped; appl. outer cytoplasm in cerebro-spinal ganglia.

antagonist (ăntăg önist) n. [Gk. antagonistes, adversary.] A muscle acting in opposition to the action produced by a prime mover or agonist; an antihormone, q.v.

antambulacral (ănt'ămbūlā'krāl) a. [Gk. anti, against; L. ambulare, to walk.] Not situated on the ambulacral area; abactinal, q.v.

antapex (ăntăp'ēks) n. [Gk. anti, opposite; L. apex, tip.] Tip of hypocone in Dinoflagellata.

antapical (ăntăp'îkăl) a. [Gk. anti, opposite; L. apex, tip.] At or pert. antapex; pert. region opposite apex.

antebrachium (ăn tebra kium) n. [L. ante, before; brachium, arm.] The fore-arm, or corresponding portion of a fore-limb.

anteclypeus (ăn'tëklīp'ëŭs) n. [L. ante, before; clypeus, shield.] Anterior portion of clypeus when differentiated by suture; cf. post-clypeus.

antecosta (ăntekos'tă) n. [L. ante, before; costa, rib.] Internal ridge of tergum, for attachment of intersegmental muscles in insects, extended to phragma in alar segments.

antecubital (ăn'tēkū'bĭtăl) a. [L. ante, before; cubitus, elbow.] Anterior to the elbow.

antedorsal (ăn'tēdôr'săl) a. [L. ante, before; dorsum, back.] Situated in front of dorsal fin in fishes.

antefrons (ăn'tēfronz) n. [L. ante, before; frons, forehead.] The portion of frons anterior to antennary base line in certain insects.

antelabrum (ăntelă'brum) n. [L. ante, before; labrum, lip.] The anterior portion of insect labrum when differentiated.

antemarginal (ăn'tëmâr'jīnăl) a. [L. ante, before; margo, edge.] Appl. sori of ferns when they lie within margin of frond.

antenna (ăntěn'ă) n. [L. antenna, sail-yard.] A jointed feeler on head of various Arthropoda.

antennary (ăntěn'ări) a. [L. antenna, sail-yard.] Like, or pert., or situated near an antenna; antennal.

antennifer (ăntěn'ifēr) n. [L. antenna, sail-yard; ferre, to carry.] Socket of antenna in arthropods; projection on rim of antennal socket, acting as a pivot, in myriopods.

antennule (ănten'ūl) n. [L. dim. from antenna.] A small antenna or feeler, specifically the first pair of antennae in Crustacea.

anteposition (ăn'tëpözish'ŭn) n. [L. ante, before; ponere, to place.] Superposition of whorls in a flower typically alternating.

anterior (ăntē'riŏr) a. [L. anterior, former.] Nearer head end; ventral in human anatomy; facing outwards from axis; previous.
anterolateral,—ventrolateral.

- antesternite (ăn'tëstër'nīt) n. [1..
 ante, before; sternum, breast-bone.]
 Anterior sternal sclerite of insects;
 basisternum, eusternum.
- anthela (anthe'la) n. [Gk. anthein, to bloom.] The cymose inflorescence of the rush family.

anthelix,—antihelix, q.v.

- anther (ăn'thër) n. [Gk. antheros, flowering.] The part of stamen which produces pollen.
- antherid,-antheridium.
- antheridia, -plu. of antheridium.
- antheridial cell,—the larger of two cells derived from a microspore and giving rise to an antheridium, or to a cell representing an antheridium.
- antheridiophore (ănthërid'iöfor) n. [Gk. anthos, flower; idion, dim.; pherein, to bear.] A gametophore bearing antheridia.
- antheridium (ănthērid'iŭm) n. [Gk. anthos, flower; idion, dim.] An organ or receptacle in which male sexual cells are produced in many cryptogams; male gametangium; cluster of microgametes, as in certain Flagellata.
- antherozoids (ăn'thërözō'idz), antherozoids (ăn'thërözō'oidz) n. plu. [Gk. anthos, flower; zoon, animal; eidos, form.] Male sexual cells in antheridia.
- anthesis (anthe'sis) n. [Gk. anthos, flower.] Stage or period at which flower-bud opens; flowering; period of flowering.
- anthoblast (ăn'thöblăst) n. [Gk. anthos, flower; blastos, bud.] In Madreporaria, a young sessile polyp producing anthocyathus.
- anthocarpous (ăn'thökâr'pŭs) a. [Gk. anthos, flower; karpos, fruit.] Appl. aggregated fruits, products of fusion of several flowers, as sorosis and syconus.
- anthocaulis (ăn'thökôl'is) n. [Gk. anthos, flower; L. caulis, stem.]
 The pedicle of a late trophozooid stage of madrepore development.
- anthocodia (ăn'thökō'dīā) n. [Gk. anthos, flower; kodeia, head.] The distal portion of a zooid bearing mouth and tentacles, in Alcyonaria.

- anthocyanin (ăn'thösī'ānĭn) n. [Gk. anthos, flower; kyanos, dark blue.]
 One of the blue or violet pigments of flowers, leaves, and stems.
- anthocyathus (ăn'thösi'ăthŭs) n. [Gk. anthos, flower; kyathos, cup.] The discoid crown of trophozooid stage in madrepore development.
- anthodium (ănthō'dĭŭm) n. [Gk. anthos, flower; eidos, form.]
 Capitulum or head of Compositae.
- anthogenesis (ăn'thöjěn'ēsīs) n. [Gk. anthos, flower; genesis, descent.] In certain aphids, production of both males and females by asexual forms.
- anthophilous (ănthôf'ilūs) a. [Gk. anthos, flower; philein, to love.] Attracted by flowers; feeding on flowers.
- anthophore (ăn'thôfôr) n. [Gk. anthos, flower; pherein, to bear.] Elongation of thalamus between calyx and corolla.
- anthophyte (ăn'thofit) n. [Gk. anthos, flower; phyton, plant.] A flowering plant; phaenogam, phanerogam, spermatophyte.
- anthostrobilus (ăn'thöstrob'îlūs) n. [Gk. anthos, flower; strobilos, fircone.] Fructification or flower of certain cycads.
- anthotaxis (ăn'thötăk'sĭs) n. [Gk. anthos, flower; taxis, arrangement.]
 Arrangement of flowers on an axis.
- anthoxanthin (ăn'thözăn'thĭn) n. [Gk. anthos, flower; xanthos, yellow.] A yellow pigment of flowers.
- anthropeic (ănthrope'îk) a. [Gk. anthropeios, by human means.]

 Due to influence of man.
- anthropogenesis (ăn'thrôpöjĕn'ĕsĭs)
 n. [Gk. anthropos, man; genesis, descent.] The ontogenesis and phylogenesis of man; descent of man.
- anthropogenetic (ăn'thrōpöjěnět'ík)
 a. [Gk. anthropos, man; genesis,
 descent.] Pert. anthropogenesis.
- anthropogenic (ăn'thrôpöjěn'ik) a. [Gk. anthropos, man; genos, descent.] Produced or caused by man.

ANT-

- anthropoid (ăn'thröpoid) a. [Gk. anthropos, man; eidos, form.] Resembling man; appl. tailless apes.
- anthropology (ăn'thröpŏl'öji) n. [Gk. anthropos, man; logos, discourse.] The natural history of man.
- anthropometry (ăn'thröpŏm'ëtrī) n.
 [Gk. anthropos, man; metron, measure.] That part of biology dealing with proportional measurements of parts of the human body.
- anthropomorphous (ăn'thröpömôr'fǔs) a. [Gk. anthropos, man;
 morphe, shape.] Resembling man.
- anthropotomy (ăn'thröpŏt'omi) n. [Gk. anthropos, man; temnein, to cut.] Human anatomy.
- antiae (ăn'tiē) n. plu. [L. antiae, forelock.] Feathers at base of bill-ridge of some birds.
- anti-apex,—lower end of axis, as in rootless plants.
- antiauxin (ăn'tiôks'în) n. [Gk. anti, against; auxein, to grow.] An organic compound which regulates or inhibits growth stimulation by auxins.
- antibiosis (ăn'tibiō'sis) n. [Gk. anti, against; bios, life.] Antagonistic association of organisms, as by production of harmful compounds.
- antiblastic (antiblas'tik) a. [Gk. anti, against; blastos, bud.] Appl. immunity due to forces which inhibit growth of invading organism.
- antibody (ăn'tibŏd'i) n. [Gk. anti, against; A.S. bodig, body.] Any substance formed in blood which reacts with a specific antigen, or antagonises toxins.
- antibrachial (ăn'tibrā'kiăl) a. [Gk. anti, against; L. brachium, arm.] Pert. forearm; appl. fascia, muscles, vein, nerves; also antebrachial.
- antibrachium,—antebrachium, q.v. anticipation (ăn'tisipā'shūn) n. [L. ante, before; capere, to take.] The manifestation of a condition or disease at a progressively earlier age in successive generations.

- anticlinal (ăn'tiklī'năl) a. [Gk. anti, against; klinein, to slope.] Appl. line of division of cells at right angles to surface of apex of a growing point; in quadrupedă, appl. one of lower thoracic vertebrae with upright spine towards which those on either side incline.
- anticoagulin (ăn'tĭkōāg'ūlĭn) n. [Gk. anti, against; L. coagulum, rennet.]
 A substance which prevents coagulation of drawn blood, as hirudin.
- anticryptic (ăn'tikrip'tik) a. [Gk. anti, against; kryptos, hidden.] Appl. protective coloration facilitating attack.
- anticubital,—antecubital, q.v.
- antidiuretic (ān'tidīūrĕt'ĭk) a. [Gk. anti, against; dia, through; ouron, urine.] Reducing the volume of urine; appl. a hormone of posterior lobe of pituitary gland.
- antidromic (ăntid'romik) a. [Gk. anti, against; dromos, running.] Contrary to normal direction; appl. conduction of impulse along axon towards body of nerve cell; antidromous; appl. stipules with fused outer margins.
- antidromy (antid'römi) n. [Gk. anti, against; dromos, running.] Condition of spiral phyllotaxis with genetic spiral changing direction after each cycle.
- anti-enzyme (ăn'tiĕn'zim) n. [Gk. anti, against; en, within; zyme, leaven.] A substance which retards or stops enzyme activity.
- antigen (ăn'tijěn) n. [Gk. anti, against; genos, birth.] Substance which causes a series of physiologico-chemical changes resulting in formation of antibodies.
- antigeny,—sexual dimorphism.
- antihelix (ăn'tihē'līks) n. [Gk. anti, against; helix, a convolution.]

 The curved prominence in front of helix of ear.
- antihormones (ăn'tihôr'monz) n. plu. [Gk. anti, against; hormaein, to excite.] Substances which prevent the effect of hormones; chalones, q.v.

- antimeres (ăn'timērz) n. plu. [Gk. anti, opposite; meros, part.] Corresponding parts, as left and right limbs, of a bilaterally symmetrical animal; a series of equal radial parts of a radially symmetrical animal.
- antineuritic (ăn'tĭnūrit'ĭk) a. [Gk. anti, against; neuron, nerve.] Appl. vitamin B₁, lack of which causes polyneuritis.
- antipepsin (ăn'tipep'sin) n. [Gk. anti, against; pepsis, digestion.] A stomach secretion which prevents action of pepsin on tissue proteins.
- antiperistalsis (ăn'tiperistăl'sis) n. [Gk. anti, against; peri, around; stalsis, contraction.] Reversed peristalsis; peristaltic action in posteroanterior direction.
- antipetalous (ăn'tĭpēt'ālŭs) a. [Gk. anti, opposite; petalon, petal.] With stamens opposite petals.
- antipodal (ăntip'ödāl) a. [Gk. anti, against; pous, foot.] Appl. group of three cells at chalazal end of embryo-sac; appl. cone of astral rays opposite spindle fibres.
- antipygidial (än'tipijid'iăl) a. [Gk. anti, against; pygidion, narrow rump.] Appl. bristles of seventh abdominal segment which extend to pygidium, in fleas.
- antirachitic (ăn'tirăkit'ik) a. [Gk. anti, against; rhachis, spine.] Appl. vitamin D, lack of which causes rickets.
- antiscorbutic (ăn'tiskôrbū'tik) a. [Gk. anti, against; L.L. scorbutus, scurvy.] Appl. vitamin C, lack of which causes scurvy.
- antisepalous (ăn'tisép'ălŭs) a. [Gk. anti, opposite; F. sépale, from L. separare, to separate.] With stamens opposite sepals.
- antiseptic (an'tisep'tik) a. [Gk. anti, against; sepsis, putrefaction.] Preventing putrefaction. n. A substance which destroys harmful micro-organisms.
- antispadix (ăn'tispā'diks) n. [Gk. anti, against; spadix, palm branch.]
 A group of four modified tentacles in internal lateral lobes of Nautilus.

- antisquama (ăn'tiskwā'mă) n. [Gk. anti, against; L. squama, scale.] Basal lobe next squama of insect wing; squama alaris or antitegula.
- antistyle (ăn'tistīl) n. [Gk. anti, against; L. stilus, pricker.] Basal projection of stylifer in certain insects.
- antitegula (ăn'titeg'ūlă) n. [Gk. anti, against; L. tegula, tile.]
 Antisquama, q.v.
- antithetic (ăn'tithět'îk) a. [Gk. antithesis, opposition.] Appl. alternation of diploid and haploid generations, or of sporophyte and gametophyte generations.
- antithrombin (ăn'tĭthrom'bin) n. [Gk. anti, against; thrombos, clot.] A substance, as formed in liver, which prevents clotting of blood.
- antitoxin (ăn'titok'sin) n. [Gk. anti, against; toxikon, poison.] A substance or antibody which neutralises or binds a toxin.
- antitragus (ăn'titrā'gūs) n. [Gk. anti, against; tragos, goat.] Prominence opposite tragus of external ear.
- antitrochanter (ăn'titrökăn'tër) n. [Gk. anti, against; trochanter, a runner.] In birds, an articular surface on ilium against which trochanter of femur plays.
- antitropic (ăn'tītrop'īk) a. [Gk. anti, against; trope, turn.] Turned or arranged in opposite directions; arranged to form bilaterally symmetric pairs, as ribs of opposite sides; cf. syntropic.
- antitropous (ăntit'röpüs) a. [Gk. anti, against; trope, turn.] Inverted; appl. embryos with radicle directed away from hilum; antitropal.
- antlia (ănt'liă) n. [L. antlia, pump.]
 The spiral suctorial proboscis of Lepidoptera.
- antorbital (ăntôr'bităl) a. [L. ante, before; orbis, circle.] Situated in front of orbit; appl. bone, cartilage, process.
- antrum (ăn'trum) n. [L. antrum, cavity.] A cavity or sinus; e.g., maxillary sinus, cavity of pylorus.

- anurous (ănū'rūs) a. [Gk. a, without; oura, tail.] Tailless.
- anus (ā'nŭs) n. [L. anus, anus.]
 Posterior opening of alimentary canal.
- aorta (āôr'tā) n. [Gk. aorte, the great artery.] The great trunk artery which carries pure blood to the body through arteries and their branches.
- aortic (āôr'tik) a. [Gk. aorte, the great artery.] Pert. aorta; appl. arch, hiatus, isthmus, lymph glands, semilunar valves, etc.
- aortic bodies,—two small masses of chromaffin cells in a capillary plexus, one on each side of foetal abdominal aorta, being part of system for controlling oxygen content and acidity of blood; Zuckerkandl's bodies.
- apatetic (ăp'ătē'tĭk) a. [Gk. apatetikos, fallacious.] Appl. misleading coloration.
- aperispermic(ăper'isper'mik)a. [Gk.
 a, without; peri, around; sperma,
 seed.] Appl. seeds without nutritive
 tissue.
- apertura piriformis,—anterior nasal aperture of skull.
- apetalous (apet'alus) a. [Gk. a, without; petalon, petal.] Without petals; monochlamydeous.
- apex (ã'pēks) n. [L. apex, summit.]
 Tip or summit, as of lungs, heart,
 nose; styloid process of fibula;
 tip of epicone in Dinoflagellata;
 wing tip in insects.

aphanipterous (ăf'ănĭp'tërŭs)a. [Gk. aphanes, unseen; pteron, wing.]
Apparently without wings.

- apheliotropism (ăfē'liöt'röpizm) n. [Gk. apo, away; helios, sun; trope, turn.] Tendency to turn away from light; aphototropism.
- aphlebia (ăflěb'yă) n. [Gk. a, without; phleps, vein.] Lateral outgrowth from base of frond-stalk in certain ferns.
- aphodal (ăf'ödăl) a. [Gk. apo, away; hodos, path.] Appl. type of canal system in sponges.
- aphodus (ăf ödus) n. [Gk. aphodos, departure.] The short tube leading

- from flagellate chamber to excurrent canal in a type of canal system in sponges.
- aphotic (ăfō'tik) a. [Gk. a, without; phos, light.] Pert. absence of light; appl. zone of deep sea where daylight fails to penetrate. Opp. photic.
- aphyllous (āfīl'ūs) a. [Gk. a. without; phyllon, leaf.] Without foliage leaves.
- aphylly (ăfil'i) n. [Gk. a, without; phyllon, leaf.] Suppression or absence of leaves.
- apical (ăp'ikăl) a. [L. apex, summit.] At tip or summit; pert. distal end; appl. cell at tip of growing point; appl. meristem; appl. style arising from summit of ovary; appl. dominance, of terminal bud; appl. aboral plates of echinoderms; appl. neural plate of trochophore and tornaria.
- apiculate (ăpĭk'ūlāt) a. [Dim. of L. apex, summit.] Forming abruptly to a small tip, as leaf.
- apiculus (ăpik'ūlūs) n. [Dim. of L. apex, summit.] A small apical termination, as in some protozoa, or of certain spores; reflexed portion of antennal club, in some Lepidoptera.
- apilary (ăpil'ări) a. [Gk. a, not; pilos, felt cap.] Having upper lip wanting or suppressed in corolla.
- apitoxin (āpitok'sin) n. [L. apis, bee; Gk. toxikon, poison.] Main toxic fraction of bee venom.
- apituitarism (ăp'îtū'îtărizm) n. [L. a, away; pituita, phlegm.] Absence or deficiency of pituitary gland secretion; hypohypophysism.
- aplacental (ắp lasen'tăl) a. [L. a, away; placenta, flat cake.] Having no placenta, as monotremes.
- aplanogametangium (ăplăn'ögămētăn'jīŭm) n. [Gk. a, not; planos, wandering; gametes, spouse; anggeion, vessel.] Cell in which aplanogametes are formed.
- aplanogamete (aplan'ogamet') n.
 [Gk. a, not; planos, wandering; gametes, spouse.] A non-motile conjugating germ-cell of various plants and animals.

- aplanospore (ăplăn'öspōr) n. [Gk. a, not; planos, wandering; sporos, seed.] A non-motile resting spore of algae; an encysted spore of fungi; aplanoplastid. Opp. planospore.
- aplasia (ăplā'zĭā) n. [Gk. a, without; plassein, to mould.] Arrested development; non-development.
- aplerotic (ăplērō'tīk) a. [Gk. a, not; pleroun, to fill.] Not entirely filling a space; appl. oospore not extended to oogonial wall. Opp. plerotic.
- aploperistomatous (ăp'löpĕr'istŏm'ătŭs) a. [Gk. aploos, single; peri, around; stoma, mouth.] Having a peristome with one row of teeth, as mosses.
- aplostemonous (ăp'löstěm'önŭs) a. [Gk. aploos, single; stemon, spun thread.] With a single row of stamens.
- apneustic (ăpnū'stīk) a. [Gk. a, without; pneustos, breath.] With spiracles closed or absent; appl. aquatic larvae of certain insects.
- apobiotic (ăp'obiot'ik) a. [Gk. apo, away; bios, life.] Causing or pert. decrease in vital energy of cells or tissue; pert. apobiosis or physiological death, opp. death of entire body.
- apocarpous (ăp'ökâr'pŭs) a. [Gk. apo, away; karpos, fruit.] Having separate or partially united carpels. Opp. syncarpous.
- apocarpy,—apocarpous condition.
- apochlorosis (ăp'öklörō'sīs) n. [Gk. apo, away; chloros, grass green.]
 The absence of chlorophyll, in Flagellata.
- apocrine (ăp'ökrin) a. [Gk. apo, away; krinein, to separate.] Appl. glands secreting only part of cell contents; cf. holocrine, merocrine.
- apocyte (ăp'ösīt) n. [Gk. apo, away; kytos, hollow.] A multinucleate cell; a plurinucleate mass of protoplasm.
- apodal (ăp'ödăl), apodous (ăp'ödŭs)
 a. [Gk. a, without; pous, foot.]

- Having no feet; without ventral fin.
- apodema (ăp'ödē'mă) n. [Gk. apo, away; demas, body.] An internal skeletal projection in Arthropoda; apodeme.
- apo-enzyme (ăp'öĕn'zim) n. [Gk. apo, away; en, in; zyme, leaven.]
 Specific protein part of an enzyme, requiring co-enzyme for action.
- apogamy (ăpŏg'ămĭ) n. [Gk. apo, away; gamos, marriage.] Reproduction without intervention of sexual organs.
- apogeotropic (āp'öjē'ötrŏp'ik) a. [Gk. apo. away; gaia, earth; trope, turn.] Turning away from the earth; ageotropic.
- apogeotropism (ăp'öjēŏt'röpizm) n. [Gk. apo, away; gaia, earth; trope, turn.] Tendency to act contrarily to law of gravity; negative geotropism.
- apolegamic (ăp'ölēgăm'īk) a. [Gk. apolegein, to choose; gamos, marriage.] Appl. mating associated with sexual selection.
- apomeiosis (ap'omiō'sis) n. [Gk. apo, away; meion, smaller.] Sporogenesis without haplosis.
- apomict (ăp'ömikt) n. [Gk. apo, away; miktos, mixed.] A biotype resulting from apogamy and vegetative propagation.
- apomixis (ăp'ömik'sis) n. [Gk. apo, away; mixis, a mixing.] A reproductive anomaly in plants akin to parthenogenesis, but including development from cells other than ovules. Opp. amphimixis.
- aponeurosis (ăp'önūrō'sis) n. [Gk. apo, from; neuron, sinew.] The flattened tendon for insertion of, or membrane investing, certain muscles.
- apopetalous (ăp'öpět'álůs) a. [Gk. apo, away; petalon, leaf.] With free petals; cf. apetalous.
- apophyllous (ăp'öfîl'ŭs) a. [Gk. apo, away; phyllon, leaf.] Appl. the parts of a single perianth whorl when they are free leaves.

- apophysis (apof'isis) n. [Gk. apo. away; phyein, to grow.] Process from a bone, usually for muscle attachment; endosternite or sternal apodeme; swelling beneath reproductive structure on fungal hypha; photosynthetic region forming swelling at base of capsule in some mosses; small protuberance at apex of ovuliferous scale in pine.
- apoplasmodial (ăp'öplăsmō'dĭāl) a. [Gk. apo, away; plasma, something moulded.] Not forming a typical plasmodium.

apopyle (ăp'öpīl) n. [Gk. apo, away; pyle, gate.] Exhalent pore of sponge.

- aporogamy (ăpörŏg'ămĭ) n. [Gk. a, without; poros, channel; gamos, marriage.] Fertilisation without entry of pollen-tube through micropyle of ovule, opp. porogamy.
- aporrhysa (ăpŏr'īsă) n. plu. [Gk. aporrhein, to flow away.] Exhalent canals in sponges; opp. epirrhysa.
- aposematic (ăp'ösēmăt'ĭk) a. [Gk. apo, away; sema, signal.] Appl. warning colours which serve to frighten away enemies.
- aposporogony (ăp'öspörŏg'öni) n. [Gk. apo, away; sporos, seed; gonos, birth.] Absence of sporogony.
- apospory (ăpŏs'pöri) n. [Gk. apo, away; sporos, seed.] Production of a gametophyte from a sporophyte without intervention of spore-formation.
- apostasis (ăpŏs'tăsis) n. [Gk. apo, away; stasis, standing.] Condition of abnormal growth of axis which thereby causes separation of perianth whorls from one another.
- apostrophe (ăpòs'tröfē) n. [Gk. apo, away; strophe, turn.] Arrangement of chloroplasts along lateral walls of leaf cells.
- apothecium (ăp'öthē'siūm, -shiūm)
 n. [Gk. apo, away; theke, cup.]
 A cup-shaped ascocarp; ascocarp
 of lichens.

apotome (ăp'ötōm) n. [Gk. apo, away; tome, a cutting.] A part appearing as if cut off, as from episternum, trochanter, etc., in Arthropoda.

apotracheal (ăpötrā'kēal) a. [Gk. apo, away; L. trachea, windpipe.] With xylem parenchyma independent of vessels, or dispersed; appl. wood.

apotropous (ăpŏt'röpŭs) a. [Gk. apo, away; trope, turn.] Anatropal and with ventrally - situated raphe.

apotype,—hypotype, q.v.

apparato reticolare,—see Golgi complex.

- appendage (ăpěn'dëj) n. [L. ad, to; pendere, to hang.] An organ or part attached to a trunk, as a limb, branch, etc.; a hyphal or rigid structure for attachment or detachment of perithecium to or from mycelium, varying in structure and function in different Ascomycetes.
- appendices colli (ăpěn'disez kŏl'ī)

 n. plu. [L. ad, to; pendere, to hang; collum, neck.] Exterior throat appendages or tassels, of goat, sheep, pig, etc.
- appendicular (ăp'ĕndĭk'ūlăr) a. [L. ad, to; pendere, to hang.] Pert. appendages; appl. skeleton of limbs, opp. axial skeleton; pert. vermiform appendix; appl. artery.

appendiculate (ăp'ĕndik'ūlāt) a. [L. ad, to; pendere, to hang.] Having a small appendage, as a stamen or filament.

- appendix (ăpěn'diks) n., appendices (ăpěn'disēz) plu. [L. ad, to; pendere, to hang.] An outgrowth, especially the vermiform appendix.
- applanate (ăpl'ānāt) a. [L. ad, to; planatus, flattened.] Flattened.
- apposition (ap'özish'un) n. [L. ad, to; ponere, to place.] The formation of successive layers in growth of a cell wall; cf. intussusception.
- appressorium (ăp'rēsō'rīum) n. [L. ad, to; pressare, to press.] Adhesive disc, as of haustorium or sucker.

- aproterodont (åprot'ërödont) a. [Gk. a, without; proteros, first; odous, tooth.] Having no premaxillary teeth.
- apteria (ăptē'riă) n. plu. [Gk. a, without; pteron, wing.] Naked or down-covered surfaces between pterylae or feather-tracts.
- apterous (ăp'tërŭs) a. [Gk. a, without; pteron, wing.] Wingless; having no wing-like expansions on stems or petioles; exalate.
- apterygial (ăp'tërij'iăl) a. [Gk. a, without; pterygion, dim. of pteron, wing.] Wingless; without fins.
- apterygotous (ăptërigō'tŭs) a. [Gk. a, without; pterygotos, winged.] Resembling or pert. primitive wingless insects.
- aptychus (apt'ikus) n. [Gk. a, without; ptyche, fold.] A horny or calcareous structure, possibly an operculum, of ammonites.
- apyrene (ăpi'rēn) a. [Gk. a, not; pyren, fruit-stone.] Appl. spermatozoa lacking nucleus; cf. eupyrene, oligopyrene; seedless, appl. certain cultivated fruits.
- aquatic (ăkwăt'îk) a. [L. aqua, water.] Pert. water; living in or frequenting water. n. An aquatic plant.
- aqueduct (ăk'wēdŭkt) n. [L. aqua, water; ducere, to lead.] A channel or passage, as that of cochlea, and of vestibule of ear; aquaeductus.
- aqueduct of Sylvius [F. de Boë or Sylvius, Flemish anatomist],—cerebral aqueduct, aqueduct of the midbrain, or iter, connecting third and fourth ventricle; mesocoele.
- aqueous (ā'kweus) a. [L. aqua, water.] Watery, appl. humour, fluid occupying space between lens and cornea; appl. tissue consisting of thin-walled watery parenchymatous cells.
- arachnactis (ărăknăk'tis) n. [Gk. arachne, spider; aktis, ray.]

- Larval stage of cerianthid Zoan-tharia.
- arachnid (ārāk'nĭd) a. [Gk. arachne, spider.] Spider-like; pert. spiders. arachnidium (ārāknīd'īum) n. [Gk. arachne, spider; idion, dim.] The spinning apparatus of a spider, including spinning-glands and

spinnerets.

brane.

- arachnoid (ărăk'noid) a. [Gk. arachne, spider, cobweb; eidos, form.] Pert. or resembling a spider; like a cobweb; consisting of fine entangled hairs; appl. the thin membrane between dura and pia mater. n. The arachnoid mem-
- arachnoideal (ărăknoid'eāl) a. [Gk. arachne, cobweb; eidos, form.] Pert. the arachnoid; appl. granulations: Pacchionian bodies, q.v.
- arbacioid,—see diadematoid.
- arborescent (âr'börĕs'ënt) a. [L. arborescens, growing like a tree.] Branched like a tree.
- arborisation (âr'börīzā'shūn) n. [L. arbor, tree.] Tree-like branching, as of nerve cell processes; arborescence.
- arboroid (âr'böroid) a. [L. arbor, tree; Gk. eidos, like.] Tree-like, designating general structure of a protozoan colony; dendritic.
- arbor vitae (ârbor vi'tē) n. [L. arbor, tree; vita, life.] The tree of life, appl. arborescent appearance of cerebellum in section.
- arbuscle (âr'būsl) n. [L. arbuscula, shrub.] A tree-like small shrub, or a dwarf tree; a branched haustorium, as in certain fungi; arbuscula.
- arbuscular (ârbŭs'kūlăr) a. [L. arbuscula, shrub.] Resembling a tree-like small shrub.
- arcade (ârkād') n. [L. arcus, arch.]
 An arched channel or passage; a
 bony arch, as supra- and infratemporal arches in skull; transverse canal connecting lateral
 canals, in Ascaris.
- Archaean (ârkē'ăn) a. [Gk. archaios, ancient.] Appl. geological era before Palaeozoic; Pre-Cambrian.

- archaeocytes (âr'kēösīts) n. plu.
 [Gk. archaios, primitive; kytos, hollow.] Cells arising from undifferentiated blastomeres and ultimately giving rise to germ-cells and gametes.
- archaeostomatous (âr'kēöstŏm'ătŭs)
 a. [Gk. archaios, primitive; stoma, mouth.] Having the blastopore persistent and forming mouth.

Archaeozoic (âr'kēözō'ik) a. [Gk. archaios, ancient; zoe, life.] Pert. earliest geological era, age of unicellular life.

arch-centra (arch'sĕn'tră) n. plu. [L. arcus, bow; centrum, centre.] Centra formed by fusion of basal growths of primary arcualia external to chordal sheath; cf. chordacentra.

archebiosis (âr'këbiō'sis) n. [Gk. arche, beginning; biosis, living.]
The origin of life; archegenesis.

- archedictyon (âr'këdik'tiŏn) n. [Gk. arche, beginning; diktyon, net.]
 An intervein network in wings of some primitive insects.
- archegoniophore (âr'kegō'nĭöfōr) n. [Gk. arche, beginning; gonos, off-spring; pherein, to bear.] Branches of bryophytes, or parts of fern prothalli, bearing archegonia.
- archegonium (âr'këgō'niŭm) n. [Gk. arche, beginning; gonos, offspring.]
 A female gametangium in which oospheres are formed, and in which the young plant begins development.
- archencephalon (ârk'čnkěf'ălŏn, -sĕf-) n. [Gk. arche, beginning; engkephalos, brain.] The primitive forebrain or cerebrum.
- archenteron (ârkěn'tërŏn) n. [Gk. arche, beginning; enteron, gut.]
 The cavity of gastrula which forms primitive gut of embryo.

archeo-,—archaeo-.

- archespore (âr'këspör) n. [Gk. arche, beginning; sporos, seed.]
 The tetrahedral or meristematic cell of a sporangium; cell of an archesporium.
- archesporium (âr'këspō'riŭm) n. [Gk. arche, beginning; sporos,

seed.] A cell or mass of cells, dividing to form spore mother-cells, or, in liverworts, spore mother-cells and elater-forming cells.

archetype, -architype, q.v.

- archiamphiaster (år'kiam'fias'tër) n.
 [Gk. archi, first; amphi, on both sides; aster, star.] The amphiaster forming first or second polar body in maturation of ovum.
- archibenthic (âr'kibĕn'thik) a. [Gk. archi, first; benthos, depths of sea.] Pert. bottom of sea from edge of continental shelf to upper limit of abyssobenthic zone, at depths of ca. 200 to 1000 metres.
- archiblast (âr'kiblăst) n. [Gk. archi, first; blastos, bud.] Egg protoplasm. archiblastic (âr'kiblăs'tik) a. [Gk. archi, first; blastos, bud.] Having total and equal segmentation.
- archiblastula (âr'kiblăs'tūlă) n. [Gk. archi, first; blastos, bud.] Typical hollow ball of cells derived from an egg with total and equal segmentation.
- archicarp (âr'kĭkârp) n. [Gk. archi, first; karpos, fruit.] Spirally coiled region of thallus, or stalk bearing oogonium, of certain fungi.
- archicerebrum (âr'kisĕr'ébrŭm) n. [Gk. archi, first; L. cerebrum, brain.] The primitive brain, as the supra-oesophageal ganglia of higher invertebrates; primary brain of arthropods.
- archichlamydeous (âr'kiklămĭd'eŭs)
 a. [Gk. archi, first; chlamys, cloak.] Having no petals, or having petals entirely separate from one another.
- archicoel (âr'kĭsēl) n. [Gk. archi, first; koilos, hollow.] The primary body-cavity or space between alimentary canal and ectoderm in development of various animals.

archidictyon,—see archedictyon. archigenesis (âr'kĭjēn'ēsĭs) n. [Gk. archi, first; genesis, descent.] Abiogenesis, q.v.

archigony (arkig'öni) n. [Gk. archi, first; gonos, begetting.] The first origin of life.

archinephric (ar'kiněf'rik) a. [Gk. archi, first; nephros, kidney.] Appl. duct into which pronephric tubules open; pert. archinephros.

archinephridium (ar'kinĕfrid'iŭm) n. [Gk. archi, first; nephros, kidney; idion, dim.] Excretory organ of certain larval invertebrates; solenocyte, q.v.

archinephros ((âr'kĭnĕf'rŏs) n. [Gk. archi, first; nephros, kidney.] The primitive kidney; Wolffian body.

- archipallium (âr'kipăl'iŭm) n. [Gk. archi, first; L. pallium, mantle.] The olfactory region of cerebral hemispheres, comprising olfactory bulbs and tubercles, pyriform lobes, hippocampus, and fornix. neopallium.
- archiplasm (âr'kiplăzm) n. [Gk. archi, first; plasma, mould.] The substance of attraction-sphere, astral rays, and spindle-fibres; also archoplasm; kinoplasm; idiosome, q.v.
- archipterygium (âr'kĭtërĭj'ĭŭm) n. [Gk. archi, first; pterygion, little wing.] Type of fin in which skeleton consists of elongated segmented central axis and two rows of jointed
- architomy (arkit'omi) n. [Gk. archi, first; tome, cutting. | Reproduction by fission with subsequent regeneration, in certain annelids; paratomy.
- architype (âr'kĭtīp) n. [Gk. archi, first; typos, type.] An original type from which others may be derived.

archoplasm,—archiplasm, q.v.

- arcicentrous (âr'sĭsĕn'trŭs) a. [L. arcus, bow; centrum, centre.] Appl. vertebral column in which centra are mainly derived from arch tissue: arcocentrous.
- arciferous (ârsĭf'ërŭs) a. [L. arcus. bow; ferre, to carry.] Appl. pectoral arch of toads, etc., where precoracoid and coracoid are separated and connected by arched epicoracoid.

arciform (âr'sĭfôrm) a. [L. arcus, bow; forma, shape.] Shaped like an arch or bow; arcuate.

- arcocentrous (ar'kösen'trus) a. [L. bow; arcus, centrum, centre.] Appl. vertebral column with inconspicuous chordal sheath and centra derived from arch tissue.
- arcocentrum (âr'kösĕn'trŭm) n. [L. arcus, bow; centrum, centre.] A centrum formed from parts of neural and haemal arches.
- Arctogaea (ârk'töjē'ă, -gâ'yă) n. Gk. Arktos, Great Bear; gaia, earth.] Zoogeographical area comprising Holarctic, Ethiopian, and Oriental regions.
- arcualia (âr'kūā'lĭă) n. plu. arcus, bow.] Small cartilaginous pieces, dorsal and ventral, fused or free, on vertebral column of fishes.
- arcuate (âr'kūāt) a. [L. arcus, bow.] Curved or shaped like a bow.
- arculus (âr'kūlūs) n. [Dim. of L. arcus, bow.] Arc formed by two wing veins of certain insects.
- ardellae (ârděl'ē) n. plu. [Gk. ardein, to sprinkle.] Small apothecia of certain lichens, having appearance of dust.
- [L. area, groundarea (ā'rēă) n. space.] A surface, as area opaca, area pellucida, area vasculosa, etc.; part enclosed by a raised ridge, as in Polyzoa; a region.
- arenaceous (ărenā'shus) a. arena, sand.] Having properties or appearance of sand; sandy; growing in sand.
- arenicolous (ărenik'olus) a. arena, sand; colere, to inhabit.] Living in sand; psammophilous.
- areola (ărē'ölă) n. [L. areola, dim. of arca, space.] A small coloured circle round a nipple; part of iris bordering pupil of eye; one of small spaces or interstices of a special kind of tissue; area defined by cracks on surface of lichens; poroids when surrounded by thickened margins; scrobicula, q.v.
- areolar (ărē'ölăr) a. L. areola, small space.] Of or like an areola; pert. an areola.
- areolate (ărē'ölāt) a. [L. areola, small space.] Divided into small areas defined by cracks or other margins.

areolation (ărēölā'shǔn) n. [L. areola, small space.] Areolar pattern or network appearance, as of cell margins in tissue.

areole (ăr'ēōl) n. [L. areola, small space.] Areola, q.v.; space occupied by a group of hairs or

spines, as in Cactus.

argentaffin (ârjěn'tăfin) a. [L. argentum, silver; affinis, related.] Staining with silver salts, appl. cells; argyrophil.

argenteal (ârjen'teăl) a. [L. argenteus, silvern.] Appl. layer of eye

containing calcic crystals.

argenteous (ârjen'teus) a. [L. argenteus, silvern.] Like silver.

argenteum (ârjĕn'tĕŭm) n. [L. argenteus, silvern.] A dermal reflecting tissue layer of iridocytes, without chromatophores, in fishes.

arginase (âr'jīnās) n.—a liver enzyme acting on the amino-acid arginine $(C_0H_{14}O_3N_4)$, urea and ornithine being separated by hydrolysis.

argyrophil (âr'jiröfil) a. [Gk. argyros, silver; philos, loving.] Staining with silver salts, appl. fibres of reticular tissue; argentaffin; argentophil, appl. basal bodies or blepharoplasts.

aril (ăr'il) n. [F. arille, Sp. arillo, a small hoop.] An additional integument formed on some seeds after

fertilisation.

arillode (ar'ilōd) n. [F. arille, hoop; Gk. eidos, like.] A false arillus arising from region of micropyle as an expansion of exostome.

arillus (ăril'ŭs) n. [L.L. arillus, aril.]

An aril, q.v.

arista (ărīs'tă) n. [L. arista, awn.]
Awn; long-pointed process as in
many grasses; a bristle borne by
antenna of many brachycerous
Diptera.

aristate (ărīs'tāt) a. [L. arista, awn.]
Provided with awns, or with a welldeveloped bristle; appl. insect

antenna.

aristogenesis (ăr'îstöjěn'ësīs) n. [Gk. aristos, best; genesis, descent.]
Process of evolving new biomechanism from the germ plasm; creative

principle or potentiality in origin of species.

aristogenic,—eugenic, q.v.

Aristotle's lantern,—masticating apparatus of sea-urchin.

arkyochrome (âr'kiökrōm) a. [Gk. arkys, net; chroma, colour.] With Nissl granules arranged like network; appl. certain neurones.

armature (âr'mătūr) n. [L. armatura, armour.] Anything which serves to defend, as hairs, prickles, thorns, spines, stings, etc.

arolium (ărō'lĭum) n. [Gk. arole, protection.] Median lobe or pad on praetarsus of many insects.

aromorph (ā'römôrf) n. [Gk. airein, to raise; morphe, form.] A character or structure resulting from aromorphosis.

aromorphosis (ā'römôr'fōsis) n. [Gk. airein, to raise; morphosis, shaping.] Evolutionary change towards an increase in life energy, e.g. evolution of a biting mouth skeleton from gill arches; opp. evolution of a merely adaptational character.

array (ărā') n. [F. arroi, retinue.] Arrangement in order of magnitude.

arrect (ărĕkt') a. [L. arrectus, set upright.] Upright; erect.

arrectores pilorum,—bundles of non-striped muscular fibres associated with hair follicles,—contraction causing hair to stand on end. Sing. arrector pili.

arrhenogenic (ărĕn'öjĕn'ik) a. [Gk. arrhen, male; genos, offspring.]
Producing offspring preponderantly

or entirely male.

arrhenoid (ăren'oid) a. [Gk. arrhen, male; eidos, form.] Exhibiting male characteristics, as genetically female animals undergoing sexreversal.

arrhenoplasm (ărĕn'öpläzm) n. [Gk. arrhen, male; plasma, mould.]
Male plasm, in reference to theory that all protoplasm consists of arrhenoplasm and thelyplasm.

arrhenotoky (ărenot'oki) n. [Gk. arrhen, male; tokos, birth.] Parthenogenetic production of males.

arrhostia (ărōstī'ā) n. [Gk. arrhostia, 1] health.] A normal condition or trend in development or evolution. which resembles a diseased condition, e.g. extreme size in certain extinct vertebrates resembling overaction of pituitary gland.

artefact (âr'tëfăkt) n. [L. ars, art; factus, made.] An appearance, or apparent structure, due to prepara-

tion and not natural.

artenkreis (âr'tënkris) n. [Ger. Art, species; Kreis, circle.] Complex of species which replace one another geographically; superspecies.

arterial (ârtē'riăl) a. [L. arteria, artery.] Pert. an artery, or system of channels by which blood issues to body from heart.

arteriole (ârtē'rĭōl) n. [L. arteriola, small artery.] A small artery.

artery (âr'tëri) n. [L. arteria, artery.]
A vessel which conveys blood from heart to body.

arthritic (ârthrit'ik) a. [Gk. arthron,

joint.] *Pert*. or at joints.

arthrobranchiae (âr'thröbrăng'kiē)
n. plu. [Gk. arthron, joint;
brangchia, gills.] Joint-gills, arising
at junction of thoracic appendage
with trunk, of Arthropoda.

arthrodia (ârthrō'diă) n. [Gk. arthron, joint.] A joint admitting of

only gliding movements.

arthrodial (ârthrō'diăl) a. [Gk. arthron, joint.] Appl. articular membranes connecting thoracic appendages with trunk, as in arthropods.

arthrogenous (ârthrŏj'ënŭs) a. [Gk. arthron, joint; genos, descent.] Formed as a separate joint, as spores; developed from separated

portions of a plant.

arthromere (âr'thrömēr) n. [Gk. arthron, joint; meros, part.] An arthropod body-segment or somite.

arthropod (âr'thrôpôd) a. [Gk. arthron, joint; pous, foot.] With jointed legs; pert. phylum including Crustacea, Myriopoda, Insecta, Arachnoidea.

arthropterous (ârthrop'tërus) a. [Gk.

arthron, joint; pteron, wing.] Having jointed fin-rays, as fishes.

arthrospore (arth'röspōr) n. [Gk. arthron, joint; sporos, seed.] A resting moniliform bacterial cell; a cell formed by segmentation of a hypha.

arthrosterigmata (âr'thröstērīg'mătă) n. plu. [Gk. arthron, joint; sterigma, support.] Jointed sterig-

mata.

arthrostracous (ârthrŏs'trākūs) a. [Gk. arthron, joint; ostrakon, shell.] Having a segmented shell.

arthrotergal (ar'thröter'gal) a. [Gk. arthron, joint; L. tergum, back.] Appl. median dorsal flexor of opisthosoma in Limulus.

articular (ârtik'ūlăr) a. [L. articulus, joint.] Pert. or situated at a joint; appl. cartilage, lamellae, surface, capsule, etc.

articularis genus, — subcrureal muscle.

articulated (ârtĭk'ūlātēd) a. [L. articulus, joint.] Jointed; articulate; separating easily at certain points.

articulation (âr'tikūlā'shūn) n. [L. articulus, joint.] A joint between bones or segments, or between segments of a stem.

artifact,-artefact, q.v.

artiodactyl (âr'tiödāk'til) a. [Gk. artios, even; daktylos, finger.] Having an even number of digits.

arytaenoid (ăr'îtē'noid) a. [Gk. arytaina, ladle; eidos, form.]
Pitcher-like; appl. two cartilages at back of larynx, also glands, muscles,

asci,—plu. of ascus.

ascidial (ăsīd'īāl) a. [Gk. askidion, dim. of askos, bag.] Sac-like; appl. certain specialised, or abnormal, floral and foliage leaves; pert. ascidium.

ascidian (ăsid'iăn) a. [Gk. askidion, little bag.] Like an ascidian or sea-

squirt.

ascidium (ăsīd'iŭm) n. [Gk. askidion, little bag.] A pitcher-leaf, as in Nepenthes. ascigerous (ăsij'ërüs) a. [Gk. askos, bag; L. gerere, to bear.] Bearing asci, as certain hyphae in fungi; asciferous.

ascocarp (ăs'kökârp) n. [Gk. askos, bag; karpos, fruit.] Asci with their protective covering; sporo-

carp of Ascomycetes.

ascogenous (ǎskŏj'ėnūs) n. [Gk. askos, bag; -genes, producing.]
Producing asci; appl. hyphae, cells.

- ascogonium (ăs'kögō'niŭm) n. [Gk. askos, bag; gonos, offspring.] A specialised hyphal branch which gives rise to ascogenous hyphae or an ascus; oogonium of Ascomycetes.
- ascoma (ăs'kōmă) n. [Gk. askoma, leather padding.] Disc-shaped ascocarp in certain fungi.
- ascophore (ăs'kofōr) n. [Gk. askos, bag; pherein, to bear.] Ascocarp. ascorbic acid,—pure vitamin C, deficiency of which in diet causes dental disorders and scurvy; hexuronic acid, C₆H₈O₆.

ascospore (ăs'köspōr) n. [Gk. askos, bag; sporos, seed.] One of the spores produced in an ascus.

ascus (as'kus) n. [Gk. askos, bag.] A membranous spore-sac, as of Ascomycetes.

asemic (åsē'mik) a. [Gk. asemos, without sign.] Without markings. aseptate (åsĕp'tāt) a. [L. a, not;

septum, partition.] Without any

septum.

asexual (åsěk'sūăl) a. [Gk. a, without; L. sexus, sex.] Having no apparent sexual organs; parthenogenetic or vegetative, as appl. reproduction.

asiphonate (ăsī'fönāt) a. [L. a, not; sipho, tube.] Appl. larvae whose respiratory tubes open directly to

exterior.

asparagine (ăspăr'ăjēn, -gin) n. [Gk. asparagos, asparagus.] A compound, first detected in asparagus, formed from amino-acids in leguminous and other seeds, of importance in nitrogen metabolism of plants; C₄H₈O₈N₂.

aspect (ăs'pěkt) n. [L. aspicere, to

look toward.] Direction facing part of a surface; appearance or look; seasonal appearance.

aspection (ăspěk shun) n. [L. aspicere, to look toward.] Seasonal succession of phytological and zoological phenomena.

asperity (ăspěr'iti) n. [L. asper, rough.] Roughness, as on a leaf.

asplanchnic (äsplängk'nik) a. [Gk. a, without; splangchna, viscera.]
Without alimentary canal.

- asporocystid (ăspŏ'rösis'tid) a. [Gk. a, not; sporos, seed; kystis, bladder; idion, dim.] Appl. oocyst of Sporozoa when zygote divides into sporozoites without sporocyst formation.
- asporous (*aspō'rus) a. [Gk. a, without; sporos, seed.] Having no spores.
- assimilation (āsīm'īlā'shūn) n. [L. ad, to; similis, like.] Conversion into protoplasm of ingested and digested nutrient material; anabolism.
- association (āsō'siā'shūn) n. [L. ad, to; socius, fellow.] A plant community forming a division of a formation or larger unit of vegetation, as of tundra, grassland, forest, and characterised by dominant species; adherence of gregarines without fusion of nuclei; appl. fibres connecting white matter of interior of brain with cortex; appl. neurons with intersegmental axons.

associes (ăsō'siēz) n. [L. ad, to; socius, fellow.] An association representing a stage in the process of succession.

astacene (ăs'tăsēn) n. [L. astacus, crayfish.] Carotenoid pigment of certain crustaceans, echinoderms, and fishes; astacin; C₄₀H₄₈O₄.

astaxanthin (ās'tāzān'thǐn) n. [L. astacus, crayfish; Gk. xanthos, yellow.] An animal carotenoid derived from ingested plant carotenoids, in chromoplasts of certain flagellates, also combining with proteins to form pigments, as of crustaceans; C₄₀H₈₂O₄.

astelic (aste'lik) a. [Gk. a, without; stele, pillar.] Not possessing a

stele.

- astely (ăstē'li) n. [Gk. a, without; stele, pillar.] Absence of a central cylinder, axis, or stele.
- aster (ăs'tër) n. [Gk. aster, star.]
 The star-shaped achromatinic structure surrounding centrosome during mitosis; star-shaped arrangement of chromosomes during mitosis.
- asterion (ăstē'riŏn) n. [Gk. aster, star,] The region of posterolateral fontanelle where lambdoid, parieto - mastoid, and occipitomastoid sutures meet.
- asteriscus (ăs'tëris'kŭs) n. [Gk. asteriskos, dim. of aster, star.] A small otolith in rudimentary cochlea of teleosts.
- asternal (aster'nal) a. [L. a, from; sternum, breastplate.] Appl. ribs whose ventral ends do not join the sternum directly.
- asteroid (ăs'tëroid) a. [Gk. aster, star; eidos, form.] Star-shaped; pert. star-fish.
- asterospondylous (ăs'tëröspŏn'dilŭs)
 a. [Gk. aster, star; sphondylos,
 vertebra.] Having centrum with
 radiating calcified cartilage; also
 asterospondylic.
- asthenic (ăsthěn'îk) a. [Gk. asthenos, feeble.] Weak; tall and slender; leptosome.
- astogeny (ăstŏj'eni) n. [Gk. astos, citizen; genos, descent.] The development of a colony by budding.
- astomatous (ăstőm'ātŭs) a. [Gk. a, without; stoma, mouth.] Not having a mouth; without epidermic pores or stomata.
- astomous (ăs'tömŭs) a. [Gk. a, without; stoma, mouth.] Without a stomium or line of dehiscence; bursting irregularly.
- astragalus (ăstrăg'ālūs) n. [Gk. astragalos, ankle-bone.] The talus, second largest tarsal bone in man; a tarsal bone in verte-brates.
- astroblast (ăs'tröblăst) n. [Gk. aster, star; blastos, bud.] A cell giving rise to protoplasmic or to fibrillar astrocytes.

- astrocentre (ăs'trösĕn'tër) n. [L. aster, star; centrum, centre.]
 Centrosome.
- astrocyte (ăs'trösīt) n. [Gk. aster, star; kytos, hollow.] A common neuroglia cell; astroglia; macroglia; Deiters' cell; a neuroglial cell with branching protoplasmic processes in grey matter; a fibrillar or spider cell in white matter.
- astropodia (ăs'tröpō'dĭā) n. plu.
 [Gk. aster, star; pous, foot.]
 Fine unbranched radiating pseudopodia, as in Heliozoa and some Radiolaria.
- astropyle (ăs'tröpīl, -pī'lē) n. [Gk. aster, star; pyle, gate.] Chief aperture of central capsule, in certain Radiolaria.
- astrosclereid (äs'trösklë'rëid) n. [Gk. aster, star; skleros, hard; eidos, form.] A multiradiate sclereid or stone cell; a spiculate or ophiuroid cell.
- astrosphere (ăs'trösfēr) n. [Gk. aster, star; sphaira, ball.] Central mass of aster without rays; aster exclusive of centrosome; astral sphere.
- asymmetrical (åsimět'rikăl) a. [Gk. asymmetros, disproportionate.] Pert. want of symmetry; having two sides unlike or disproportionate; appl. structures or organs which cannot be divided into similar halves by any plane; asymmetric.
- asynapsis (ăsinăp'sis) n. [Gk. a, not; synapsis, union.] Absence of pairing of chromosomes in meiosis; asyndesis.
- atactostele (ătăk'töstē'lē) n. [Gk. ataktos, irregular; stele, post.] A complex stele having bundles scattered in the ground tissue, as in monocotyledons.
- atavism (ăt'ăvizm) n. [L. atavus, ancestor.] Reversion, occurrence of an ancestral characteristic not observed in more immediate progenitors.
- atavistic (ăt'ăvis'tik) a. [L. atavus, ancestor.] Pert., marked by, or tending to atavism.

- ateleosis (ătěleō'sis) n. [Gk. ateles, imperfect.] Dwarfism where individual is a miniature adult; cf. achondroplasia.
- atelia (ătěl'ĭă) n. [Gk. ateles, ineffectual.] The apparent uselessness of a character of unknown biological significance; incomplete development.
- atelomitic (ătělömīt'īk) a. [Gk. a, not; telos, end; mitos, thread.]

 Appl. other than terminal attachment of chromosome to spindle.

athalamous (åthäl'ämŭs) a. [Gk. a, without; thalamos, inner room.]
Lacking a thalamus.

- athrocytosis (ăth'rösītō'sīs) n. [Gk. athroos, collected; kytos, hollow.]
 The capacity of cells to selectively absorb and retain solid particles in suspension, as dyes.
- atlanto-occipital,—occipito-atlantal, q.v.
- atlas (ăt'lăs) n. [Gk. Atlas, a Titan.] The first cervical vertebra.
- atokous (ăt'ökŭs) a. [Gk. atokos, childless.] Without offspring.
- atoll (ăt'ôl, ătôl') n. [Mal. atoll.] A coral reef surrounding a central lagoon.
- atopy (ăt'öpĭ) n. [Gk. atopia, unusual nature.] Idiosyncrasy, genetic sensitivity to poisonous effects of particular antigens or atopens, as of certain proteins, pollen, etc.
- atretic (atre'tik) a. [Gk. a, not; tretos, perforated.] Appl. vesicles resulting from degeneration of Graafian follicles, spurious corpora lutes
- atrial (ā'triăl) a. [L. atrium, central room.] *Pert*. atrium; appl. cavity, pore, canal, siphon, lobes.
- atrichic (ătrikik) a. [Gk. a, not, thrix, hair.] Having no flagella; atrichous.
- atriocoelomic (ā'trīösēlŏm'īk) a. [L. atrium, central room; Gk. koiloma, a hollow.] Connecting atrium and coelom; appl. funnels, of uncertain function, in Cephalochorda.
- atriopore (ā'trīöpōr) n. [L. atrium, central room; porus, channel.]
 The opening from atrial cavity to

exterior in Cephalochorda; spiracle in tadpole.

atrioventricular (ā'trīovēntrīk'ūlār)
a. [L. atrium, chamber; ventriculus,
small cavity.] Pert. atrium and
ventricle of heart; appl. bundle,
groove, node, openings.

atrium (ā'trium) n. [L. atrium, chamber.] Anterior cavity of heart; tympanic cavity; a division of the vestibule at end of bronchiole; chamber surrounding pharynx in Tunicata and Cephalochorda.

atrochal (ăt'rökăl) a. [Gk. a, without; trochos, wheel.] Without preoral circlet of cilia; appl. trochophore when preoral circlet is absent and surface is uniformly ciliated.

atropal,-atropous.

atrophy (ăt'röfi) n. [Gk. a, without; trophe, nourishment.] Emaciation; diminution in size and function.

- atropous (ât'röpŭs) a. [Gk. a, without; trope, turn.] Appl. ovule in proper position, i.e. not inverted.
- attachment, the spindle attachment; a lasting fusion of two chromosomes.
- attenuated (ătěn'ūātėd) a. [L. attenuare, to thin.] Thinned; reduced in density, strength, or pathogenic activity.
- atterminal (ătter'minăl) a. [L. ad, to; terminus, end.] Towards a terminal; appl. current directed toward thermal cross-section.
- attic (ăt'îk) n. [Gk. attikos, Athenian.] The epitympanic recess. attraction-particle,—centriole, q.v. attraction sphere, centrosphere, q.v.
- auditory (ôd'itôri) a. [L. audire, to hear.] Pert. hearing apparatus, appl. organ, nucleus, ossicle, capsule, canal, meatus, nerve, vesicle, etc.; pert. sense of hearing.
- Auerbach's plexus [L. Auerbach, German anatomist]. A gangliated plexus of non-medullated nervefibres, found between the circular and longitudinal layers of muscular coat of small intestine; plexus myentericus.
- augmentation (ôgměntā'shun) n. [L. augere, to increase.] Increase in number of whorls; cf. chorisis.

- augmentor (ôgmĕn'tŏr) a. [I.. augere, to increase.] Appl. nerves rising from sympathetic system and acting on heart, with antagonistic relation to vagi; accelerator.
- aulophyte (ôl'öfit) n. [Gk. aulon, hollow way; phyton, plant.] A non-parasitic plant growing in hollow of another.
- aulostomatous (ôl'östőm'átús) a.
 [Gk. aulos, tube; stoma, mouth.]
 Having a tubular mouth or snout.
- aural (ôr'ăl) a. [L. auris, ear.] Pert. ear or hearing.
- auricle (ôr'ikl) n. [L. auricula, small ear.] Any ear-like lobed appendage; the external ear; atrium or anterior chamber of heart.
- auricula (ôrīk'ūlă) n. [L. auricula, small ear.] An auricle.
- auricular (ôrìk'ūlăr) n. [L. auricula, small ear.] Ear covert of birds. a. Pert. an auricle; appl. artery, nerve, tubercle, vein.
- auricularia (ôrik'ūlā'rīā) n. [I.. auricula, small ear.] A type of larva found among Holothuria.
- auricularis (ôrikūlā'rīs) n. [L. auricula, earlap.] Superior, anterior, posterior, extrinsic muscles of the external ear.
- auriculate (ôrlk'ūlāt) a. [L. auricula, small ear.] Eared; appl. leaf with expanded bases surrounding stem; appl. leaf with lobes separate from rest of blade; hastate-auricled.
- auriculo-ventricular,—pert. or connecting auricle and ventricle of heart; appl. bundle, valve.
- auriform (ô'rifôrm) a. [L. auris, ear; forma, shape.] Resembling the external ear in shape, as shell of Haliotis.
- aurophore (ôr'öfōr) n. [L. auris, ear; Gk. pherein, to bear.] An organ projecting from base of pneumatophore of certain Siphonophora.
- austral (ôs'trăl) a. [L. australis, southern.] Appl. or pert. southern biogeographical region, or restricted to North America between transitional and tropical zones.

- Australian (ôstrā'liān) a. [L. australis, southern.] Appl. or pert. a zoogeographical region including Papua, Australia, New Zealand, and Pacific islands.
- autacoid (ôt'ākoid) n. [Gk.autos, self; akos, remedy; eidos, form.] Internal secretion, a hormone or a chalone.
- autarticular (ôt'ârtĭk'ūlăr) n. [Gk. autos, self; L. articulus, joint.] Gonial, q.v.
- autecology (ôt'ēkŏl'öjī) n. [Gk. autos, self; oikos, household; logos, discourse.] The biological relations between a single species and its environment; ecology of an individual organism; auto-ecology.
- autobiology,—idiobiology, q.v.
 autoblast (ôt'oblăst) n. [Gk. autos, self; blastos, bud.] An independent micro-organism or cell.
- autocarp (ôt'ökârp) n. [Gk. autos, self; karpos, fruit.] Fruit resulting from self-fertilisation.
- autocatalysis (ôt'ökätäl'isis) n. [Gk. autos, self; kata, down; lysis, loosing.] Dissolution or reaction of a cell or substance due to influence of a product or secretion of its own. autochthon (ôtök'thön) n. [Gk.
- autochthon (ötök'thön) n. [Gk. autos, self; chthon, the ground.]

 An indigenous species.
- autochthonous (ôtôk'thônūs) a. [Gk. autos, self; chthon, ground.] Aboriginal; indigenous; inherited or hereditary, native, appl. characteristics; originating within an organ, as pulsation of excised heart; formed where found. Opp. allochthonous.
- autocoid,—autacoid, q.v.
- autocyst (ôt'ösist) n. [Gk. autos, self; kystis, bladder.] A thick membrane formed by Neosporidia separating them from host tissues.
- autodermalia (ôt'ödermā'līā) n. plu.
 [Gk. autos, self; derma, skin.]
 Dermal spicules with axial cross,
 within dermal membrane.
- autodont (ôt'ödönt) a. [Gk. autos, self; odous, tooth.] Designating or pert. teeth not directly attached to jaws, as in cartilaginous fishes.

autoecious (ôtë'sĭŭs) a. [Gk. autos, self; oikos, house.] Passing different stages of life history in the same host; appl. parasitic fungi; autoxenous.

autogamous (ôtŏg'ămŭs) a. [Gk. autos, self; gamos, marriage.] Self-

fertilising, opp. allogamous.

autogamy (ôtŏg'āmī) n. [Gk. autos, self; gamos, marriage.] Self-fertilisation, opp. allogamy; conjugation of nuclei within a single cell; conjugation of two protozoa originating from division of the same individual.

autogenesis (ôt'öjĕn'ĕsĭs) n. [Gk. autos, self; genesis, birth.] Spontaneous generation; autogeny, autogony.

autogenetic (ôt'öjēnět'ik) a. [Gk. autos, self; genesis, birth.] Reproducing spontaneously, as body-cells.

- autogenic (ôtöjěn'ík) a. [Gk. autos, self; gennaein, to produce.] Caused by reactions of organisms themselves; appl. plant successions, opp. allogenic; autonomic or spontaneous, appl. movements.
- autogenous (ôtôj'enŭs) a. [Gk. autos, self;-genes, producing.] Produced in the same organism; appl. enzymes; appl. graft reimplanted in same animal; appl. vaccine injected into same animal; appl. variations due to changes within chromosomes.

autogony (ôtŏg'önĭ) n. [Gk. autos, self; gonos, offspring.] Autogenesis, q.v.

- autoheteroploid (ô'töhĕt'éröploid) n.
 [Gk. autos, self; heteros, other; aploos, onefold; eidos, form.]
 Heteroploid derived from a single genome or multiplication of some of its chromosomes.
- autoinfection (ôt'öĭnfĕk'shun) n. [Gk. autos, self; L. inficere, to taint.] Reinfection from host's own parasites.
- autointoxication (ôt'öintök'sikā'-shūn) n. [Gk. autos, self; L. in, in; Gk, toxikon, poison.] Reabsorption of toxic substances produced by the body.
- autolysis (ôtŏl'īsīs) n. [Gk. autos,

self; *lysis*, loosing.] Self-digestion; cell or tissue disintegration by action of autogenous enzymes.

autolytic (ôt'ölīt'īk) a. [Gk. autos, self; lysis, loosing.] Causing or pert. autolysis; appl. enzymes.

- automixis (ôt'ömik'sis) n. [Gk. autos, self; mixis, mingling.] The union, in a cell, of chromatin derived from common parentage; self-fertilisation.
- autonarcosis (ôt'önârkō'sīs) n. [Gk. autos, self; narke, numbness.] State of being poisoned, rendered dormant, or arrested in growth, owing to self-produced carbon dioxide.
- autonomic (ôt'önŏm'îk) a. [Gk. autos, self; nomos, law.] Autonomous; self-governing, spontaneous; appl. the involuntary nervous system as a whole, comprising parasympathetic and sympathetic systems; induced by internal stimuli, as movements of development, growth, unfolding, etc., opp. paratonic; internal, appl. environment, opp. choronomic.

autopalatine (ôt opăl'ătin) n. [Gk. autos, self; L. palatum, palate.] In a few teleosts, an ossification at anterior end of pterygoquadrate.

autoparasite (ô'töpār'asīt) n. [Gk. autos. self; parasitos, one who subsists on another.] A parasite growing on another parasite.

- autoparthenogenesis (ôt'öpâr'thënöjĕn'esis) n. [Gk. autos, self; parthenos, virgin; genesis, descent.] Development from unfertilised eggs activated by a chemical or physical stimulus.
- autophagous (ôtŏf'agus) a. [Gk. autos, self; phagein, to eat.] Appl. birds capable of running about and securing food for themselves when newly hatched.

autophilous (ôtôf'îlŭs) a. [Gk. autos, self; philein, to love.] Self-pollinating; autogamous.

autophya (ôt'öfi'ā) n. plu. [Gk. autos, self; phyein, to produce.]
Elements in formation of shell secreted by animal itself; cf. xenophya.

autophyllogeny (ôt'öfilŏj'eni) [Gk. autos, self; phyllon. leaf; genos, birth.] Growth of one leaf upon or out of another.

autophyte (ôt'öfīt) n. [Gk. autos, phyton, plant.] A selfnourished plant; plant nourished directly by inorganic matter; cf. saprophyte.

autophytic (ôtöfĭt'ĭk) a. [Gk. autos, self; *phyton*, plant.] Autotrophic,

q.v.; pert. autophytes.

autoplasma (ôt'öplăz'mă) n. ſGk. autos, self; plasma, mould.] Plasma from same animal used as medium for tissue culture; homoplasma, heteroplasma.

autoplastic (ôt'öplăs'tĭk) a. [Gk. autos, self; plastos, formed.] Appl. graft to another position in the same

individual.

autopodium (ôt'öpō'dĭŭm) n. autos, self; pous, foot.] The hand or foot.

- autopolyploid (ô'töpŏl'ĭploid) [Gk. autos, self; polys, many; eidos, aploos, onefold; form.] An organism having more than two sets of homologous chromosomes.
- autopotamic (ô'töpŏtăm'ĭk) a. [Gk. autos, self; potamos, river.] Thriving in a stream, not in its backwaters; appl. potamoplankton.
- autoradiography (ô'torādiŏg'răfi) n. [Gk. autos, self; L. radius, ray; Gk. graphein, to write.] Method of demonstrating the presence specific chemical substances by first making them radioactive, then recording on a photographic film their distribution in the body, organs, or tissues.

autoskeleton (ôt'öskěl'ëtŏn) n. [Gk. autos, self: skeletos, dried.] A true skeleton formed within the animal.

- autosome (ôt'ösōm) n. [Gk. autos. soma, body.] self; A typical chromosome, or euchromosome; opp. sex-chromosome; cf. allosome.
- autospore (ô'töspōr) n. [Gk. autos, self; sporos, seed.] An aplanospore which resembles the parent cell.

autostoses (ôt'ŏstō'sēz) n. plu. [Gk. autos, self; osteon, bone.] Bones formed in cartilage; cf. allostoses.

autostylic (ôt'östīl'īk) a. [Gk. autos, self; stylos, pillar.] With mandibular arch self-supporting, articulating directly with skull; hyostylic.

autosynapsis (ô'tösĭnăp'sĭs) n. [Gk. autos, self; synapsis, union.] Auto-

syndesis.

- autosyndesis (ô'tösĭn'dësĭs) n. [Gk. autos, self; syndesis, a binding together.] Pairing of chromosomes from the same parent, in a polyploid or allopolyploid; pairing of homogenetic chromosomes; allosyndesis.
- autotilly (ô'tötĭl'ĭ) n. [Gk. autos, self; tillesthai. to pluck.] Autotomy, as in certain spiders.
- autotomy (ôtŏt'ömĭ) n. [Gk. autos, self; tome, cutting.] Self-amputation of a part, as in certain worms, arthropods, and lizards.
- autotransplantation, transplantation of tissue or organ to another part of same organism; cf. homoiotransplantation.
- autotrophic (ôt'ötrŏf'ĭk) a. [Gk. autos, self; trephein, to nourish.] Procuring food independently; appl. plants which form carbohydrates and proteins from carbon dioxide and inorganic compounds; neither saprophytic nor parasitic; autophytic. Opp. heterotrophic.
- autotropism (ôtěť röpřem) n. [Gk. autos, self; trope, turn.] Tendency to grow in a straight line; appl. plants unaffected by external influence; tendency of organs to resume original form, after bending or straightening due to external factors; rectipetality.

autoxenous (ôt'özĕn'ŭs) a. [Gk. autos, self; xenos, host.] Parasitic on the same host at different stages in life-history; autoe-

cious.

autozooid (ôt'özō'oid) n. [Gk. autos, self; zoon, animal; eidos, form.] An independent alcyonarian zooid or individual.

- auxenolonic acid,—auxin B, q.v. auxentriolic acid,—auxin A, q.v.
- auxesis (ôksē'sīs) n. [Gk. auxesis, growth.] Growth; increase in size owing to increase in cell size; induction of cell division; cf. merisis.
- auxetic (öksĕt'ik) n. [Gk. auxein, to increase.] Any agent which induces cell-division. a. Stimulating cell proliferation.
- auxilia (ôgzīl'yă) n. plu. [L. auxilium, assistance.] Two small sclerites between unguitractor and claws, in insects.
- auxiliary cells,—two or more cells adjoining guard cells, or surrounding stomata; accessory or subsidiary cells.

auximone (ôk'simōn) n. [Gk. auximos, promoting growth.] An accessory growth-stimulating factor in food of plants.

- auxins (ôk'sinz) n. plu. [Gk. auxein, to increase.] Growth-regulating hormones of plants; auxin A isolated from growing tips of oat seedlings and human urine, C₁₈H₃₂O₅; auxin B, from vegetable sources and urine, accelerates mycelium growth, C₁₈H₃₀O₄; heteroauxin (q.v.), and a number of other substances.
- auxocyte (ôks'ösīt) n. [Gk. auxein, to increase; kytos, hollow.] Androcyte, sporocyte, oocyte, or spermatocyte at growth period.

auxospireme (ôks'öspī'rēm) n. [Gk. auxein, to increase; speirema, coil.]
Spireme formed after syndesis.

- auxospore (ôk'söspōr) n. [Gk. auxein, to increase; sporos, seed.]
 Zygote of diatoms, formed by union of two individuals at limit of decrease in size.
- auxotonic (ôk'sötŏn'îk) a. [Gk. auxein, to increase; tonos, strain.] Induced by growth; appl. movements of immature plants; cf. allassotonic; appl. contraction against an increasing resistance.
- avicularium (āvīkūlā'rīūm) n. [L. avicula, dim. of avis, bird.] In Polyzoa a modified zooecium with

muscular movable attachments resembling a bird's beak.

avifauna (ā'vifô'na) n. [L. avis, bird; Faunus, rural deity.] All the bird species or birds of a region or period; ornis.

- avitaminosis (ăvī'tămīnō'sis) n. [L. a, from; vita, life; ammoniacum, resinous gum.] A condition or disease resulting from vitamindeficiency.
- awn (ôn) n. [Icel. ögn, chaff.] The 'beard' of grasses; point of leaf, in certain Lycopsida.
- axial (ăk'siăl) a. [L. axis, axle.] Pert. axis or stem.
- axial filament,—central filament, as of a stiff radiating pseudopodium or of a flagellum.
- axial sinus,—a nearly vertical canal in echinoderms, opening into internal division of oral ring sinus, and communicating with stone canal.
- axial skeleton,—skeleton of head and trunk, opp. appendicular skeleton.
- axiate pattern,—arrangement of parts with reference to a definite axis.
- axil (ăk'sĭl) n. [L. axilla, arm-pit.] The angle between leaf or branch and axis from which it springs.
- axile [ăk'sīl) a. [L. axis, axle.] Pert., situated in, or belonging to the axis; appl. placentation, free central, q.v.
- axilemma (ăk'sĭlĕm'ă) n. [L. axis, axle; Gk. lemma, husk.] In medullated nerve fibres, the sheath surrounding axial cylinder.
- axilla (ăksĭl'ā, āk'sĭlā) n. [L. axilla, arm-pit.] The arm-pit; an axil.
- axillary (ăk'silări, ăksīl'āri) a. [L. axilla, arm-pit.] Pert. axil; growing in axil, as buds; pert. arm-pit; appl. seventh longitudinal or anal vein of insect wing. n. One of the pteralia, q.v.

axinost,—axonost, q.v.

axipetal (ăksip'ētāl) a. [L. axis, axle; petere, to seek.] Passing towards attachment of axon, appl. nerve impulses. axis (åk'sis) n. [L. axis, axle.] The main stem or central cylinder; the fundamentally central line of a structure; rachis of trilobites; epistropheus, q.v.; structure at base of insect wing.

axis cylinder,—the central tract of a nerve fibre, the impulse transmitter; axon and its myelin sheath.

axodendritic (āk'söděndrit'ik) a. [Gk. axon, axle; dendron, tree.] Appl. synapse in which end-brush of

axon is in contact with dendritic

processes.

axon (ăk'sŏn) n. [Gk. axon, axle.]
The axis-cylinder process of a
nerve-cell normally transmitting
excitations from its cell body;
axone, neuraxon, neurite.

axon hill or hillock,—cone of origin,

q.v

- axoneme (ăk'sönēm) n. [Gk. axon, axle; nema, thread.] A thread of strand forming infusorian stalk; an axostyle; the axial filament of a flagellum; axial thread or genoneme of a chromosome.
- axonost (ăk'sönŏst) n. [Gk. axon, axle; osteon, bone.] The basal portion of rods supporting dermotrichia of fin-rays; axinost; interspinal.
- axoplast (ăk'söplăst) n. [Gk. axon, axle; plastos, formed.] A filament extending from kinetoplast to end of body in some trypanosomes.

axopodium (ăk'söpo diŭm) n. [Gk. axon, axle; pous, foot.] A pseudopodium with axial filament.

axosomatic (ăk'sösōmăt'ik) a. [Gk. axon, axle; soma, body.] Appl. synapse in which end-brush of axon terminates about nerve-cell body.

axospermous (ăk'söspěr'mŭs) a. [Gk. axon, axle; sperma, seed.]

With axile placentation.

axostyle (āk'söstīl) n. [Gk. axon, axle; stylos, pillar.] A slender flexible rod of organic substance forming a supporting axis for the body of many Flagellata.

azoic (ăzō'îk) a. [Gk. a, without; zoikos, pert. life.] Uninhabited; without remains of organisms or

of their products; appl. Pre-Cambrian era or rocks.

azurophil (åzū'röfil, āzh'ūröfil) a.
[F. azur, from Ar. al azurd, lapis lazuli; Gk. philein, to love.]
Staining readily with blue aniline dyes.

azygobranchiate (ăz'īgöbrăng'kiāt)
 a. [Gk. a, without; zygon, yoke;
 brangchia, gills.] Having gills or ctenidia not developed on one side.

azygoid (az'īgoid) a. [Gk. a, without; zygon, yoke; eidos, form.] Not zygoid; haploid; appl. parthenogenesis.

azygomatous (åzīgŏm'átŭs) a. [Gk. a, without; zygoma, a bar.] Without a zygoma or cheek-bone arch.

azygomelous (ăz'īgŏměl'ūs, a. [Gk. a, without; zygon, yoke; melos, limb.] Having unpaired appendages; appl. fin of Acrania and Cyclostomata.

azygos (ăzī'gŏs) n. [Gk. a, without; zygon, yoke.] An unpaired muscle,

artery, vein, process.

azygosperm (az'īgöspērm), azygospore (az'īgöspēr) n. [Gk. a, without; zygon, yoke; sperma, sporos, seed.] A spore developed directly from a gamete without conjugation; parthenospore.

azygote (äzī'gōt) n. [Gk. a, without; zygon, yoke.] An organism resulting from haploid parthenogenesis.

azygous (az'igŭs) a. [Gk. a, without; zygon, yoke.] Unpaired.

В

Babes-Ernst bodies,—metachromatic or volutin granules, in bacteria. bacca (băk'ă) n. [L. bacca, berry.]

A pulpy fruit; berry.

baccate (băk'āt) a. [L. bacca, berry.]

Pulpy, fleshy; berried.

bacciferous (băksĭf'ërŭs) a. [L. bacca, berry; ferre, to bear.]
Berry-producing, or -bearing.

bacciform (băk'sifôrm) a. [L. bacca, berry; forma, shape.] Berry-shaped. bacillary (băsil'āri) a. [L. bacillum, small staff.] Appl. layer of rods and cones of retina.

- bacillus (băsĭl'ŭs) n. [L. bacillum, small staff.] A rod-like bacterium; a single-celled fungus.
- back-cross,-to mate a cross or hybrid to a member of one of the parental stocks; a resulting hybrid.

back mutation,—reversion of a mutant gene to its original state; reverse mutation.

bactericidin (băk'terisī'din) n. [Gk. bakterion, small rod; L. caedere, to kill.] A substance that kills bacteria without causing lysis.

bacteriochlorin (băktē'rĭöklō'rĭn) n. [Gk. bakterion, small rod; chloros, green.] Green pigment, related to chlorophyll, in sulphur bacteria.

bacteriochlorophyll (băktē'rĭöklō'röfil) n. [Gk. bakterion, small rod; chloros, green; phyllon, leaf. A photosynthetic pigment of bacteria, from which chlorophyll-a may be derived; bacteriochlorin.

bacteriology (băk'tēriŏl'öji) n. [Gk. bakterion, small rod; logos, discourse.] The science dealing with

bacteria.

- bacteriolysin (băktē'riölī'sin) n. [Gk. bakterion, small rod; lysis, loosing.] A substance which causes dissolution of bacteria.
- bacteriolysis (băk'tēriŏl'isis) n. [Gk. bakterion, small rod; lysis, loosing.] The disintegration and dissolution of bacteria.

bacteriophage (băktē'rĭöfāj') [Gk. bakterion, small rod; phagein, to devour.] A destroyer of bacteria; a bacteriolytic agent; phage.

(bāktē'rĭöpŭr'bacteriopurpurin pūrin) n. [Gk. bakterion, small L. purpura, purple.] complex of photosynthetic pigments causing the red, purple, or violet appearance of certain bacteria.

bacteriostatic (băktē'rĭöstāt'īk) a. [Gk. bakterion, small rod; statikos, causing to stand.] Inhibiting de-

velopment of bacteria.

bacteriotropin (băk'tērio'tropin) n. [Gk. bakterion, small rod; trope, An ingredient of blood serum which renders bacteria more readily phagocytable; opsonin.

- bacteroid (băk'tëroid) n. [Gk. bakterion, small rod; eidos, form.] An irregular form of certain bacteria.
- baculiform (băk'ūlifôrm) a. baculum, rod; forma, shape.] Rodshaped; appl. chromosomes; appl. ascospores.
- baculum (băk'ūlŭm) n. [L. baculum, rod.] The penis bone; os pri-
- Baillarger's line [J. F. G. Baillarger, French neurologist]. Outer and inner layer of white fibres parallel to surface of cerebral cortex.
- balanced lethals,—heterozygotes in which different lethal genes are in such close proximity on a pair of homologous chromosomes that is usually no there crossing over.
- balancers (băl'ănsërz) n. plu. [L. bilanx, having two scales. | Halteres or poisers of Diptera; paired larval head appendages functioning as props until forelegs are developed in certain salamanders.
- balanoid (băl'ănoid) a. [Gk. balanos, acorn; eidos, like.] Pert. barnacles; acorn-shaped.
- balanus (băl'ănus) n. [L. balanus, acorn.] Glans penis; a genus of barnacles.
- balausta (bălôs'tă) n. [Gk. balaustion, blossom.] A many-celled, manyseeded, indehiscent fruit with tough pericarp; fruit of pomegranate.
- **baleen** (bălēn') n. [L. balaena, whale.] Horny plates attached to upper jaw of true whales; whalebone.
- ballast (băl'ăst) n. [Sw. barlast.] Appl. elements present in plants and which are not apparently essential for growth, e.g., Al, Si.
- ballistospores (bălĭs'töspōrz) n. plu. [Gk. ballein, to throw; sporos, seed.] Asexual spores, formed on sterigmata and suddenly discharged with excretion of droplet, in Sporobolomycetes.

balsamiferous (băl'sămif'ërus) a. [L. balsamum, balsam; ferre, to bear.] Producing balsam.

bar of Sanio,—crassula, q.v.

49 BAR-BAS-

- baraesthesia (băr'ēsthē'zĭā) n. [Gk. baros, weight; aisthesis, sensation.] The sensation of pressure.
- barb (bârb) n. [L. barba, beard.] One of delicate thread-like structures extending obliquely from a feather rachis, and forming the vane; a hooked hair-like bristle.
- barbate (bâr'bāt) a. [L. barbatus, bearded.] Bearded; having hair tufts.
- barbel (bâr'bĕl) n. [L.L. barbellus, barbel.] A tactile process arising from the head of various fishes.
- barbellate (bârběl'āt, bâr'bëlāt) a. [L. barba, beard.] With stiff hooked hair-like bristles; appl. pappus.
- barbicel (bâr'bisĕl) n. [L. barba, beard.] Small process on a feather barbule.
- barbula (bâr'būlă) n. [L. barbula, dim. of barba, beard.] Row of teeth in peristome of certain mosses.
- barbule (bâr'būl) n. [L. barbula, dim. of barba, beard.] One of small hooked processes fringing barbs of feather; appendage of lower jaw in some teleosts.
- bark (bârk) n. [Dan. bark.] The tissues external to the vascular cambium, collectively; phloem. cortex, and periderm; outer dead tissues and cork.
- barotaxis (bărötăk'sis) n. [Gk. baros, weight; taxis, arrangement.] The reaction to a pressure stimulus.
- Bartholin's duct [C. Bartholin, jr., Danish anatomist]. The larger duct of the sublingual gland.
- Bartholin's glands,—the greater vestibular glands on each side of vagina, homologues of male bulbourethral glands.
- basal (bā'săl) a. [L. basis, base.] Pert., at, or near the base.
- **basal bone,**—os basale, basale, q.v.basal ganglia,—ganglia connecting cerebrum with other centres.
- basal granule,—a thickening, or body, at base of a flagellum in certain protozoa.
- basal knobs,—swellings or granules at points of emergence of cilia in ciliated epithelial cells.

- basal leaf, one of the leaves produced near base of stem.
- basal metabolic rate, --- rate of metabolism of a resting organism, expressed as percentage of normal heat production per hour square metre surface area.
- basal metabolism,-standard metabolism, tissue activity or physicochemical changes of a resting organism.
- basal placenta,-arises from proximal end of ovary.
- plates,-certain plates in echinoderms, situated at or near top of stalk in crinoids, in echinoids forming part of apical disc; fused parachordal plates in skull development; of placentae, outer wall of intervillous space.
- basal wall,—the first plane of division of oospores of ferns and mosses. basalar (bāsā'lăr) a. [L. basis, base; ala, wing.] Appl. sclerites below
- wing base in insects. basale (băsā'lē) n. [L. basis, base.] A bone of variable structure arising from fusion of pterygrophores and supporting fish fins; os basale, the fused basioccipital and parasphenoid in Gymnophiona.
- basement membrane,-the membrane of a gland containing acini or special secreting portions.
- baseost (bā'seŏst) n. [Gk. basis, base; osteon, bone.] Distal element of pterygiophore of teleosts.
- basialveolar (bā'sĭāl'vēölār) a. basis, base; alveolus, small pit.] Extending from basion to centre of alveolar arch.
- basibranchial (bā'sibrang'kial) n. [Gk. basis, base; brangchia, gills.] Median ventral or basal skeletal portion of branchial arch.
- basic (bā'sīk) a. [Gk. basis, base.] Appl. stains which act in general on nuclear contents of cell; cf. acidic; appl. number, the minimum haploid chromosome number occurring in a series of euploid species of a chromosome number in gametes of diploid ancestor of a polyploid organism.

basichromatin (bā'sīkrō'mātīn) n. [Gk. basis, base; chroma, colour.] The deeply staining substance of nuclear network; chromatin.

basiconic (bā'sikon'ik) a. [Gk. basis, base; konos, cone.] Having, or consisting of, a conical process above general surface; appl. sensillae.

basicoxite (bā'sĭkŏks'īt) n. [L. basis, base; coxa, hip.] Basal ring of coxa.

basicranial (bā'sikrā'nĭāl) a. [Gk. basis, base; kranion, skull.] Situated at or relating to base of skull.

basidia,—plu. of basidium.

basidial (băsid'iăl) a. [Gk. basis, base; idion, dim.] Pert. basidia or a basidium.

basidiocarp (băsīd'iökârp) n. [Gk. basis, base; idion, dim.; karpos, fruit.] The fruit-body of Basidiomycetes.

basidiophore (băsīd'iöför) n. [Gk. basis, base; idion, dim.; pherein, to bear.] A sporophore which carries basidia.

basidiospore (băsĭd'ĭöspōr) n. [Gk. basis, base; idion, dim.; sporos, seed.] A spore or gonidium abstricted from a basidium; a secondary conidium; a basidiogonidium.

basidium (băsid'iŭm) n. [Gk. basis, base; idion, dim.] A special cell or row of cells, of certain fungi, forming spores by abstriction.

basidorsal (bā'sidôr'săl) a. [L. basis, base; dorsum, back.] Appl. small cartilaginous neural plate.

basifixed (bā'sĭfīksd) a. [L. basis, base; figere, to make fast.] Attached by base; innate, having filament attached to anther base.

basifugal (bāsif'ūgăl) a. [L. basis, base; fugere, to flee.] Growing away from base.

basifuge (bā'sifūj) n. [L. basis, base; fugere, to flee.] A plant unable to tolerate basic soils; calcifuge. a. Oxyphilous.

basigamous (bāsīg'āmūs) a. [Gk. basis, base; gamos, marriage.]

Having oosphere reversed in embryo-sac.

basigynium,—podogynium, q.v.

basihyal (bā'sihī'āl) n. [Gk. basis, base; hyoeides, Y-shaped.] Broad median plate, the basal or median ventral portion of hyoid arch.

basilabium (bā'sīlā'bīum) n. [L. basis, base; labium, lip.] Sclerite formed by fusion of labiostipites in insects.

basilar (bās'īlār) a. [L. basis, base.] Pert. near or growing from base; as artery, crest, membrane, plexus, plate, process, style.

basilic (băsil'îk) a. [Gk. basilikos, royal.] Appl. a large vein on inner

side of biceps of arm.

basilingual (bā'siling'gwăl) a. [L. basis, base; lingua, tongue.] Appl. a broad cartilaginous plate, the body of the hyoid, in crocodiles, turtles, and amphibians.

basimandibula (bā'sīmāndīb'ūlā) n. [L. basis, base; mandibulum, lower jaw.] A small sclerite, on insect head, at base of mandible.

basimaxilla (bā'sīmāksīl'ā) n. [L. basis, base; maxilla, upper jaw.] A sclerite at base of maxilla in insects.

basioccipital (bā'siŏksip'ītāl) n. [L. basis, base; occiput, back of head.] The median basilar bone or element in occipital region of skull.

basion (bā'siŏn) n. [Gk. basis, base.]
The middle of anterior margin of foramen magnum.

basiophthalmite (bā'siŏfthăl'mīt) n.
[Gk. basis, base; ophthalmos, eye.]
The proximal joint of eye-stalk in crustaceans.

basiotic (bā'siŏt'ik) a. [Gk. basis, base; ous, ear.] Mesotic, q.v.

basipetal (bāsip'ētăl) a. [L. basis, base; petere, to seek.] Developing from apex to base; appl. leaves and inflorescences.

basipharynx (bā'sĭfăr'ingks) n. [Gk. basis, base; pharyngx, gullet.] In insects, epipharynx and hypopharynx united.

basiphil (bā'sifil) a. [Gk. basis, base;
philein, to love.] Basophil, q.v.
n. A basiphil cell; a mast cell, q.v.

51 B-CO-BAS-

basipodite (bāsīp'ödīt) n. [Gk. basis, base; pous, foot.] The second or distal joint of the protopodite of certain limbs of Crustacea; trochanter of spiders.

basipodium (bā'sĭpō'dĭŭm) n. [Gk. basis, base; pous, foot.] Wrist or

ankle.

basiproboscis (bā'sĭpröbŏs'ĭs) n. [Gk. basis, base; proboskis, trunk.] Membranous portion of proboscis of some insects, consisting of mentum, submentum, and maxillary cardines and stipites.

basipterygium (bā'sītërīj'ĭŭm) n. [Gk. basis, base; pterygion, little wing.] A large flat triangular bone in pelvic fin of teleosts, and a bone or cartilage in other fishes.

basipterygoid (bā'sĭtĕr'ĭgoid) [Gk. basis, base; pteryx, wing; eidos, form.] A process of the basisphenoid in some birds.

basiscopic (bā'sĭskŏp'ĭk) a. [Gk. basis, base; skopein, to view.] Facing towards the base; opp, acroscopic.

basisphenoid (bā'sīsfē'noid) n. [Gk. basis, base; sphen, wedge; eidos, form.] Cranial bone between basioccipital and presphenoid.

basisternum (bā'sĭstĕr'nŭm) [L. basis, base; sternum, breast-The principal sclerite of insect sternum; antesternite, eusternum.

basistyle (bā'sīstīl) n. [Gk. basis, base; stylos, pillar.] Proximal part or coxite of gonostyle in mosquitoes; cf. dististyle.

basitarsus (bā'sĭtâr'sŭs) n. [Gk. basis, base; tarsos, sole of foot.] Proximal tarsomere or tarsus' of spiders. Cf. telotarsus.

basitemporal (bā'sĭtěm'pörăl) n. [L. basis, base; tempora, temples.] A broad membrane bone covering basisphenoidal region of skull.

basitonic (bā'sĭtŏn'ĭk) a. [Gk. basis, base; tonos, brace.] Having anther united at its base with rostellum; basitonous. Opp. acrotonic.

basivertebral (bā'sĭvēr'tēbrăl) a. [L. basis, base; vertebra, vertebra.] Appl. veins within bodies of vertebrae and communicating vertebral plexuses.

basket cells, -- myo-epithelial cells surrounding glandular cells; cerebellar cortical cells with axon branches surrounding Purkinje cells.

[Gk. basis, basophil (bā'söfil) a. base; philein, to love.] Having a strong affinity for basic stains; also basiphil, basiphilic, basophile, basophilic, basophilous. n. A cell which stains with basic dyes.

bast (băst) n. [A.S. baest, bast.] The inner fibrous bark of certain trees: liber.

bastard merogony,—activation of an enucleated egg fragment by spermatozoon of a different species.

bastard wing,—the alula or ala spuria, consisting of three quill feathers borne on first digit of bird's wing.

bathylimnetic (băth'îlĭmnět'ik) a. [Gk. bathys, deep; limnetes, living in marshes.] Living or growing in the depths of lakes or marshes.

bathymetric (băth'imět'rĭk) a. [Gk. bathys, deep; metron, measure.] Pert. vertical distribution of organisms in space.

bathypelagic (băth'ipělăj'ik) a. [Gk. bathys, deep; pelagos, sea.] Pert., or inhabiting, the deep sea.

bathysmal (băthĭz'măl) a. [Gk. bathys, deep.] Pert. deepest depths of the sea.

batonette (bătönět) n. [F. bâtonnet, small stick.] An element of the Golgi apparatus, q.v.

batrachian (bătrā'kĭăn) a. batrachos, frog.] Relating to frogs and toads.

B-cells,—beta cells of islets Langerhans.

B-chromosome, — supernumerary chromosome in maize.

B-complex,—a group of accessory food factors comprising thiamine or vitamin B_1 , riboflavin (B_2) , pantothenic acid (B₃), niacin or P-P factor, pyridoxin (B₆), biotin (H), inositol, choline, para-amino benzoic acid, and folic acid (M), and B_{12} anti-pernicious-anaemia factor. bdelloid (děl'oid) a. [Gk. bdella, leech; eidos, form.] Having the appearance of a leech.

BDE-

beard (berd) n. [A. S. beard, beard.]
Any of the arrangements of hairs which resemble a man's beard, on heads of animals; barbed or bristly hair-like outgrowths on grain; awn.

bedeguar (běď'ēgăr) n. [From Persian through F. bédeguar, thistle.] A mossy gall formed on rose-bushes.

behaviorism (bēhā'viörizm) n. [A.S. behabban, to hold in.] Theory that the manner in which animals act may be explained in terms of conditioned neuromotor and glandular reactions.

belemnoid (běl'ěmnoid, bělěm'noid)

a. [Gk. belemnon, dart; eidos, form.] Shaped like a dart; appl. styloid process.

Bellini's ducts [L. Bellini, Italian anatomist]. Tubes opening at apex of kidney papilla, and formed by union of smaller straight or collecting tubules.

bell nucleus,—a solid mass of cells, derived from ectoderm and lying between ordinary ectoderm and mesogloea at apex of medusoid bud.

belonoid (běl'önoid) a. [Gk. belone, needle; eidos, form.] Shaped like a needle; styloid.

benthic (ben'thik) a. [Gk. benthos, depths of sea.] Pert., or living on, sea-bottom; benthal.

benthopotamous (běn'thöpŏt'āmūs)
a. [Gk. benthos, depths; potamos, river.] Pert., growing, or living, on bed of a river or stream.

benthos (běn'thôs) n. [Gk. benthos. depths of sea.] The fauna and flora of the sea-bottom.

Berlese's organ [A. Berlese, Italian zoologist]. A glandular organ in haemocoel on right side of female abdomen in Cimex, secreting during passage of spermatozoa to spermatheca.

berry (ber'i) n. [A.S. berie, berry.] Inferior, indehiscent, many-seeded fruit, usually with fleshy pericarp; egg of lobster, or crayfish; dark knob-like structure on bill of swan.

beta (β) cells,—basophil cells in pars glandularis of pituitary gland; cells with alcohol-soluble granules, in islets of Langerhans, B cells.

beta (β) granules,—granules in peripheral region of protoplast, a protein reserve in blue-green algae; cyanophycin.

betaine (be'taĭen) n. [L. beta, beet.]

A basic decomposition product of lecithin, occurring in beet and other plants, and in animals; C₅H₁₁O₂N.

Betz cells [V. A. Bets, Russian histologist]. Giant pyramidal cells in motor area of cerebral cortex.

B-horizon,—the lower, illuvial soil layers.

biacuminate (bī'ākū'mĭnāt) a. [L. bis, twice; acumen, point.] Having two tapering points.

biarticulate (bī'artīk'ūlāt) a. [L. bis, twice; articulus, joint.] Twojointed.

bicapsular (bīkāp'sūlăr) a. [L. bis, twice; capsula, little box.] Having two capsules or vessels.

bicarinate (bīkār'ināt) a. [L. bis, twice; carina, keel.] With two keel-like processes.

bicarpellate (bikâr'pelāt) a. [L. bis, twice; Gk. karpos, fruit.] With two carpels; bicarpellary.

bicaudate (bīkô'dāt) a. [L. bis, twice; cauda, tail.] Possessing two tail-like processes.

bicellular (bīsēl'ūlăr) a. [L. bis, twice; cellula, little cell.] Composed of two cells.

bicentric (bī'sĕntrīk) a. [L. bis, twice; centrum, centre.] Pert. two centres; appl. distribution of species, etc., discontinuous owing to alteration in the intervening

biceps (bī'sēps) n. [L. bis, twice; caput, head.] A muscle with two heads or origins, as biceps brachii and femoris.

biciliate (bīsīl'īāt) a. [L. bis, twice; cilium, eyelash.] Furnished with two cilia.

bicipital (bīsĭp'ītăl) a. [L. bis, twice; caput, head.] Pert. biceps; appl. fascia, or lacertus fibrosus, an aponeurosis of distal tendon of the biceps brachii; a groove, the intertubercular sulcus, on upper part of humerus; ridges, the crests of the greater and lesser tubercles of the humerus; appl. a rib with dorsal tuberculum and ventral capitulum.

bicollateral (bikolat'eral) a. [L. bis, twice; con, together; latus, side.] Having the two sides similar; appl. a vascular bundle with phloem on both sides of xylem, as in Cucurbitaceae and Solanaceae.

bicolligate (bīkŏl'igāt) a. [L. bis, twice; cum, together; ligare, to bind.] With two stretches of webbing on the foot.

biconjugate (bīkŏn'joogāt) a. bis, twice; cum, with; jugum, yoke.] With two similar sets of pairs.

bicornute (bīkôrnūt') a. [L. bis, twice; cornutus, horned.] With two horn-like processes.

bicostate (bīkŏs'tāt) a. [L. bis, twice, costa, rib.] Having two longitudinal ridges or ribs, as a leaf.

bicrenate (bīkrē'nāt) a. [L. bis, twice; crena, notch.] Doubly crenate, as crenate leaves with

notched toothed margins.

bicuspid (bīkŭs'pid) a. [L. bis, twice; cuspis, point.] Having two cusps or points; appl. valve consisting of anterior and posterior cusps attached to circumference of left atrioventricular orifice, mitral valve; appl. teeth, premolar.

bicyclic (bīsīk'līk) a. [L. bis, twice; Gk. kyklos, circle.] Arranged in

two whorls.

Bidder's ganglia [F. H. Bidder, Estonian anatomist]. A collection of nerve-cells in region of the auriculo-ventricular groove.

rudimentary Bidder's organ,—a ovary attached to anterior end of generative organs in the toad.

bidental (bīděn'tăl) a. [L. bis, twice; dens, tooth.] Having two teeth, or tooth-like processes; bidentate.

bidenticulate (bī'děntik'ūlāt) a. [L. bis, twice; dim. of dens, tooth.] With two small teeth or tooth-like processes, as some scales.

biennial (bīĕn'ĭăl) a. [L. bis, twice; annus, year.] Lasting for two

years.

bifacial (bīfā'siăl, bīfā'shăl) a. [L. bis, twice; facies, face.] Appl. leaves with distinct upper and lower surfaces; dorsiventral.

bifarious (bīfā'rĭŭs) a. [L. bis, twice; fariam, in rows.] Arranged in two rows, one on each side of axis.

bifid (blf'id) a. [L. bis, twice; findere, to split.] Forked, opening with a median cleft; divided nearly to middle line.

biflabellate (bī'flăběl'āt) a. [L. bis, twice; flabellum, fan.] Doubly flabellate, each side of antennal joints sending out flabellate pro-

biflagellate (bīflăj'elāt) a. [L. bis, twice; flagellum, whip] Having two flagella.

biflex (bī'flěks) a. [L. bis, twice; flectere, to bend.] Twice curved.

biflorate (bīflo'rāt) a. [L. bis, twice: flos, flower.] Bearing two flowers.

bifoliar (bīfō'lĭăr) a. [L. bis, twice; folium, leaf.] Having two leaves. bifoliate (bīfō'lĭāt) a. [L. bis, twice;

folium, leaf.] Appl. palmate compound leaf with two leaflets.

biforin (bif'orin) n. [L. bis, twice; foris, door.] An oblong raphidian cell opening at each end.

biforous (bif'orus) a. [L. bis, twice; foris, door.] With two openings; appl. spiracles in larvae of certain beetles.

[L. bis, bifurcate (bīfŭr'kāt) a. twice; furca, fork.] Forked; having two prongs; having two joints, the distal V-shaped and attached by its middle to the proximal.

bigeminal (bījěm'īnăl) a. [L. bis, twice; geminus, double.] With structures arranged in double pairs; appl. arrangement of pore-pairs in two rows, in ambulacra of some echinoids; cf. corpora bigemina.

bigeminate (bījem'īnāt) a. [L. bis, twice; geminus, double.] Doublypaired; twin-forked.

bigeneric (bī'jēnēr'ik) a. [L. bis, twice; genus, race.] Appl. hybrids between two distinct genera.

bijugate (bijoo'gāt) a. [L. bis, twice; jugare, to join.] With two pairs of leaflets.

bilabiate (bīlā'bĭāt) a. [L. bis, twice; labium, lip.] Two-lipped; appl. calyx, corolla, dehiscence.

bilamellar (bīlamel'ar) a. [L. bis, twice; lamella, plate.] Formed of two plates; having two lamellae.

(bīlăm'ĭnăr) bilaminar a. bis, twice; lamina, thin plate.] Having two plate-like layers; diplo-

bilateral (bīlăt'ërăl) a. [L. bis, twice; latus, side.] Having two sides symmetrical about an axis.

bile (bīl) n. [L. bilis, bile.] The secretion of the liver, passing to duodenum and assisting digestion.

biliary (bil'iări) a. [L. bilis, bile.]

Conveying or pert. bile.

bilicyanin (bil'isī'anin) n. [L. bilis, bile; Gk. kyanos, dark blue.] A blue pigment resulting from oxidation of biliverdin.

bilipurpurin (bĭl'ĭpŭr'pūrĭn) n. [L. purpura, bilis, bile : purple.]

Phylloerythrin.

bilirubin (bil'iroo'bin) n. [L. bilis, bile; ruber, red.] A reddishyellow pigment of bile and blood, end-product of hæmoglobin metabolism; so also bilifulvine, biliphaein; $C_{32}H_{36}N_4O_6$.

biliverdin (bil'iver'din) n. [L. bilis, bile; F. vert, green.] A green bile pigment formed by oxidation of bilirubin; $(C_{16}H_{18}N_2O_4)_n$.

bilobate (bīlō'bāt) a. [L. bis, twice; L.L. lobus, from Gk. lobos, rounded flap.] Having two lobes.

bilobular (bīlob'ūlar) a. [L. bis. twice; L. lobulus, dim. of lobus, lobe.] Having two lobules.

bilocellate (bī'lösĕl'āt) a. [L. bis, twice; locellus, dim. of locus, Divided into two complace.] partments.

bilocular (bīlŏk'ūlăr), biloculine (bīlok'ūlin) a. [L. bis, twice; locus, place.] Containing two cavities or chambers; cf. loculus.

bilophodont (bīlŏf'ödŏnt) a. [L. bis, twice; Gk. lophos, ridge; odous, tooth.] Appl. molar teeth of tapir, which have ridges joining the two anterior and two posterior cusps.

bimaculate (bīmāk'ūlāt) a. [L. bis, twice; macula, spot.] Marked with two spots or stains.

bimanous (bim'ănus) a. [L. bis, twice; manus, hand.] Having two hands; appl. certain Primates.

bimastism (bīmăs'tĭzm) n. [L. bis, twice; Gk. mastos, breast.] Condition of having two mammae.

bimuscular (bīmus'kūlar) a. [L. bis, twice; musculus, muscle.] Having two muscles.

binary fission,—division of a cell into two by an apparently simple division of nucleus and cytoplasm.

binary nomenclature, -- binomial nomenclature, q.v.

binate (bī'nāt) a. [L. bini, two by two.] Growing in pairs; appl. leaf composed of two leaflets.

binocular (binok'ūlar) a. [L. bini, pair; oculus, eye.] Having or pert. two eyes; stereoscopic, appl. vision.

binodal (bīnō'dăl) a. [L. bis, twice; nodus, knob.] Having two nodes.

as stem of plant.

binomial (bīno'mial) a. [L. bis, twice; nomen, name.] Consisting of two names; appl. nomenclature, the system of double names given to plants and animals,—first generic name, then specific, as Felis (genus) tigris (species).

binomialism (bīnō'mĭālĭzm) n. bis, twice; nomen, name.] The system of binomial nomenclature.

binovular (binôv'ūlăr) a. [L. bini, pair; ovum, egg.] Pert. two ova; dizygotic; appl. twinning.

binuclear (bīnū'klear), binucleate (bīnū'klëat) a. [L. bis, twice; nucleus, small nut.] Having two nuclei.

bioblast (bī'öblāst) n. [Gk. bros, life;
 blastos, bud.] A hypothetical unit, q.v.
 biocatalyst (bī'ökāt'ālīst) n. [Gk. bios, life; katalysis, dissolving.] An

enzyme; a ferment.

biocellate (bīōs'ēlāt) a. [L. bis, twice; ocellus, dim. of oculus, eye.] Having two ocelli.

biocenosis, -biocoenosis, q.v.

biochemistry (bī'ökēm'īstrī) n. [Gk. bios, life; chemeia, transmutation.] The chemistry of living organisms.

biochore (bī'ökōr) n. [Gk. bios, life; choris, separate.] Boundary of a floral or faunal region; climatic boundary of a floral region; a group of similar biotopes.

biochrome (bī'ökrōm) n. [Gk. bios, life; chroma, colour.] Any natural colouring matter of plants and animals; biological pigment.

biocoenosis (bī'ösēnō'sis) n. [Gk. bios, life; koinos, common.] A community of organisms inhabiting a biotope; biocenosis.

biocycle (bi'ösikl) n. [Gk. bios, life; kyklos, place of assembly.] One of the three main divisions of the biosphere: marine, or fresh-water, or terrestrial habitat.

biodemography (bīödēmŏg'răfi) n. [Gk. bios, life; demos, people; graphein, to write.] Science dealing with the integration of ecology and genetics of populations.

biodynamics (bi'ödinam'iks) n. [Gk. bios, life; dynamis, power.] The science of the active vital phenomena of organisms.

bioecology (bi'öekŏl'öji) n. [Gk. bios, life; oikos, household; logos, discourse.] Ecology of plants and

animals.

bioelectric (bī'öēlēk'trīk) a. [Gk. bios, life; elektron, amber.] Appl. currents produced in living organisms.

bioenergetics (bi'öenerjet'iks) n. [Gk. bios, life; energeia, action.] Study of energy transformations in living organisms.

biogen (bī'öjēn), biogene (bīöjēn) n. [Gk. bios, life; genos, descent.]

A hypothetical unit, q.v.; a large living molecule.

biogenesis (bī'ojēn'ësīs) n. [Gk. bios, life; genesis, descent.] The theory of the descent of living matter from living matter—omne vivum e vivo. Opp. abiogenesis.

biogenetic law, - recapitulation

theory, q.v.

biogenous (bīŏj'ënŭs) a. [Gk. bios, life; genos, offspring.] Inhabiting living organisms, as parasites.

biogeny (biŏj'eni) n. [Gk. bios, life; genesis, descent.] The science of the evolution of organisms, comprising ontogeny and phylogeny.

biogeochmeistry (bī'öjē'ökĕm'īstrī)

n. [Gk. bios, life; ge, earth; chemeia,
transmutation.] The study of the
distribution and migration of
chemical elements present in living
organisms and in interaction with
their geographical environment.

biogeography (bī'öjēŏg'răfi) n. [Gk. bios, life; ge, earth; graphein, to write.] The part of biology dealing with the geographical distribution of plants (phytogeography) and animals (zoogeography); chorology.

biological (biölöj'ikal) a. [Gk. bios, life; logos, discourse.] Relating to

the science of life.

biology (biŏl'öji) n. [Gk. bios, life; logos, discourse.] The science of life and living.

bioluminescence (bī'ölūmīnes'ëns, -loo-) n. [Gk. bios, life; L. luminescere, to grow light.] Light-production, as in many groups of animals, and in bacteria and fungi.

biolysis (bīŏl'īsīs) n. [Gk. bios, life; lysis, loosing.] The decomposition of organic matter resulting from activity of living organisms; disintegration of life.

biomass (bī'öniās) n. [Gk. bios, life; massein, to squeeze.] Total weight of organisms per unit area.

biome (biōm) n. [Gk. bios, life.] A community of living organisms; a complex of climax communities of plants and animals in a major region, as tundra, forest, grassland, desert.

biometeorology (bī'ömětëōrŏl'öji) n. [Gk. bios, life; meteorologia, treatise on the heavenly bodics.] The study of the effects of atmospheric conditions upon plants and animals.

biometrics (biömět'riks) n. [Gk. bios, life; metron, measure.] The statistical study of living organisms and their variations; biometry.

bion (bī'ŏn), biont (bī'ŏnt) n. [Gk. bion, living.] An independent living organism; an individual organism.

bionergy (bi'ŏnërji) n. [Gk. bios, life; energeia, action.] Vital force. bionomics (biönŏm'íks) n. [Gk. bios, life; nomos, law.] The study

of organisms in relation to their environment; bionomy; ecology. biophore (bi'öför) n. [Gk. bios, life;

biophore (bi'öfōr) n. [Gk. bios, life; pherein, to carry.] A hypothetical unit, q.v.

biophotogenesis (bī'öfōtöjēn'ēsĭs) n. [Gk. bios, life; phos, light; genesis, origin.] The production and emission of light by plants or by animals; bioluminescence.

biophysics (biöfiz'iks) n. [Gk. bios, life; physis, nature.] Study of biological phenomena interpreted in terms of physical principles; physics as applicable to biology.

biophyte (bī'öfit) n. [Gk. bios, life; phyton, plant.] A plant which gets sustenance from living organisms.

bioplasm (bī'öplăzm) n. [Gk. bios, life; plasma, mould.] Living

matter; protoplasm.

bioplast (bī'öplāst) n. [Gk. bios, life; plastos, formed.] A minute quantity of living protoplasm capable of reproducing itself.

biopsy (bī'öpsī) n. [Gk. bios, life; opsis, sight.] Examination of living organisms, organs, or tissues.

biorgan (bī'ôrgan) n. [Gk. bios, life; organon, instrument.] An organ in the physiological sense, not necessarily a morphological unit.

bios (bi'os) n. [Gk. bios, life.]
Organic life, plant or animal; a complex mixture of vitamins or growth factors; B complex, q.v.

bioseries (bi'ösērīēz) n. [Gk. bios, life; L. series, row.] A succession

of changes of any single heritable character.

biosomes (bī'ösōmz) n. plu. [Gk. bios, life; soma, body.] Structural and functional units in cytoplasm, as chondriosomes, chromidia and plastids.

biosphere (bī'ösfēr) n. [Gk. bios, life; sphaira, globe.] The part of the globe containing living organisms.

biostatics (bi'ostat'iks) n. [Gk. bios, life; statos, stationary.] The science of structure in relation to function of organisms.

biosystematics,—genonomy; taxonomy.

biota (bīō'tă) n. [Gk. bios, life.] The fauna and flora of a region.

biotic (bīŏt'īk) a. [Gk. biotikos, pert. life.] Pert. life; vital.

biotic community,—a community of plants and animals as a whole.

biotic potential,—highest possible rate of population increase, resulting from maximum natality and minimum mortality.

biotin (bī'ötīn) n. [Gk. bios, life.] Vitamin H, a growth substance of yeast, also obtained from liver; or coenzyme R, required by nitrogenfixing bacteria; antiperosis factor; antiavidin; C₁₀H₁₆O₃N₂S.

biotonus (bīot'onus) n. [Gk. bios, life; tonos, tension.] The ratio between assimilation and dissimila-

tion of biogens.

biotope (bi'ötōp) n. [Gk. bios, life; topos, place.] An area in which the main environmental conditions and biotypes adapted to them are uniform; a place where organisms can survive; also, microhabitat.

biotype (bī'ötīp) n. [Gk. bios, life; L. typus, image.] Type of plant or animal; all the individuals of equal genotype.

biovulate (bīōv'ūlăt) a. [L. bis, twice; orum, egg.] Containing two ovules. bipaleolate (bīpā'lēölāt) a. [L. bis, twice; palea, chaff.] Furnished

with two small paleae.

bipalmate (bipăl'māt) a. [L. bis, twice; palma, palm of hand.] Lobed with the lobes again lobed. biparietal (bi'pări'etăl) a. [L. bis, twice; paries, wall.] Connected with the two parietal eminences.

biparous (bĭp'arus) a. [L. bis, twice; parere, to bear.] Having two young at a time; dichotomous, appl. branching.

bipectinate (bīpēk'tīnāt) a. [L. bis, twice; pecten, comb.] Having the two margins furnished with teeth like a comb.

biped (bī'pēd) n. [L. bis, twice; pes, foot.] A two-footed animal.

bipenniform (bipěn'īfôrm) a. [L. bis, twice; penna, feather; forma, shape.] Feather-shaped, with sides of vane of equal size; appl. some muscles.

bipetalous (bīpět'ālŭs) a. [I.. bis, twice; Gk. petalon, leaf.] With two petals.

bipinnaria (bīpĭnā'rīǎ) n. [L. bis, twice; pinna, feather.] An asteroid larva with two bands of cilia.

bipinnate (bīpīn'āt) a. [L. bis, twice; pinna, feather.] Having leaflets growing in pairs on paired stems.

bipinnatifid (bi'pinăt'ifid) a. [L. bis, twice; pinna, feather; findere, to cleave.] With leaves segmented and these segments again divided.

bipinnatipartite (bī'pināt'īpârtīt) a. [L. bis, twice; pinna, feather; partiri, to divide.] Bipinnatifid, but with divisions extending nearly to midrib.

bipinnatisect (bī'pĭnăt'isĕkt) a. [L. bis, twice; pinna, feather; secare, to cut.] Bipinnatifid, but with divisions extending to midrib.

biplicate (bī'plīkāt) a. [L. bis, twice; plicare, to fold.] Having two folds.

bipocillus (bī'pösīl'ūs) n. [L. bis, twice; pocillum, little cup.] A microsclere with curved shaft and cup-shaped expansion at each end.

bipolar (bipō'lăr) a. [L. bis, twice; polus, pole.] Having, located at, or pert. two ends or poles; appl. nerve cells having a process at each end; appl. allied species

occurring towards Arctic and Antarctic regions.

bipolarity (bi'pölăr'îti) n. [L. bis, twice; polus, pole.] The condition of having two polar processes; condition of having two distinct poles, as vegetative and animal poles in an egg.

biradial (bīrā'dīāl) a. [L. bis, twice; radius, ray.] Symmetrical both radially and bilaterally, as some coelenterates; disymmetrical.

biramose (birā'mos), biramous (birā'mus) a. [L. bis, twice; ramus, branch.] Divided into two branches.

birostrate (bīrŏs'trāt) a. [L. bis, twice; rostrum, beak.] Furnished with two beak-like processes.

birth pore,—uterine pore of trematodes and cestodes; birth-opening of redia of trematodes.

biseptate (bīsep'tāt) a. [L. bis, twice; septum, fence.] With two partitions.

biserial (bīsē'rĭāl) a. [L. bis, twice; series, row.] Arranged in two rows or series; biseriate.

biserrate (biser'āt) a. [L. bis, twice; serra, saw.] Having marginal teeth which are themselves notched. bisexual (bisek'sūāl, a. [L. bis, twice; sexus, sex.] Having both male and female reproductive organs; herma-

phrodite; amphisporangiate, q.v. bisporangiate (bi'sporan'jiāt) a. [L. bis, twice; Gk. sporos, seed; anggeion, vessel.] Having both micro- and megasporangia; appl. strobilus consisting of both micro- and megasporophylls.

bistephanic (bī'stĕfān'īk) a. [L. bis, twice; Gk. stephanos, crown.] Joining two points where coronal suture crosses superior temporal ridges.

bistipulate (bīstīp'ūlāt) a. [L. bis, twice; stipula, stem.] Provided with two stipules.

bistrate (bī'strāt) a. [L. bis, twice; stratum, layer.] Having two layers; appl. e.g. indumentum.

bistratose (bīstrāt'ōs) a. [L. bis, twice; stratum, layer.] With cells arranged in two layers.

bisulcate (bīsŭl'kāt) a. [L. bis, twice; sulcus, groove.] Having two grooves.

bitemporal (bītěm'pörăl) a. [L. bis, twice; tempora, temples.] Appl. two temporal bones; a line joining posterior ends of two zygomatic processes.

biternate (bītěr'nāt) a. [L. bis, twice; terni, three by three.] Ternate with each division itself again ternate.

bivalent (bīvā'lēnt, bĭv'alēnt) a. [L. bis, twice; valere, to be strong. Appl. paired homologous chromosomes.

bivalve (bī'vălv) a. [L. bis, twice; valvae, folding doors.] Consisting of two plates or valves, as a mussel shell; or appl. a seed-capsule of similar structure.

biventer cervicis (bīvēn'tër sërvī'sīs) [L. bis, twice; venter, belly; cervix, neck.] The spinalis capitis, or medial part of semispinalis, a muscle of neck, consisting of two fleshy ends with narrow tendinous portion in middle.

biventral (bīvěn'trăl) a. [L. bis, twice; venter, belly.] Appl. muscles of the biventer type; digastric; appl. a lobule of the cerebellum.

bivittate (bīvǐt'āt) a. [L. bis, twice; vitta, band.] With two oil receptacles; with two stripes.

bivium (biv'ium) n. [L. bis, twice; via, way.] Generally the posterior pair of ambulacral areas in certain Echinoidea; the two rays between which the madreporite lies.

bladder (blad'er) n. [A.S. blaedre, A membranous sac filled with air or fluid; a cyst.

bladderworm stage, — cysticercus stage in tape-worms.

blade (blād) n. [A.S. blaed, leaf.] The flat part of leaf of grasses; lamina.

Blandin's glands [P.-F. Blandin, French surgeon]. Anterior lingual glands; glands of Nuhn.

blastaea (blaste'a) n. [Gk. blastos, bud.] A planaea or ciliated planula, a hypothetical stage in evolution.

blastelasma (blast'ělas'ma) n. [Gk. blastos, bud; elasma, plate.] Any

germ layer formed after formation of epiblast and hypoblast.

blastema (blast'ēmā) n. [Gk. blastema, bud.] Formative substance in an egg; primordium of an organ; thallus of a lichen.

blastic (blăs'tik) a. [Gk. blastos, bud.] Pert. or stimulating enlargement by cell-division; opp. trophic.

blastocarpous (blăs'tökâr'pŭs) a. [Gk. blastos, bud; karpos, fruit.] Developing while still surrounded by pericarp.

blastocheme (blas'tökēm) n. [Gk. blastos, bud; ochema, vessel.] A reproductive individual in some Medusae.

blastocholines (blăs'tökō'lēnz) n. plu. [Gk. blastos, bud; cholos, halting.] Various substances, present in sporangia, seeds, and fruits, which prevent premature germination; germination inhibitors.

blastochyle (blăs'tokīl) n. blastos, bud; chylos, juice.] fluid in a blastocoel or segmenta-

tion-cavity.

blastocoel (blăs'tösēl) n. [Gk. blastos, bud; koilos, hollow.] The segmentation-cavity, cavity blastula.

blastocolla (blăs'tökŏl'ă) n. blastos, bud; kolla, glue.] gummy substance coating certain buds.

blastocone (blăs'tökōn) n. [Gk. blastos, bud; konos, cone.] outer larger cell of first circumferential division, in segmentation of certain eggs.

blastocyst (blas'tösist) n. [Gk. blastos, bud; kystis, bladder.] The

germinal vesicle.

blastocyte (blas'tösīt) n. [Gk. blastos, bud; kytos, hollow.] undifferentiated embryonic cell.

blastoderm (blas'töderm) n. blastos, bud; derma, skin.] germinal disc.

blastodermic vesicle, --- hollow sphere of cells, an early stage in development of a fertilised ovum.

- blastodisc (blas'tödisk) n. [Gk. blastos, bud; diskos, disk.] The germinal area of a developing ovum.
- blastogenesis (blas'töjen'esis) n. [Gk. blastos, bud; genesis, descent.] Gemmation or reproduction by budding; transmission of inherited characters by means of germ-plasm only.
- blastogenic (blas'töjen'ik) a. [Gk. blastos, bud; genos, offspring.] Appl. inactive idioplasm unalterable till time and place of activity are reached; arising from changes in germ cells; appl. characteristics of germinal constitution.

blastokinesis (blas'tökinē'sis) n. [Gk. blastos, bud; kinesis, movement.] Movement of embryo in the egg, as in certain insects and cephalo-

pods.

blastomere (blas'tömer) n. blastos, bud; meros, part.] One of the cells formed during primary divisions of an egg; cleavage cell.

blastoneuropore (blas'tönū'röpōr) n. [Gk. blastos, bud; neuron, nerve; passage.] A temporary passage connecting blastopore and neuropore.

blastophore (blas'töför) n. blastos, bud; pherein, to bear.] Embryonic origin of plumule; the reproductive body in Alcyonaria; central part of spermocyte mass which remains unchanged through spermatogenesis in Annelida.

blastophthoria (blas'töfthō'rĭā) n. [Gk. blastos, bud; phthora, corruption.] Any injurious effect on germ

cells or on germ plasm.

blastopore (blas'töpör) n. ſGk. blastos, bud; poros, passage.] Channel leading into archenteron of gastrula.

blastosphere (blăs'tösfēr) n. [Gk. blastos, bud; sphaira, globe.] The blastula; blastodermic vesicle; a hollow ball of cells.

blastospore (blăs'töspōr) n. blastos, bud; sporos, seed.] An attached thallospore developed by budding and itself capable of budding, as of yeast cells.

blastostyle (blas'töstil) n. blastos, bud; stylos, pillar.] In Hydrozoa, a columniform zooid with or without mouth and tentacles, bearing gonophores.

blastozoite (blăs'tözō'īt) n. blastos, bud; zoe, life.] An individual organism produced by

budding.

blastozooid (blăs'tözō'oid) n. [Gk. blastos, bud; zoon, animal; eidos, form.] A larval bud in precocious budding in ascidians.

blastula (blăs'tūlă) n. [L. dim. of Gk. blastos, bud.] A hollow ball of cells, with wall usually one layer thick; blastosphere.

blastulation (blăs'tūlā'shūn) n. [L. blastula, little bud.] Formation of blastulae.

bleeder,-an individual subject to haemophilia, q.v.

- bleeding, of plants, exudation of watery sap from vessels at a cut surface, due to root-pressure.
- blended inheritance,-mixed race or descent; mingling or nonsegregation of parental characteristics.
- blendling (blen'dling) n. [A.S. blandan, to mix.] A racial hybrid.
- blennoid (blen'oid) a. [Gk. blennos, mucus; eidos, form.] Resembling mucus.
- (blĕf'ărăl) blepharal a. blepharon, eyelid.] Pert. eyelids.
- blepharoplast (blěf'ăröplăst) [Gk. blepharis, eyelash; plastos, formed.] A basal granule in relation with a motor cell organ, as the flagellum of Flagellata; blepharoblast.
- blight (blit) n. [A.S. blaecan, to grow pale.] An insect or fungus producing a plant disease; the disease itself.
- blind pit,—a cell-wall pit which is not backed by a complementary pit.
- blind spot,—region of retina devoid of rods and cones and where optic nerve enters; optic disc.

- blister (blis'ter) n. [A.S. blowan, to blow.] A subcutaneous bubble or bladder filled with fluid; a certain plant disease.
- blood (blud) n. [A.S. blod, blood.]
 The fluid circulating in the vascular system of animals, distributing food-material and oxygen and collecting waste products.

blood cells,—cells derived by mitosis from ordinary mesoderm cells; primitive haematoblasts.

blood crystals,—crystals of haemoglobin, haemin, or haematoidin, which form when blood is shaken up with chloroform or ether.

blood dust,—fine droplets of neutral fats present in the blood stream; haemokonia.

blood gills,—delicate blood-filled sacs functioning in uptake of salts, in certain insects.

blood groups,—types of blood depending on presence or absence of two agglutinogens (A and B) in the red corpuscles and two agglutinins (a or anti-A, and β or anti-B) in serum or plasma: A cells agglutinate with B type serum, B with A type, AB with A and B type, and O cells not agglutinating with A and B types; cf. universal donor, universal recipient.

blood islands, — isolated reddish patches in mesoderm, in which primitive erythroblasts are found enclosed in a mesodermal syncytium; blood anlage, haemangioblast.

blood platelets,—colourless bodies about one-third the size of red corpuscles, and formed from megakaryocytes, and agglutinating in shed blood; thrombocytes; thromboplastids.

blood plates,—minute amoeboid protoplasmic bodies found in blood. blood serum,—fluid or plasma left after removal of corpuscles and

fibrin.

blood vessel,—any vessel or space in which blood circulates; strictly used only in regard to special vessels with well-defined walls. bloom,—a layer of wax particles on external surface of certain fruits, as grapes, peaches; blossom or flower

blubber (blub'ër) n. [M.E. blober, a bubble.] Fat of whales, seals, etc., lying between outer skin and muscle layer.

blue timber,—a wood disease produced by fungus, causing a bluish discoloration.

body blight,—fungal disease of trees.
body cavity,—coelom or space in which viscera lie, mesodermal in origin, and schizocoelic or enterocoelic in development; considered primarily, the generative cavity.

body cell,—a somatic cell as distinct from a germ cell; an antheridial cell.

body stalk,—a band of mesoderm connecting caudal end of embryo with chorion.

Bojanus, organ of [L. H. Bojanus, Alsatian zoologist]. Excretory organ in lamellibranchs.

bone (bon) n. [A.S. ban, bone.]
Connective tissue in which the ground-substance contains salts of lime.

bone-beds,—deposits formed largely by remains of bones of fishes and reptiles, as Liassic bone-beds.

bones of Bertin [E. J. Bertin, French anatomist]. Thin anterior coverings of sphenoidal sinuses.

bonitation (bonita'shun) n. [L. bonitas, goodness.] The evaluation of the numerical distribution of a species in a particular locality or season, in relation to agricultural, veterinary, or medical implications.

book gill,—a gill composed of delicate leaf-like lamellae placed one over the other like leaves of a book, as seen in Limulus.

book lung,—a gill similar to a book gill, but modified for air-breathing, and open to exterior only by a small slit, as in scorpions.

booted (boot'ëd) a. [O.F. boute, boot.] Equipped with raised horny plates of skin, as feet of some birds. bordered pit,—a form of pit, developed on walls of tracheids and wood-vessels, with overarching border of secondary cell-wall.

boreal (bō'rëăl) a. [L. boreas, north wind.] Appl. or pert. northern biogeographical region; holarctic except Sonoran, or restricted to nearctic; pert. post-glacial age with continental type of climate.

bosselated (bŏs'ëlātēd) a. [M.E. bosse, knob.] Covered with knobs. bosset (bŏs'ēt) n. [M.E. bosse,

knob.] The beginning of horn formation in deer in the first year. bostryx (bŏs'trĭks) n. [Gk. bostry-

chos, curl.] A helicoid cyme, cymose inflorescence with blooms on only one side of axis.

Botallo's duct [L. Botallo, Italian surgeon]. Ductus arteriosus, a small blood vessel representing sixth gill arch and connecting pulmonary with systemic arch.

botany (bŏt'ánĭ) n. [Gk. botane, pasture.] The branch of biology dealing with plants; phytology.

bothridium (bothrid'ium) n. [Gk. bothros, trench; idion, dim.] A muscular cup-shaped outgrowth from scolex of tape-worms; a phyllidium.

bothrium (both'rium) n. [Gk. bothros, trench.] A sucker; a sucking groove in scolex of tape-

worms.

botryoidal (hötrioid'āl) a. [Gk. botrys, bunch of grapes; eidos, form.] In the form of a bunch of grapes; appl. tissue of branched canals surrounding enteric canal in leeches.

botryose (bŏt'rīōs) a. [Gk. botrys, bunch of grapes.] Racemose; botryoidal.

botuliform (bŏt'ūlifôrm) a. [L. botulus, sausage; forma, form.]
Sausage-shaped; allantoid.

bouillon (booyong) n. [F. bouillon, broth.] An infusion or broth, containing watery extract of meat, also peptone, for the cultivation of bacteria.

bouquet (booka', book'a) n. [F.

bouquet, nosegay. Arrangement of chromosomes in loops with their ends near one side of nuclear wall during zygotene and pachytene in some organisms.

bouton (bootông) n. [F. bouton, bud.] Terminal bulb of arborisation of an axon; labellum, in

Hymenoptera.

Bowman's capsule [Sir W. Bowman, English histologist]. The vesicle of a renal tubule; capsula glomeruli.

Bowman's glands,—serous glands in corium of olfactory mucous membrane.

Bowman's membrane,—anterior 'elastic' membrane of cornea.

braccate (brāk'āt) a. [L. braccae, breeches.] Having additional feathers on legs or feet, appl. birds.

brachelytrous (brăkěl'ítrůs) a. [Gk. brachys, short; elytron, sheath.] Having short wing-covers.

brachia (brāk'īā) n. plu. [L. brachium, arm.] The arms; two spirally coiled structures, one at each side of mouth, in Brachiopoda; cerebellar peduncles; white lateral bands of colliculi of corpora quadrigemina. Sing. brachium.

brachial (brāk'iăl) a. [L. brachium, arm.] Pert. arm; arm-like.

brachialis (brākiā'līs) n. [L. brachium, arm.] A flexor muscle of the forearm, from lower half of front of humerus to coronoid process of ulna; brachialis anticus.

brachiate (brā'kiāt) a. [L. brachium, arm.] Branched; having opposite paired branches on alternate sides.

brachidia (brăkid'iă) n. plu. [Gk. brachion, arm; idion, dim.] Calcareous skeleton supporting brachia in certain Brachiopoda.

brachiferous (brākif'ērūs), brachigerous (brākij'erūs) a. [L. brachium, arm; ferre, gerere, to carry.] Branched.

brachiocephalic (brāk'iökĕfāl'īk, -sĕf-) a. [L. brachium, arm; Gk. kephale, head.] Pert. arm and head; appl. artery.

62

brachiocubital (bråk'iökū'bītǎl) a. [L. brachium, arm; cubitum, forearm.] Pert. arm and forearm.

brachiolaria (brākīölā'rīā) n. [L. brachiolum, small arm.] A larval stage in metamorphosis of some starfishes.

brachiole (bră'kiōl) n. [L. brachiolum, small arm.] A pinnule-like structure on ambulacral margin in Blastoidea.

brachiorachidian (bråk'iöråkid'iän)
a. [Gk. brachion, arm; rhachis, spine.] Pert. arm and spine.

brachioradialis (brāk'iörādiā'līs) n.
[L. brachium, arm; radius, ray.]
The supinator longus muscle of forearm.

brachium (brāk'iŭm) n. [L. brachium, arm.] Arm or branching structure; upper limb of vertebrates; a bundle of fibres connecting cerebellum to cerebrum or to pons.

brachyblast,—brachyplast, q.v.

brachycephalic (brǎk'îkĕfāl'îk, -sĕf-)
a. [Gk. brachys, short; kephale, head.] Short-headed; with cephalic index of over eighty; cf. dolichocephalic.

brachycerous (brākis'ērūs) a. [Gk. brachys, short; keras, horn.] Shorthorned; with short antennae.

brachycnemic (brāk'īknē'mīk) a. [Gk. brachys, short; kneme, tibia.] Appl. arrangement of mesenteries of Zoantharia where the sixth protocneme is imperfect.

brachydactyly (brăk'îdăk'tili) n.
[Gk. brachys, short; daktylos, digit.]
Brachydactylous condition, viz.
having digits abnormally short.

brachydont (brāk'idönt) a. [Gk. brachys, short; odous, tooth.] Appl. molar teeth with low crowns; also brachyodont.

brachymeiosis (brāk'imīō'sīs) n. [Gk. brachys, short; meion, smaller.] A third karyokinetic or second reduction division, as in asci; meiosis involving only one division.

brachyodont,—brachydont, q.v. brachyourous,—brachyural, q.v.

brachyplast (brak'iplast) n. [Gk. brachys, short; plastos, formed.]
A short branch or spur bearing leaf tufts, occurring with normal branches on the same plant.

brachypleural (brăk'iploo'răl) a. [Gk. brachys, short; pleuron, side.] With short pleura or side plates.

brachypodous (brăkip'ödus) a. [Gk. brachys, short; pous, foot.] With short legs, or stalk.

brachypterous (brākip'tërūs) a. [Gk. brachys, short; pteron, wing.] With short wings.

brachysclereid (bråk'isklē'rēĭd) n. [Gk. brachys, short; skleros, hard; eidos, form.] A stone cell.

brachysm (bräk'ism) n. [Gk. brachys, short.] Dwarfism in plants caused by shortening of internodes.

brachystomatous (brāk'īstŏm'ātūs)

a. [Gk. brachys, short; stoma, mouth.] With short proboscis; appl. certain insects.

brachytic (brākīt'īk) a. [Gk. brachytes, shortness.] Dwarfish, appl. plants; exhibiting or pert. brachysm.

brachyural (brăk'iū'răl) a. [Gk. brachys, short; oura, tail.] Having short abdomen usually tucked in below thorax, appl. certain crabs.

brachyuric (brāk'īū'rīk) a. [Gk. brachys, short; oura, tail.] Short-tailed.

bract (bråkt) n. [L. bractea, thin plate of metal.] A floral leaf; a modified leaf in whose axil a flower arises; a hydrophyllium in Siphonophora; distal exite of sixth appendage of Apus.

bract scales,—small scales developed directly on axis of cones; cf. ovu-

liferous scales.

bracteal (brăk'tēăl) a. [L. bractea, thin metal plate.] Like a bract; bracteiform.

bracteate (brāk'tēāt) a. [L. bractea, thin metal plate.] Having bracts.

bracteolate (brăk'tëölāt) a. [L. bractea, thin metal plate.] Appl. flowers with bracteoles.

bracteole (brāk'tēōl) n. [L. bractea, thin metal plate.] Secondary bract at base of flower; bractlet.

bracteose (brăk'tëōs) a. [L. bractea, thin metal plate.] With many bracts.

bractlet,-bracteole.

bradyauxesis (brad'íôksē'sīs) n. [Gk. bradys, slow; auxesis, growth.] Relatively slow growth; growth of a part at a slower rate than that of the whole; opp. tachyauxesis.

bradygenesis (brăd'ijĕn'ësis) n. [Gk. bradys, slow; genesis, descent.] Retarded development, in phylogeny; opp. tachygenesis.

bradytelic (brădĭtěl'ík) a. [Gk. bradys, slow; telos, fulfilment.] Evolving at a rate slower than the standard rate; opp. tachytelic; cf. horotelic.

brain (brān) n. [O.E. braegen, brain.]
Centre of nervous system; mass of
nervous matter in vertebrates at
anterior end of spinal cord, lying in
cranium; in invertebrates, supraoesophageal or suprapharyngeal
ganglia.

brain sand,—granular bodies of calcium and ammonium and magnesium phosphates, occurring in pineal gland and pia mater; corpora amylacea; acervulus cerebri.

branchia (brăng'kiă) n., branchiae (brăng'kiē) plu. [Gk. brangchia, gills.] Gills.

branchiac (brăng'kiăk), branchial (brăng'kiăl) a. [Gk. brangchia, gills.] Pert. gills.

branchial arch,—one of the bony or cartilaginous arches on side of the pharynx posterior to hyoid arch, and supporting gill bars.

branchial grooves,—outer pharyngeal grooves or visceral clefts, q.v. branchiate (brăng'kiāt) a. [Gk. brangchia, gills.] Having gills.

branchicolous (brăngkik'ölüs) a. [Gk. brangchia, gills; L. colere, to inhabit.] Parasitic on fish gills; appl. certain crustaceans.

branchiform (brang'kifôrm) a. [Gk.

brangchia, gills; L. forma, shape.] Gill-like.

branchihyal (brăng'kĭhī'āl). [Gk. brangchia, gills; hyoeides, Y-shaped.]
An element of a branchial arch.

branchiocardiac (brăng'klökâr'dlăk)

a. [Gk. brangchia, gills; kardia, heart.] Pert. gills and heart; appl. vessel given off ventrally from ascidian heart; appl. vessels conveying blood from gills to pericardial sinus in certain crustaceans.

branchiomere (brăng'kiömēr) n. [Gk. brangchia, gills; meros, part.] A branchial segment.

branchiopallial (brang'kiöpäl'iăl) a. [Gk. brangchia, gills; L. pallium, mantle.] Pert. gill and mantle of molluses.

branchiostegal (brăng'kiŏs'tëgăl) a. [Gk. brangchia, gills; stege, roof.] With or pert. a gill cover; appl. membrane, rays.

branchiostege (brăng'kĭöstēj') n. [Gk. brangchia, gills; stege, roof.] The branchiostegal membrane.

branchiostegite (brăng'kiŏs'tējīt) n. [Gk. brangchia, gills; stege, roof.] Expanded lateral portion of carapace forming gill cover in certain Crustacea.

branchireme (brăng'kĭrēm) n. [L. branchiae, gills; remus, oar.] A branchiate limb; locomotory and respiratory limb of Branchiopoda.

brand (brand) n. [A.S. beornan, to burn.] A burnt appearance on leaves, caused by fungi.

bregma (breg'ma) n. [Gk. bregma, fore-part of head.] That part of skull where frontals and parietals meet; intersection of sagittal and coronal sutures.

brephic (bref'ik) a. [Gk. brephikos, childish.] Appl. a larval phase preceding that of adult form; neanic.

brevicaudate (brev'ikô'dāt) a. [L. brevis, short; cauda, tail.] With a short tail.

brevifoliate (brev'ifō'liāt) a. [L. brevis, short; folium, leaf.] Having short leaves.

- brevilingual (brev'îlĭng'gwăl) a. [L. brevis, short; lingua, tongue.]
 With short tongue.
- breviped (brev'iped) a. [L. brevis, short; pes, foot.] Having short legs; appl. certain birds.
- brevipennate (brev'ipen'āt) a. [L. brevis, short; penna, feather.] With short wings.
- brevirostrate (brev'iros'trāt) a. [L. brevis, short; rostrum, beak.] With short beak or rostrum.
- **brevissimus oculi,** obliquus inferior, shortest muscle of eye.
- bridge corpuscle,—desmosome, q.v. Broca's area [P. P. Broca, French surgeon]. Parolfactory area of brain.
- Broca's gyrus,—left inferior frontal gyrus, speech centre in cerebral cortex.
- brochidodrome (brŏkĭd'ödrōm) a. [Gk. brochos, loop; dromein, to run.] Appl. veins in leaves when they form loops within the blade.
- brochonema (brökönē'mā) n. [Gk. brochos, loop; nema, thread.] The spireme in loops to the number of chromosome pairs to be formed.
- bromatium (brömā'shĭum) n. [Gk. broma, food.] A swelling on a fungus cultivated by ants, and serving as food.
- bronchi (brong'ki) n. plu. [Gk. brongchos, windpipe.] Tubes connecting trachea with lungs. Sing. bronchus.
- bronchia (bröng'kiă) n. plu. [Gk. brongchos, windpipe.] The subdivisions or branches of each bronchus.
- bronchial (brŏng'kĭăl) a. [Gk. brong-chos, windpipe.] Pert. bronchi.
- bronchiole (brong'kiōl) n. [Gk. brongchos, windpipe.] A small terminal branch of bronchi.
- bronchopulmonary (brong'köpŭl'mönārī) a. [Gk. brongchos, windpipe; L. pulmo, lung.] Pert.
 bronchi and lungs.
- bronchotracheal (brong'kötră'kēal)
 a. [Gk. brongchos, windpipe; L. trachea, trachea.] Pert. bronchi and trechea.

- bronchovesicular (brong'kövesik'ūlar) a. [Gk. brongchos, windpipe; L. vesicula, little sac.] Pert. bronchial tubes and lung cells.
- bronchus,-sing. of bronchi.
- brood bud,—a spore of certain types of sporangia; a soredium; a bulbil.
- brood cells,—gonidia, q.v.
- brood pouch,—a sac-like cavity in which eggs or embryos are placed; a space formed by overlapping plates attached to bases of thoracic limbs in certain Crustacea.
- brown body,—a brown, rounded mass of compacted degenerate organs in some polyzoa; nephrocyte in ascidians.
- brown funnels,—a single pair of organs on dorsal aspect of posterior end of pharynx, in Amphioxus; atrio-coelomic funnels; brown canals.
- Brownian movements [R. Brown, Scottish botanist]. The passive vibratory movements of fine granules when suspended in a fluid.
- Bruch's membrane [C. W. L. Bruch, German anatomist]. The basal membrane, inner layer of choroid; lamina basalis.
- Brunner's glands [J. C. Brunner, Swiss anatomist]. Small tubuloracemose glands containing a proteolytic enzyme, in submucous coat of small intestine; duodenal glands.
- bryology (brīŏl'öjĭ) n. [Gk. bryon, moss; logos, discourse.] The science dealing with mosses, also with liverworts; muscology.
- bryophyte (brī'öfīt) n. [Gk. bryon, moss; phyton, plant.] Any of the mosses, or liverworts.
- bryozoon (brī'özō'ŏn) n. [Gk. bryon, moss; zoon, animal.] A polyzoon, so named from moss-like appearance.
- **B-substance**,—intermedin, q.v.
- buccae (bŭk'ē) n. plu. [L. bucca, cheek.] The cheeks.
- buccal (bŭk'āl) a. [L. bucca, cheek.] Pert. the cheek or mouth.
- buccinator (būk'sīnā'tŏr) n. [L. buccinator, trumpeter.] A broad thin muscle of the cheek.

buccolabial (bŭk'ölā'bĭāl) a. [L. bucca, cheek; labium, lip.] Pert. mouth cavity and lips.

buccolingual (bŭk'öling'gwăl) a. [L. bucca, cheek; lingua, tongue.]

Pert. cheeks and tongue.

bucconasal (būk'onā'zāl) a. [L. bucca, cheek; nasus, nose.] Pert. cheek and nose; appl. membrane closing posterior end of olfactory

buccopharyngeal (bǔk'öfārĭn'jēāl) a. [L. bucca, cheek; Gk. pharyngx, throat.] Pert. cheeks and pharynx; appl. membrane and fascia.

bud (bud) n. [M.E. budde, bud.] A rudimentary shoot, or flower; a

gemma.

budding (bud'ing) n. [M.E. budde, bud.] A method of reproduction seen in many primitive animals.

buffer (but'er) n. [O.F. buffe, blow.] Appl. salt solution which minimises changes in pH when an acid or alkali is added; appl. genes controlling the action of an allelomorph, i.e. polygenes.

bufotoxins (bū'fotŏk'sĭnz) n. plu. [L. bufo, toad; Gk. toxikon, poison.] Toad venom, as bufotoxin, C₈₄H₄₆O₁₀, and bufonin,

C34H54O2.

bulb (bulb) n. [L. bulbus, globular root.] A specialised underground bud with thick fleshy leaves; a part resembling a bulb; a bulb-like dilatation; basal part of intromittent organ in spiders; the medulla oblongata.

bulbar (bul'bar) a. [L. bulbus, globular root.] Pert. bulb or bulb-like part; pert. medulla oblon-

gata.

bulbiferous (bulbif'ërus) a. [L. bulbus, bulb; ferre, to carry.]

Bulb-bearing.

bulbil (būl'bīl) n. [L. bulbus, bulb.]
A fleshy axillary bud which may
fall and produce a new plant, as in
some lilies; aerial bulb; any
small bulb-shaped structure or
dilatation.

bulbocavernosus (bul'bökåverno'sus) n. [L. bulbus, bulb; cavernosus, cavernous.] A muscle of perinaeum, ejaculator urinae in the male; sphincter of vagina.

bulbonuclear (būl'bonū'klēār) a. [L. bulbus, bulb; nucleus, kernel.]
Pert. medulla oblongata and nuclei

of cranial nerves.

bulbo-urethral (bul'boūrē'thrăl) a. [L. bulbus, bulb; Gk. ourethra, urethra.] Appl. two racemose glands, Cowper's or Méry's glands, opening into bulb of male urethra; also appl. the greater vestibular glands, Bartholin's glands, in the female.

bulbous (būl'būs) a. [L. bulbus, bulb.] Like a bulb; developing from a bulb; having bulbs.

bulbus (būl'būs) n. [I.. bulbus, bulb.]
A bulb; swollen base of stipe in agaries; the knob-like part found in connection with various nerves; a dilatation of base of aorta.

bulla (bool'a) n. [L. bulla, bubble.]

Appl. rounded prominence formed
by bones of ear, tympanic bulla;

appl. prominence of middle eth-

moidal air cells.

bullate (bool'āt) a. [L. bulla, bubble.]
Blistered-like; puckered like a

savoy-cabbage leaf.

bulliform (bool'ifôrm) a. [L. bulla, bubble; forma, shape.] Bubble-shaped; appl. thin-walled cells which cause rolling, folding, or opening of leaves by turgor changes.

bundle-sheath,—a layer of large parenchymatous cells surrounding

vascular tissue of leaf-vein.

bunodont (bū'nödŏnt) a. [Gk. bounos, mound; odous, tooth.] Having molar teeth with low conical cusps.

bunoid (bū'noid) a. [Gk. bounos, mound; eidos, form.] Appl. cusps of cheek-teeth, low and conical.

bunolophodont (bū'nölŏf'ödŏnt) a. [Gk. bounos, mound; lophos, crest; odous, tooth.] Between bunodont and lophodont in structure, appl. cheek-teeth.

- bunoselenodont (bū'nösēlē'nodŏnt)

 a. [Gk. bounos, mound; selene,
 moon; odous, tooth.] Having
 internal cusps bunoid, external
 selenoid; appl. cheek-teeth.
- bursa (bur'sa) n. [L. bursa, purse.] A sac-like cavity; a sac with viscid fluid to prevent friction at joints.

bursa copulatrix,—a genital pouch of various animals.

bursa entiana,—the short duodenum in Chondropterygii.

bursa Fabricii,—a sac opening into dorsal part of posterior region of cloaca in birds, and usually degenerating during adolescence.

bursicule (bŭr'sĭkūl) n. [L. dim. of bursa, pursc.] A small sac.

buttress-roots,—branch roots given off above ground, arching away from stem before entering soil, forming additional props.

butyrinase (bū'tirĭnās) n. [L. butyrum, butter.] An enzyme occurring in blood serum.

byssal (bis'ăl) a. [Gk. byssos, fine flax.] Pert. the byssus.

byssogenous (bisŏj'enŭs) a. [Gk. byssos, fine flax; genos, birth.] Byssus-forming; appl. glands.

byssoid (bis'oid) a. [Gk. byssos, fine flax; eidos, shape.] Resembling a byssus; formed of fine threads; byssaceous.

byssus (bīs'ūs) n. [Gk. byssos, fine flax.] The tuft of strong filaments secreted by a gland of certain bivalve molluscs, by which they

become attached; the stalk of certain

fungi.

C

- cacogenesis (kăk'öjēn'ësĭs) n. [Gk. kakos, bad; genesis, descent.] Inability to hybridise; kakogenesis.
- cacogenic (kāk'öjēn'ik) a. [Gk. kakos, bad; genos, birth.] Dysgenic, q.v.
- cadophore (kad'öför) n. [Gk. kados, cask; pherein, to bear.] A dorsal

bud-bearing outgrowth in certain tunicates.

caducibranchiate (kădū'sībrăng'kīāt) a. [L. caducus, falling; branchiae, gills.] With temporary gills.

caducous (kădū'kŭs) a. [L. caducus, falling.] Pert. parts that fall off early, e.g. calyx, stipules; fugacious; cf. deciduous.

caecal (sē'kāl) a. [L. caecus, blind.] Ending without outlet; appl. stomach with cardiac part prolonged into blind sac; pert. caecum.

caecum (sē'kŭm) n. [L. caecus, blind.] A blind diverticulum or pouch from some part of alimentary canal.

caecum cupulare,—the closed apical end of the cochlear canal.

Caenogaea (sẽ'nöjē'a) n. [Gk. kainos, recent; gaia, earth.] A zoogeographical region which includes the Nearctic, Palearctic, and Oriental regions; cf. Eogaea; also Cainogea, Kainogaea.

caenogenesis (sē'nojēn'ēsīs) n. [Gk. kainos, recent; genesis, origin.]
The non-phylogenetic processes in development of an individual; development of transitory adaptations in early stages of an individual.

caenogenetic (sē'nöjēnět'ĭk) a. [Gk. kainos, recent; genesis, origin.]
Of recent origin.

Caenozoic (sēnözō'ĭk) a. [Gk. kainos, recent; zoe, life.] Pert. age of mammals, geological era from Mesozoic to recent times; Tertiary and Quaternary periods; also Cainozoic, Cenozoic, Kainozoic.

caespitose (sĕs'pĭtōs) a. [L. caespes, turf.] Pert. turf; having low, closely matted stems; growing densely in tufts; caespitulose, cespitose.

caisson (kā'sŏn) n. [F. caisson, coffer.] Box-like arrangement of longitudinal muscle fibres in Lumbricidae.

calamistrum (kål'amis'trum) n. [L. calamister, curling-iron.] A comblike structure on metatarsus of certain spiders.

- calamus (kāl'āmūs) n. [L. calamus, reed.] A hollow reed-like stem without nodes; the quill of a feather; calamus scriptorius, the tip of posterior part of floor of fourth ventricle.
- calcaneus (kālkā'nēŭs) n. [L. calx, heel.] The heel; large bone or os calcis of tarsus which forms heel; calcaneum; process on metatarsus of birds.
- calcar (kăl'kâr) n. [L. calcar, spur.]
 A hollow prolongation or tube at base of sepal or petal; spur-like process on leg or wing of birds; tibial spine in insects; process of calcaneus which supports web between leg and tail in bats; prehallux of frog; internal bony plate strengthening neck of femur; calcar avis, eminence in posterior part of lateral ventricle.

calcarate (kăl'kărāt) a. [L. calcar, spur.] Spurred; appl. petal, corolla.

- calcareous (kălkā'rēŭs) a. [L. calcareous, limy.] Limy; growing on soil derived from decomposition of calcareous rocks; pert. limestone.
- calcariform (kălkăr'ifôrm) a. [L. calcar, spur; forma, shape.] Spurlike.
 calcarine (kăl'kărin) a. [L. calcar, spur.] Pert. calcar avis; appl.

fissure extending to hippocampal gyrus, on medial surface of cerebral

hemisphere.

calcicole (kāl'sīkōl) n. [L. calx, lime; colere, to dwell.] A plant which thrives in soils rich in calcium salts; calcipete, calciphile, calciphyte, gypsophyte. a. Calcicolous.

calciferol,—vitamin D₂, occurring in fish liver oils, egg yolk, milk, etc., and conserving body calcium and phosphorus; antirhachitic vitamin; isomeric with ergosterol from which it is formed by a series of photochemical reactions; C₂₈H₄₄O.

calciferous (kālsīf'ērūs), calcigerous (kālsīj'ērūs) a. [L. calx, lime; ferre, gerere, to carry.] Containing

or producing lime salts.

calcification (kālsifikā'shūn) n. [L. calx, lime; facere, to make.] The

- deposition of lime salts in tissue; the process of accumulation of lime salts in soil development.
- calcifuge (kāl'sifūj) n. [L. calx, lime; fugere, to flee.] A plant which thrives only in soils poor in calcium carbonate; calciphobe.

calcipete (kăl'sĭpēt) n. [L. calx, lime; petere, to go towards.] A calcicole, g.v.; a calciphil plant.

calciphile,—calciphyte.

calciphobe, -- calcifuge.

- calciphyte (kăl'sifit) n. [L. calx, lime; Gk. phyton, plant.] A plant which thrives only on calcareous soils; calcicole, calcipete, calciphile, gypsophyte.
- calcivorous (kălsīv'örūs) a. [L. calx, lime; vorare, to devour.] Appl. plants which live on limestone.
- calcospherites (kăl'kösfē'rīts) n. plu. [L. calx, lime; sphaera, globe.] Concentrically laminated granules of calcium carbonate in Malpighian tubes of some insects, in cells associated with fat-body in certain larval Diptera.

calicle,—calycle, q.v.

- calines (kălēnz) n. plu. [Gk. kalein, to summon.] Plant hormones influencing growth of specific parts, as of root, stem, or leaf.
- callosal (kălō'săl) a. [L. callosus, hard.] Pert. corpus callosum.
- callose (kal'ōs) n. [L. callum, hard skin.] An occasional carbohydrate or periodic component of plant cell walls, as on sieve-plates. a. Having callosities.
- callosity (kălŏs'iti) n. [L. callositas, hardness.] Hardened and thickened area on skin, or on bark.
- callow (kăl'ō) n. [A.S. calu, bald.]
 A newly hatched worker ant. a.
 Unfledged.
- callus (kāl'ūs) n. [L. callum, hard skin.] Tissue that forms over cut or damaged plant surface; deposit of callose on sieve-plates; small hard outgrowth at base of spikelet or of floret, in some grasses; a growth of shell-like material within umbilicus of shell; a mesonotal swelling in some insects; callosity.

- caloricity (kălöris'iti) n. [L. calere, to be warm.] In animals, the power of developing and maintaining a certain degree of heat.
- calorie (kăl'öri) n. [L. calere, to be warm.] Amount of heat required to raise temperature of one gramme of water one degree centigrade (small calorie); one large calorie equals one thousand small calories.

calorigenic (kăl'örijen'ik) a. [L. calor, heat; genere, to beget.]
Promoting oxygen consumption and

heat production.

- calotte (kălŏt') n. [F. calotte, skull-cap.] An outer cell group or polar cap in Dicyemidae, for adhesion to kidney of Cephalopoda; a retractile disc with sensory cilia, in larval Bryozoa.
- caltrop (kal'trop) n. [A.S. coltraeppe, thistle.] A sponge spicule with four rays so disposed that any three being on the ground the fourth projects vertically upwards; also calthrop.
- calvarium (kălvā'rīŭm) n. [L. calvus, bald.] The dome of the skull.
- calx (kălks) n. [L. calx, lime, heel.] Lime; calcaneus, q.v.
- calycanthemy (kål'ikån'thëmi) n. [Gk. kalyx, calyx; anthemon, flower.] Abnormal development of parts of calyx into petals.

calyces,—plu. of calyx.

- calyciflorous (kål'isiflö'rūs) a. [L. calyx, calyx; flos, flower.] Appl. flowers in which stamens and petals are adnate to the calyx.
- calyciform (kălis'ifôrm) a. [L. calyx, calyx; forma, shape.] Calyx-like in shape.
- calycine (kăl'īsīn) a. [L. calyx, calyx.]

 Pert. a calyx; cup-like.
- calycle (kål'ikl) n. [L. calyculus, little calyx.] An epicalyx; a cupshaped cavity in a coral; a theca in a hydroid; calyculus; also calicle.
- calyculus (kălīk'ūlūs) n. [L. calyculus, little calyx.] Cup-shaped or bud-shaped structure; calycle.
- calyculus gustatorius,—a taste-bud

- or taste-bulb, an ovoid buccal sense organ composed of gustatory cells supported and surrounded by sustentacular cells.
- calyculus ophthalmicus, optic cup, formed by invagination of the optic bulb and developing into the retina.

calymma,—kalymma, q.v.

- calypter (kălip'tër) n. [Gk. kalyptos, hidden.] Antitegula or modified alula covering haltere in certain Diptera; calyptron.
- calyptoblastic (kālīp'töblās'tīk) a. [Gk. kalyptos, hidden; blastos, bud.] Pert. hydroids in which gonophore is enclosed in a gonotheca.
- calyptobranchiate (kălĭp'töbrăng'kīāt) a. [Gk. kalyptos, hidden; brangchia, gills.] With gills not visible from exterior.
- calyptopsis (kăliptop'sis) n. [Gk. kalyptos, hidden; opsis, sight.]
 A larva with short-stalked eyes, as of some arthropods.
- calyptra (kălīp'tră) n. [Gk. kalyptra, covering.] Tissue enclosing developing sporogonium in liverworts; remains of archegonium which surround apex of capsule in mosses; neck of archegonium in prothallus of some pteridophytes; root-cap; cf. calyptrogen.
- calyptrate (kālip'trāt) a. [Gk. kalyptra, covering.] Appl. caducous calyx separating from its lower portion or from thalamus; operculate; appl. Diptera with halteres hidden by squamae.
- calyptrogen (kălip'tröjen) n. [Gk. kalyptra, covering; gennaein, to produce.] The special layer of cells lying at apex of growing root and giving origin to root-cap.

calyptron (kălip'trön) n. [Gk. kalyptra, covering.] The squama of Calypterae; calypter.

calyx (kål'iks) n. [Gk. kalyx, calyx.]
The outer whorl of floral leaves;
cup-like portion of pelvis of kidney;
theca of certain hydroids; cuplike body of crinoids; cup or
head of pedunculate bodies in
insects.

- cambial (kăm'bĭāl) a. [L. cambium, change.] Pert. cambium.
- cambiform (kăm'bĭfôrm) a. [L. cambium, change; forma, shape.] Similar to cambium cells.
- cambiogenetic (kăm'biöjénět'ík) a. [L. cambium, change; Gk. genesis, origin.] Appl. cells which produce cambium.
- cambium (kăm'biŭm) n. [L. cam-bium, change.] The tissue from which secondary growth arises in stems and roots.
- Cambrian (kăm'briăn) a. [L. Cambria, Wales.] Pert. earliest period, or system of rocks, of Palaeozoic era.
- cameration (kămërā'shun) n. [L. cameratio, vaulting.] Division into a large number of separate chambers.
- camerostome (kăm'ërostōm') n. [L. camera, chamber; Gk. stoma, mouth.] Hollowed-out under surface of 'hood' of certain Trogulidae.
- campaniform (kāmpān'īfôrm) a. [L.L. campana, bell; forma, shape.] Bell- or dome-shaped; appl. sensilla.
- campanula Halleri [Dim. of L.L. campana, bell; A. von Haller, Swiss anatomist]. Expansion of falciform process at lens in many fishes.
- campanulate (kămpăn'ūlāt) a. [Dim. of L.L. campana, bell.] Bellshaped; appl. corolla.
- campodeiform (kămpō'dēifôrm) a. [Gk. kampe, caterpillar; eidos, form; L. forma, shape.] Appl. larva resembling a Campodea; thysanuriform.
- camptodrome (kămp'tödrōm) a. [Gk. kamptos, flexible; dromos, course.] Pert. leaf venation in which secondary veins bend forward and anastomose before reaching margin.
- camptotrichia (kamp'tötrikya) n. plu.
 [Gk. kamptos, flexible; thrix, hair.]
 Jointed dermal fin-rays in certain
 primitive fishes.
- campylospermous (kămpilöspěr'mus) a. [Gk. kampylos, curved;

- sperma, seed.] Appl. seeds with groove along inner face.
- campylotropous (kăm'pilot'röpūs) a. [Gk. kampylos, curved; trope, turning.] Pert. ovules in which nucellus and embryo-sac are bent so that micropyle points almost back to placenta.
- canalicular (kănălīk'ūlăr) a. [L. canaliculus, small channel.] Pert. canals, or canaliculi.
- canalicular apparatus,—the Golgi bodies, regarded as a system of canals.
- canaliculus (kănălĭk'ūlūs) n. [L. canaliculus, small channel.] One of the small canals containing cell-processes of bone-corpuscles and connecting lacunae in Haversian system; small channel for passage of nerves through various bones.
- canaliform (kănăl'ĭfôrm) a. [L. canalis, canal; forma, shape.]
 Canal-like.
- cancellous (kăn'sëlūs) a. [L. cancellosus, chambered.] Consisting of slender fibres and lamellae, which join to form a reticular structure; cancellated; appl. inner, more spongy, portion of bony tissue; appl. anterior portion of cuttle-bone.
- cancrisocial (kăng'krīsō'shāl) a. [L. cancer, crab; socius, ally.] Appl. commensals with crabs.
- canine (kănîn', kā'nîn) n. [L. canis, dog.] The tooth next to incisors.
 a. Pert. canine tooth, or to a fossa and eminence on anterior surface of maxilla.
- caninus (kănī'nŭs) n. [L. caninus, canine.] Muscle from canine fossa to angle of mouth; levator anguli oris
- cannon bone,—bone supporting limb from hock to fetlock, enlarged and fused metacarpals or metatarsals; in birds, the tarsometatarsus.
- canopy (kăn'öpi) n. [Gk. konopeion, curtained bed.] Topmost layer of leaves, twigs, and branches of forest trees, or of other woody plants.

cantharidin (kănthăr'idin) n. [Cantharidae, blister-beetles, from Gk. kantharos.] Poison from accessory glands of genital tract and blood of blister-beetles; C₁₀H₁₈O₄.

canthus (kăn'thus) n. [Gk. kanthos, corner of eye.] The angle where upper and lower eyelids meet; com-

missura palpebrarum.

capillary (kāpīl'ārī) a. [L. capillus, hair.] Hair-like; appl. moisture held between and around particles of soil. n. One of minute thin-walled vessels which form networks in various parts of body, e.g. blood, lymph, or biliary capillaries.

capillitium (kāp'īlīt'ium, kāp'īlīsh'ium) n. [L. capillus, hair.] A protoplasmic network of elaters or filaments embedding spores within

sporangia of certain fungi.

capitate (kăp'îtât) a. [L. caput, head.] Enlarged or swollen at tip; gathered into a mass at apex, as compound stigma, some inflorescences; appl. a bone, os capitatum.

capitatum (kăpitā'tum) n. [L. caput, head.] The third carpale; os

magnum.

- capitellum (kāp'ītēl'ūm) n. [Dim. of L. caput, head.] A capitulum or articulatory protuberance at end of a bone.
- capitulum (kăpīt'ūlūm) n. [L. caput, A knob-like swelling at head.] end of a bone, e.g. on humerus for articulation with radius; part of cirripede body enclosed in mantle, opp. peduncle; swollen end of hair or tentacle; enlarged end of insect proboscis, or antenna; exsert part of head in ticks; part of column above parapet in sea-anemones; spherical cell at inner end of manubrium in Characeae; head or anthodium, an inflorescence sessile flowers or florets crowded together on a receptacle and usually surrounded by an involucre.
- capreolate (kăprē'ölāt, kăp'rëölāt) a. [L. capreolus, tendril.] Supplied with tendrils; tendril-shaped.
- caprification (kăp'rifikā'shun) n. [L.

caprificus, wild fig-tree.] Pollination of flowers of fig-tree by Chalcid insects.

- capsule (kāp'sūl) n. [L. capsula, little box.] A sac-like membrane enclosing an organ; thickened slime layer surrounding certain bacteria; any closed box-like vessel containing spores, seeds, or fruits; sporogonium, in Bryophyta; a superior, one or more celled, many-seeded, dehiscent fruit; membrane surrounding nerve-cells of sympathetic ganglia.
- capsuliferous (kăp'sūlif'ērūs) a. [L. capsula, little box; ferre, to carry.] With, or forming, a capsule; capsuligerous, capsulogenous.
- captacula (kăptăk'ūlă) n. plu. [L. captare, to lie in wait for.] Exsertile filamentous tactile organs near mouth of Scaphopoda.
- caput (kăp'ŭt) n. [L. caput, head.] Head; knob-like swelling at apex; peridium of certain fungi.
- caput caecum coli,— former name of caecum.
- carapace (kăr'ăpās) n. [Sp. carapacho, covering.] A chitinous or bony shield covering whole or part of back of certain animals.
- carbohydrates (kâr'böhī'drāts) n. plu. [L. carbo, coal; Gk. hydor, water.] Compounds of carbon, hydrogen, and oxygen, aldehydes or ketones constituting sugars, or condensation products thereof.
- carbon dioxide (kâr'bŏn diŏk'sīd)

 n. [L. carbo, coal; Gk. di-, two;
 oxys, sharp.] Carbonic acid gas,
 a heavy, colourless gas present in the
 atmosphere, assimilated by plants
 and produced by decomposition of
 organic substances; CO₂.

Carboniferous (kâr'bŏnĭf'ërŭs) a. [L. carbo, coal; ferre, to carry.] Pert. period of late Palaeozoic era including formation of coal measures.

carcerule (kâr'sërūl) n. [L. carcer, prison.] A superior, dry, many-celled fruit, with indehiscent one-or few-seeded carpels cohering by united styles to a central axis.

- carcinology (kâr'sinŏl'öjī) n. [Gk. karkinos, crab; logos, discourse.]
 The study of Crustacea.
- cardia (kar'dia) n. [Gk. kardia, stomach.] The opening between oesophagus and stomach.
- cardiac (kâr'dĭāk) a. [Gk. kardiakos, pert. heart, stomach.] Pert., near, or supplying heart; appl. cycle, etc.; pert. anterior part of stomach.
- cardiac impulse,—motion caused by rapid increase in tension of ventricle.
- cardinal (kâr'dĭnăl) a. [L. cardo, hinge.] Pert. that upon which something depends or hinges; pert. hinge of bivalve shell, or to cardo of insects; appl. points for plant growth: minimum, optimum, and maximum temperatures or temperature ranges.
- cardinal sinuses and veins,—veins uniting in Cuvier's duct, persistent in most fishes, embryonic in other vertebrates.
- cardioblast (kâr'dĭöblăst) n. [Gk. kardia, heart; blastos, bud.] One of embryonic cells destined to form walls of heart.
- cardiobranchial (kâr'diöbrăng'kiăl)
 a. [Gk. kardia, heart; brangchia, gills.] Appl. enlarged posterior basibranchial cartilage ventral to heart in elasmobranchs.
- cardo (kâr'dō) n. [L. cardo, hinge.] The hinge of a bivalve shell; basal sclerite of maxilla in insects, itself divided into eucardo and paracardo.
- carina (kărī'nă, kărē'nă) n. [L. carina, keel.] A keel-like ridge on certain bones, as breast-bone of birds; median dorsal plate of a barnacle; the two coherent anterior petals of a leguminous flower; ridge on bracts of certain grasses.
- carinal (kărī'năl) a. [L. carina, keel.] Like or pert. a keel or ridge; appl. median strand of xylem passing from stem to leaf; appl. canals in protoxylem beneath ridges of stem in Equisetales; appl. dots or puncta on keel of diatom valves.

- carinate (kār'īnāt) a. [L. carina, keel.] Having a ridge or keel.
- cariniform (kărîn'ĭfôrm) a. [L. carina, keel; forma, shape.] Keel-shaped.
- carnassial (kârnăs'iăl) a. [L. caro, flesh.] Pert. cutting teeth of Carnivora, fourth premolar above and first molar below, in upper the protocone is reduced, in lower the metaconid.
- carnivorous (kârnĭv'örŭs) a. [L. caro, flesh; vorare, to devour.] Flesh-eating; appl. Carnivora, and to certain plants which feed on entrapped insects.
- carotenase (kăr'ŏtēnās) n. [L. carota, carrot.] A liver enzyme which activates vitamin A formation from carotenes.
- carotene (kăr'ötēn) n. [L. carota, carrot.] A yellow pigment synthesised by plants and present in milk, liver oils, egg yolk, etc.; provitamin A; C₄₀H₅₈.
- carotenoids (kār'ŏtēnoidz) n. plu.

 [I.. carota, carrot; Gk. eidos, form.] Pigments occurring in plants and some animal tissues, and including carotene, xanthophylls, and other fat-soluble pigments.
- carotid (kărôt'id) a. [Gk. karos, heavy sleep.] Pert. chief arteries in the neck; appl. arch, ganglion, nerve, etc.
- carotid bodies,—two small masses of chromaffin cells associated with carotid sinus, and being part of system for controlling oxygen content and acidity of blood; glomera carotica.
- carotin,—carotene, q.v.
- carotinoids,—carotenoids, q.v.
- carpal (kâr'păl) n. [L. carpus, wrist.]
 A wrist bone. a. Pert. wrist.
- carpel (kâr'pēl) n. [Gk. karpos, fruit.] A division of the seed-vessel; a simple pistil. Plu. Sporophylls which carry megasporangia; megasporophylls.
- carpocerite (kâr'pösërīt) n. [L. carpus, wrist; Gk. keras, horn.]
 Fifth antennal joint in certain
 Crustacea.

- carpogenic (kârpöjĕn'îk) a. [Gk. karpos, fruit; gennaein, to produce.] Appl. those cells in red algae which form the carpogonium; appl. cell: oogonium of archicarp; carpogenous.
- carpogonium (kâr'pögō'nĭŭm) n. [Gk. karpos, fruit; gonos, birth.] Lower portion of procarp, which contains female nucleus, in some thallophytes; female gametangium in red algae.

carpolith (kâr'pölith) n. [Gk. karpos, fruit; lithos, stone.] A fossil fruit.

- carpometacarpus (kâr'pömětăkâr'pūs) n. [Gk. karpos, wrist; meta, after.] Portion of wing skeleton formed by fusion of carpal and metacarpal bones, in birds.
- carpophagous (kârpŏf'ăgūs) a. [Gk. karpos, fruit; phagein, to eat.] Feeding on fruit.
- carpophore (kâr'pöför) n. [Gk. karpos, fruit; pherein, to bear.] Part of flower axis to which carpels are attached; stalk of sporocarp.
- carpophyte (kâr'pöfit) n. [Gk. karpos, fruit; phyton, plant.] A
 thallophyte which forms sporocarps.
- carpopodite (kâr'pöpödīt) n. [L. carpus, wrist; Gk. pous, foot.] The third joint of endopodite in certain Crustacea; patella in spiders.

carposperm (kâr'pösperm) n. [Gk. karpos, fruit; sperma, seed.] The fertilised oosphere in certain Thallophyta.

carposporangium (kâr'pöspörăn'jĭŭm) n. [Gk. karpos, fruit; sporos, seed; anggeion, vessel.] The terminal cells of filaments developed from fertilised carpogonium in some Thallophyta.

carpospore (kar'pöspör) n. [Gk. kar-pos, fruit; sporos, seed.] A spore of those formed at end of filaments developed from carpogonium.

carposporophyte (kār'pöspō'röfīt) n.
[Ĝk. karpos, fruit; sporos, seed; phyton, plant.] The diploid generation of red algae, which consists of filaments forming carpospores at their apices.

carpus (kâr'pŭs) n. [L. carpus, wrist.]

The wrist; region of fore-limb between forearm and metacarpus.

cartilage (kâr'tIlej) n. [L. cartilago, cartilage.] Gristle, a translucent, bluish-white tissue, firm and elastic, found generally in connection with bones; cartilaginous structure.

caruncle (kărŭng'kl) n. [L. caruncula, small piece of flesh.] A naked, fleshy excrescence; small conical body at inner junction of upper and lower eyelids, caruncula lacrimalis; a fleshy outgrowth on head of certain birds, and on certain caterpillars; a little horny elevation at end of beak of embryo chicks; piston-like structure within acetabulum of dibranchiate Cephalopoda; sucking-disc on tarsi of certain mites; one of outgrowths from various regions of testa of a seed, a strophiole.

caryo-,-also karyo-, q.v.

caryolite (kăr'iölit) n. [Gk. karyon, nut; lytikos, loosing.] A nucleated muscle fragment undergoing phagocytosis in development of insects.

caryopsis (kār'iŏp'sīs) n. [Gk. karyon, nut; opsis, appearance.] A superior, one-celled, one-seeded, indehiscent fruit with a thin dry membranous pericarp inseparably united with the seed; grain.

casein (kā'sein) n. [L. caseus, cheese.] A phosphoprotein of milk, formed from caseinogen (casein in U.S.A.) by action of rennet; paracasein (U.S.A.)

Casparian band [R. Caspary, German botanist]. A cork- or wood-like strip encircling radial walls of endodermis cells.

cassideous (kăsid'eŭs) a. [L. cassis, helmet.] Helmet-like.

caste (kast) n. [L. castus, pure.] One of the distinct forms found among certain social insects.

castrate (kăs'trāt) a. [L. castrare, to castrate.] Pert. flowers from which androecium has been removed.

n. An animal deprived of functional gonads. v. To deprive of testes; to gonadectomise; to inhibit development of gonads.

cata-,-also kata-, q.v.

catadromous (kătăd'römus) a. [Gk. kata, down; dromein, to run.] Tending downward; having branches arising from lower side of pinnae, in ferns; having first set of nerves in a frond segment given off on basal side of midrib; appl. fishes which migrate from fresh to salt water annually, opp. anadromous.

catalase,-an enzyme occurring in plant and animal tissues, which decomposes hydrogen peroxide into

water and oxygen.

catalepsis (kătălěp'sis) n. katalepsis, seizure.] A so-called shamming - dead reflex. spiders; cf. kataplexy.

catalysis (kătăl'isis) n. [Gk. katalysis, dissolving.] Acceleration or retardation of reaction due to

presence of a catalyst.

catalyst (kăt'ălĭst) n. [Gk. katalysis, dissolving.] An agent, e.g. an enzyme, which can accelerate or retard, or initiate, a reaction and apparently remains unchanged; catalysor.

cataphoresis (kăt'ăförē'sis) n. [Gk. katapherein, to carry down.] Migration of particles in suspension, as of living cells, under influence of electric current, the rate depending on voltage; electrophoresis.

cataphyll (kăt'ăfil) n. [Gk. kata, phyllon, leaf.] Simple down; form of leaf on lower part of plant, as cotyledon, bud-scale, scale-leaf; cataphyllary leaf. Opp. hypsophyll.

cataphyllary (kàt'āfìl'ārī) a. kata, down; phyllon, leaf.] Appl. rudimentary or scale-like leaves which act as covering of buds.

catapleurite (kăt'aploor'īt) n. [Gk. kata, down; pleura, side.] Thoracic pleurite between anapleurite and trochantin, as in certain Thysanura; coxopleurite.

catelectrotonus (kătëlěk'trŏt'önŭs, kăt'ëlěktrŏt'önŭs) n. [Gk. kata, down; elektron, amber; tonos, tension.] Increase in irritability of a

nerve under influence of non-polarising electric current; katelectrotonus.

[L. catena, catena (kătē'nă) n. chain.] A sequence of soil types which is repeated in a corresponding sequence of topographical sites, as between ridges and valleys of a region.

catenation (kătënā'shun) n. catenatus, chained.] End-to-end arrangement of chromosomes; ring formation of alternating paternally and maternally derived chromosomes; a chain, as of diatom frustules.

catenoid (kătē'noid) a. [L. catena, chain; Gk. eidos, form.] Chain-like; appl. certain protozoan colonies.

catenular (kătěn'ūlăr), catenulate (kătěn'ūlāt) a. [L. catena, chain.] Chain-like; appl. colonies of bacteria, colour-markings on butterfly wings, shells, etc.

caterpillar (kăt'erpil'ăr) n. [L.L. cattus, cat; L. pilosus, hairy.] Young worm-like insect larva, particularly of Lepidoptera; eruca.

cathammal (kăth'ămăl) a. kathamma, anything tied.] Appl. plates forming endoderm lamella in some Coelenterata.

catkin (kăt'kin) n. [A.S. catkin, little cat.] A spike with unisexual flowers and pendulous rachis; amentum.

cauda (kô'dă) n. [L. cauda, tail.] A tail, or tail-like appendage; posterior part of an organ, e.g. cauda equina, cauda epididymis; a tube at posterior end of abdomen of certain insects, suggesting presence of a further segment.

caudad (kô'dăd) adv. [L. cauda, tail; ad, toward. Towards tail region or posterior end.

caudal (kô'dăl) a. [L. cauda, tail.] Of or pert. a tail, e.g. caudal fin.

caudate (kô'dāt) a. [L. cauda, tail.] Having a tail, e.g. caudate nucleus; appl. a lobe of the liver.

caudatolenticular (kôdā'tölčntĭk'ūlăr) a. [L. cauda, tail; lens, lentil.] Appl. caudate and lenticular nuclei of corpus striatum.

- caudex (kô'dēks) n. [I.. caudex, dry stump.] The axis or stem of a woody plant, as of tree-ferns, palms, etc.
- caudicle (kô'díkl) n. [Dim. of L. cauda, tail] Stalk of pollinium in orchids.
- caudihaemal (kô'dìhē'māl) a. [L. cauda, tail; Gk. haima, blood.] Appl. posterior lower portion of a sclerotome.
- caudineural (kôdinū'iăl) a. [L. cauda, tail; Gk. neuron, nerve.]
 Appl. posterior upper portion of a sclerotome.
- caudostyle (kô'döstīl) n. [L. cauda, tail; Gk. stylos, column.] A terminal structure in certain parasitic amoebae.
- caul (kôl) n. [M.E. calle, covering.]
 An enclosing membrane; amnion; omentum.
- caulescent (kôlěs'ënt) a. [L. caulis, stalk.] With leaf bearing stem above ground.
- caulicolous (kôlīk'ölūs) a. [I.. caults, stalk; colere, to inhabit.] Appl. fungi growing on plant-stems.
- cauliflory (kôl'íflō'ri) n. [L. caulis, stalk; flos, flower.] Condition of having flowers arising from axillary buds on main stem or older branches; cauliflorous habitus.
- cauliform (kôl'ifôrm) a. [I.. caulis, stalk; forma, shape.] Stem-like.
- cauligenous (kôlij'ënŭs) a. [Gk. kaulos, stem; genos, birth.] Borne on the stem.
- cauline (kô'līn) a. [I.. caulis, stalk.] Pert. stem; appl. leaves growing on upper portion of a stem; appl. vascular bundles not passing into leaves.
- caulis (kô'lĭs) n. [L. caulis, stalk.] The stem, in herbaceous plants.
- caulocaline (kôl'ökălēn') n. [Gk. kaulos, stem; kalein, to summon.] A plant hormone, possibly elaborated in roots, which stimulates growth of stem.
- caulocarpous (kô'lökâr'pŭs) a. [Gk. kaulos, stem; karpos, fruit.] With fruit-bearing stem.
- caulocystidium (kôl'ösistid'iŭm) n.

- [Gk. kanlos, stalk; kystis, bag; idion, dim.] One of the cystidium-like structures on stipe of certain Basidiomycetes.
- caulome (kô'lōm) n. [Gk. kaulos, stem.] The stem structure of a plant as a whole.
- cavernosus (kăv'ërnō'sūs) a. [L. cavernosus, chambered.] Full of cavities; hollow, or resembling a hollow; appl. tissue, nerve, arteries.
- cavicorn (kav'íkôrn) a. [L. cavus, hollow; cornu, horn.] Hollow-horned; appl. certain ruminants.
- cavum (kā²vum) n. [L. cavus, hollow.] The lower division of concha caused by origin of helix; cavity of mouth, larynx, long bones, etc.; any hollow or chamber.
- cecal, -- caecal, q.v.
- cecum,-caecum, q.v.
- celiac,—coeliac, q.v.
- cell (sĕl) n. [L. cella, compartment.]
 A small cavity or hollow; a loculus; a unit mass of protoplasm, usually containing a nucleus or nuclear material; originally, the cell wall; space between veins of insect wings.
- cellifugal (sĕlĭf'ūgăl) a. [L. cella, cell; fugere, to flee.] Moving away from a cell.
- cellipetal (sĕlĭp'étăl) a. [L. cella, cell; petere, to seek.] Moving towards a cell.
- cell lineage,—the derivation of a tissue or part from a definite blastomere of embryo.
- cell organ,—a part of a cell having a special function, as a centrosome; organoid.
- cell plate,—equatorial thickening of spindle fibres from which partition wall arises during division of plant cells.
- cell sap,—the more fluid ground substance of the cell.
- cellular (sĕl'ūlăr) a. [L. cellula, small cell.] Pert. or consisting of cells.
- cellulase (sĕl'ūlās) n. [L. cellula, small cell.] An enzyme which hydrolyses cellulose, occurring in bacteria and fungi.

cellulin (sĕl'ūlin) n. [L. cellula, little cell.] A carbohydrate found in constrictions of hyphae.

cellulose (sěl'ūlōs) n. [L. cellula, small cell.] A carbohydrate forming main part of plant cell walls, also found in tests of tunicates; (C₈H₁₀O₅)_x.

cell-wall,—investing portion of cell.

celo-,—coelo-, q.v.

cement (sëměnt') n. [L. caementum, mortar.] A substance chemically and physically allied to bone, investing parts of teeth; crusta petrosa; a uniting substance secreted by certain animals.

cenchrus (seng'krus) n. [Gk. kengchros, millet.] A pale-coloured area on mesothorax of saw-flies.

ceno-,-see caeno-, coeno-.

censer mechanism,—method of seed distribution by which seeds are jerked out from fruit by high wind.

- centradenia (sĕn'trădē'niā) n. [Gk. kentron, centre; aden, gland.] The type of siphonophore colony in Disconectae.
- central (sĕn'trăl) a. [L. centrum, centre.] Situated in the centre; pert. a vertebral centrum. n. A bone in wrist or ankle, situated between proximal and distal rows.

central body,—centrosome, q.v.

central cylinder,—stele, q.v.

- centric (sĕn'trīk) a. [L. centrum, centre.] Appl. leaves which are cylindrical or terete; having a centromere.
- centrifugal (sĕntrif'ūgăl) a. [L. centrum, centre; fugere, to flee.] Appl. compact cymose inflorescences having youngest flowers towards outside; appl. nerves transmitting impressions from nerve centre to parts supplied by nerve.
- centriole (sĕn'triōl) n. [L. centrum, centre.] The central particle of the centrosome; the centrosome itself.
- centripetal (sěntrip'étăl) a. [L. centrum, centre; petere, to seek.]
 Appl. racemose inflorescences having youngest flowers at apex;
 appl. nerves transmitting impres-

sions from peripheral extremities to nerve centres.

- centripetal canals,—blind canals growing from circular canal backwards towards apex of bell in certain Trachomedusae.
- centro-acinar (sen'troas'inar) a. [L. centrum, centre; acinus, berry.]

 Pert. centre of an alveolus, as in pancreas.
- centrodesmose (sĕn'trödĕs'mōs), centrodesmus (sĕn'trödĕs'mūs) n. [Gk. kentron, centre; desmos, bond.] The fibril or system of fibrils temporarily connecting two centrosomes; also centrodesm.
- centrodorsal (sĕn'trödôr'săl) a. [L. centrum, centre; dorsum, back.] Appl. plate in middle of aboral surface of unstalked crinoids.
- centrogenous (sĕntrŏj'ēnŭs) a. [Gk. kentron, centre; gennaein, to produce.] Appl. a skeleton of spicules which meet in a common centre and grow outwards.
- centrolecithal (sĕn'trölĕs'ithăl) a. [Gk. kentron, centre; lekithos, yolk.] With yolk aggregated in the centre, appl. ovum.
- centromere (sĕn'trömēr) n. [Gk. kentron, centre; meros, part.] The part of the chromosome located at the point lying on the equator of the spindle at metaphase and dividing at anaphase, controlling chromosome activity; spindle-attachment region, achromite, kinetochore.

centron (sĕn'trŏn) n. [Gk. kentron, centre.] Cyton, q.v.

- centrophormium (sen'tröfôr'miŭm)

 n. [Gk. kentron, centre; phormis,
 small basket.] The Golgi-bodies
 when in round basket like
 form.
- centroplasm (sĕn'tröplăzm) n. [Gk. kentron, centre; plasma, mould.]
 Substance of centrosphere; a more or less definite concentric zone round the aster in mitosis.
- centroplast (sěn'tröpläst) n. [Gk. kentron, centre; plastos, formed.]
 An extranuclear spherical body forming division centre of mitosis in certain lower organisms.

- centrosome (sěn'trösôm) n. [Gk. kentron, centre; soma, body.] A cell-organ, the centre of dynamic activity in mitosis, consisting of centriole and attraction-sphere.
- centrosphere (sĕn'trösfēr) n. [Gk. kentron, centre; sphaira, ball.] The central mass of aster and centrosome; astrosphere; attraction-sphere.
- centrotaxis (sĕn'trotăk'sīs) n. [Gk. kentron, centre; taxis, arrangement.] Orientation of chromatin thread towards cytocentrum during leptotene stage.
- centrotheca (sĕn'tröthē'kă) n. [Gk. kentron, centre; theke, case.] Idiozome, q.v.
- centrum (sěn'trům) n. [L. centrum, centre.] The main body of a vertebra, from which neural and haemal arches arise; centrosome, centrosphere, q.v.

cephal-,-also kephal-.

- cephalad (kĕf'ālăd, sĕf-) adv. [Gk. kephale, hcad; L. ad, towards.] Towards head region or anterior end.
- cephalanthium (kĕf'ălăn'thĭum, sĕf-)
 n. [Gk. kephale, head; anthos, flower.] The capitulum in composite plants; anthodium.
- cephaletron (kĕfālē'trŏn, sĕf-) n. [Gk. kephale, head; etron, belly.] The anterior region of Xiphosura.
- cephalic (kĕfăl'īk, sĕf-) a. [Gk. kephale, head.] Pert. head; in head region.
- cephalic index,—one hundred times maximum breadth divided by maximum length of skull.
- cephalin (kĕf'ālīn, sĕf-) n. [Gk. kephale, head.] A phospholipide present in nerve fibres and egg-yolk; kephalin; an epimerite bearing trophozoites.
- cephalis (kĕf'ālīs, sĕf-) n. [Gk. kephalis, little bulb.] The uppermost chamber of monaxonic shells of Radiolaria.
- cephalisation (kĕf'ālīzā'shun, sĕf-) n. [Gk. kephale, head.] Increasing differentiation and importance

- of anterior end in animal development.
- cephalon (kĕf'ālŏn, sĕf-)n. [Gk. kephale, head.] The head of arthropods; head shield of trilobites.
- cephalont (kĕf'ālŏnt, sĕf-) n. [Gk. kephale, head.] A sporozoan about to proceed to spore-formation.
- cephalopod (kěf'álöpöd, sěf-) n. [Gk. kephale, head; pous, foot.] Marine mollusc with muscular suckerbearing arms on head region, e.g. cuttle-fish, octopus.
- cephalopodium (kĕf'ălöpō'dĭŭm, sĕf-)
 n. [Gk. kephale, head; pous, foot.]
 The head and arms constituting
 the head-region in cephalopods.
- cephalopsin (kěfálŏp'sin, sěf-) n. [Gk. kephale, head; opsis, sight.] A photopigment resembling visual purple, in eyes of cephalopods and some other invertebrates.
- cephalostegite (kĕf'ălŏs'tëjīt, sĕf-)

 n. [Gk. kephale, head; stege, roof.]

 Anterior part of cephalothoracic shield.
- cephalostyle (kĕf'ālostīl, sĕf-) n. [Gk. kephale, head; stylos, pillar.] Anterior end of notochord enclosed in sheath, in Chondrocrania.
- cephalotheca (kĕf'ālöthē'kā, sĕf-) n. [Gk. kephale, head; theke, case.] Head integument in insect pupa.
- cephalothorax (kĕf'ălöthō'rāks, sĕf-)

 n. [Gk. kephale, head; thorax,
 breast.] The body-region formed
 by fusion of head and thorax in
 Arachnida and Crustacea.
- cephalotrocha (kĕfālŏt'rökā, sĕf-) n. [Gk. kephale, head; trochos, wheel.] A turbellarian larva with eight processes round mouth.
- cephalula (kĕfāl'ūlă, sĕf-) n. [Gk. kephale, head.] Free-swimming embryonic stage in certain brachiopods. cer-,—also ker-.
- cerata (sĕr'ātā, kĕr-) n. plu. [Gk. keras, horn.] Lobes or leaf-like processes acting as gills on back of nudibranch molluscs.
- ceratium (sërā'shīŭm) n. [Gk. keration, little horn.] A siliqua without the replum.

- ceratobranchial (ker'atöbrang'kiāl, ser-). [Gk. keras. horn; brangchia, gills.] An element of branchial arch.
- ceratohyal (kĕr'ātöhī'āl, sĕr-) n. [Gk. keras, horn; hyoeides, T-shaped.]
 The component of hyoid arch next below epihyal.
- ceratotrichia (kěr'átötrĭk'ĭā, sĕr-) n. plu. [Gk. keras, horn; thrix, hair.]
 Horny and non-cellular actinotrichia of elasmobranchs.
- cercal (sĕr'kăl) a. [Gk. kerkos, tail.] Pert. the tail; pert. cerci, appl. hairs, nerve.
- cercaria (sĕrkā'rīă) n. [Gk. kerkos, tail.] A heart-shaped trematode larva with tail.
- cerci,—plu. of cercus.
- cercid (sĕr'sid) n. [Gk. kerkis, shuttle.] One of minute wandering cells produced by division of archaeocytes in certain sponges.
- cercoid (sĕr'koid) n. [Ğk. kerkos, tail; eidos, shape.] One of paired appendages on ninth, or tenth, abdominal segment of certain insect larvae.
- cercus (sĕr'kŭs) n. [Gk. kerkos, tail.]
 A jointed appendage at end of abdomen in many arthropods; appendage bearing acoustic hairs in some insects.
- cere (sēr) n. [L. cera, wax.] A swollen fleshy patch at proximal end of bill in birds.
- cerebellar (sĕr'ēbĕl'ar) a. [L. cerebrum, brain.] Pert. the cerebellum or hind-brain.
- cerebellum (sĕr'ebĕl'ŭm) n. [L. cerebrum, brain.] The fourth division of brain, arising from differentiation of anterior part of third primary vesicle.
- cerebral (sĕr'ĕbrăl) a. [L. cerebrum, brain.] Pert. the brain; pert. anterior part of brain or cerebral hemispheres.
- cerebral organs,—chemical sense organs, paired ciliated tubes associated with dorsal ganglion and opening to exterior, in nemertines.
- cerebrifugal (sĕrëbrif'ūgăl) a. [L. cerebrum, brain; fugere, to flee.]

- Appl. nerve fibres which pass from brain to spinal cord.
- cerebroganglion (sĕr'ëbrögăng'gliŏn) n. [L. cerebrum, brain; Gk. ganglion, swelling.] The supraoesophageal ganglia of invertebrates.
- cerebropedal (sĕr'ebröpēd'ăl) a. [L. cerebrum, brain; pes, foot.] Appl. nerve strands connecting cerebral and pedal ganglia in molluscs.
- cerebrospinal (sĕr'ëbrospī'năl) a. [L. cerebrum, brain; spina, spine.]
 Pert. brain and spinal cord.
- cerebrovisceral (ser'ebrovis'eral) a. [L. cerebrum, brain; viscera, viscera.] Appl. connective joining cerebral and visceral ganglia in molluscs.
- cerebrum (sĕr'ebrum) n. [L. cerebrum, brain.] The fore-brain, arising from differentiation of first primary vesicle.
- cereous (sē'reŭs) a. [L. cereus, waxen.] Wax-like.
- ceriferous (sērĭf'ērŭs) a. [L. cera, wax; ferre, to carry.] Wax-producing; appl. organs.
- cernuous (ser'nūŭs) a. [L. cernuus, with face turned downwards.]
 Drooping; pendulous.
- ceroma (sē'rōmă) n [Gk. keroma, ointment.] The cere of birds.
- cerous (sē'rŭs) a. [L. cera, wax.]
 Appl. structure resembling a cere.
- certation (sĕrtā'shŭn) n. [L. certatio, contest.] Competition in growth rate of pollen tubes of genetically different types.
- cerumen (serū'mēn) n. [L. cera, wax.] Wax-like secretion from ceruminous glands of ear; wax secreted by scale insects; wax of nest of certain bees.
- cervical (ser'vikal, servi'kal) a. [L. cervix, neck.] Appl. or pert. structures connected with neck, as nerves, bones, blood-vessels, also to cervix or neck of an organ.
- cervicum (sĕr'vĭkŭm) n. [L. cervix, neck.] The neck-region of Arthropoda.
- cervix (sĕr'vĭks) n. [L. cervix, neck.]
 The neck or narrow mouth of an organ, as cervix uteri.

- centrosome (sěn'trösōm) n. [Gk. kentron, centre; soma, body.] A cell-organ, the centre of dynamic activity in mitosis, consisting of centriole and attraction-sphere.
- centrosphere (sĕn'trosfēr) n. [Gk. kentron, centre; sphaira, ball.]
 The central mass of aster and centrosome; astrosphere; attraction-sphere.
- centrotaxis (sĕn'tròtăk'sīs) n. [Gk. kentron, centre; taxis, arrangement.] Orientation of chromatin thread towards cytocentrum during leptotene stage.
- centrotheca (sĕn'tröthē'kă) n. [Gk. kentron, centre; theke, case.] Idiozome, q.v.
- centrum (sen'trum) n. [L. centrum, centre.] The main body of a vertebra, from which neural and haemal arches arise; centrosome, centrosphere, q.v.

cephal-,-also kephal-.

- cephalad (kĕf'ālăd, sĕf-) adv. [Gk. kephale, head; L. ad, towards.] Towards head region or anterior end.
- cephalanthium (kĕf'ălăn'thĭŭm, sĕf-)
 n. [Gk. kephale, head; anthos, flower.] The capitulum in composite plants; anthodium.
- cephaletron (kĕfālē'trŏn, sĕf-) n. [Gk. kephale, head; etron, belly.] The anterior region of Xiphosura.
- cephalic (kĕfăl'ĭk, sĕf-) a. [Gk. kephale, head.] Pert. head; in head region.
- cephalic index,—one hundred times maximum breadth divided by maximum length of skull.
- cephalin (kĕf'ālĭn, sĕf-) n. [Gk. kephale, head.] A phospholipide present in nerve fibres and egg-yolk; kephalin; an epimerite bearing trophozoites.
- cephalis (kĕf'ālĭs, sĕf-) n. [Gk. kephalis, little bulb.] The uppermost chamber of monaxonic shells of Radiolaria.
- cephalisation (kĕf'ālīzā'shŭn, sĕf-) n. [Gk. kephale, head.] Increasing differentiation and importance

- of anterior end in animal development.
- cephalon (kĕf'ālŏn, sĕf-)n. [Gk. kephale, head.] The head of arthropods; head shield of trilobites.
- cephalont (kĕf'ālŏnt, sĕf-) n. [Gk. kephale, head.] A sporozoan about to proceed to spore-formation.
- cephalopod (kĕf'ălöpŏd, sĕf-) n. [Gk. kephale, head; pous, foot.] Marine mollusc with muscular suckerbearing arms on head region, e.g. cuttle-fish, octopus.
- cephalopodium (kĕf'ălöpō'dĭŭm, sĕf-)
 n. [Gk. kephale, head; pous, foot.]
 The head and arms constituting
 the head-region in cephalopods.
- cephalopsin (kĕfālŏp'sĭn, sĕf-) n. [Gk. kephale, head; opsis, sight.] A photopigment resembling visual purple, in eyes of cephalopods and some other invertebrates.
- cephalostegite (kĕf'ălŏs'tëjīt, sĕf-)
 n. [Gk. kephale, head; stege, roof.]
 Anterior part of cephalothoracic shield.
- cephalostyle (kĕf'ālöstīl, sĕf-) n. [Gk. kephale, head; stylos, pillar.] Anterior end of notochord enclosed in sheath, in Chondrocrania.
- cephalotheca (kĕf'álöthē'kā, sĕf-) n. [Gk. kephale, head; theke, case.] Head integument in insect pupa.
- cephalothorax (kĕf'ālöthō'rāks, sĕf-)

 n. [Gk. kephale, head; thorax,
 breast.] The body-region formed
 by fusion of head and thorax in
 Arachnida and Crustacea.
- cephalotrocha (kĕfālŏt'rökā, sĕf-) n. [Gk. kephale, head; trochos, wheel.] A turbellarian larva with eight processes round mouth.
- cephalula (kĕfăl'ūlă, sĕf-) n. [Gk. kephale, head.] Free-swimming embryonic stage in certain brachiopods. cer-,—also ker-.
- cerata (sēr'ātā, kēr-) n. plu. [Gk. keras, horn.] Lobes or leaf-like processes acting as gills on back of nudibranch molluscs.
- ceratium (sërā'shǐum) n. [Gk. keration, little horn.] A siliqua without the replum.

CER-77 CER-

- ceratobranchial (kčr'atöbrang'kĭal, sèr-). [Gk. keras, horn; brangchia, gills.] An element of branchial arch.
- ceratohyal (ker'atöhī'al, ser-) n. [Gk. keras, horn; hyoeides, Y-shaped.] The component of hyoid arch next below epihyal.
- ceratotrichia (kěr'ătötrik'iă, sěr-) n. plu. [Gk. keras, horn; thrix, hair.] Horny and non-cellular actinotrichia of elasmobranchs.
- cercal (sĕr'kăl) a. [Gk. kerkos, tail.] Pert. the tail; pert. cerci, appl. hairs, nerve.
- cercaria (sěrkā'rĭă) n. [Gk. kerkos, tail.] A heart-shaped trematode larva with tail.
- cerci,-plu. of cercus.
- cercid (ser'sid) n. [Gk. kerkis, shuttle.] One of minute wandering cells produced by division of archaeocytes in certain sponges.
- cercoid (sĕr'koid) n. [Gk. kerkos, tail; eidos, shape.] One of paired appendages on ninth, or tenth, abdominal segment of certain insect larvae.
- cercus (ser'kus) n. [Gk. kerkos, tail.] A jointed appendage at end of abdomen in many arthropods; appendage bearing acoustic hairs in some insects.
- cere (sēr) n. [L. cera, wax.] A swollen fleshy patch at proximal end of bill in birds.
- cerebellar (sěr'ëběl'ăr) a. [L. cerebrum, brain.] Pert. the cerebellum or hind-brain.
- cerebellum (sěr'eběl'ŭm) n. [L. cerebrum, brain.] The fourth division of brain, arising from differentiation of anterior part of third primary vesicle.
- cerebral (sĕr'ëbrăl) a. [L. cerebrum, Pert. the brain; pert. anterior part of brain or cerebral hemispheres.
- cerebral organs,-chemical sense paired ciliated organs, tubes associated with dorsal ganglion and opening to exterior, in nemer-
- cerebrifugal (serebrif'ūgăl) a. [L. cerebrum, brain; fugere, to flee.]

Appl. nerve fibres which pass from brain to spinal cord.

- cerebroganglion (sĕr'ëbrögăng'glion) n. [L. cerebium, brain; Gk. ganglion, swelling.] The supraoesophageal ganglia of invertebrates.
- cerebropedal (sěr'ebröpěd'al) a. [L. cerebrum, brain; pes, foot.] Appl. nerve strands connecting cerebral and pedal ganglia in molluscs.
- cerebrospinal (sěr'ebröspī'năl) a. [L. cerebrum, brain; spina, spine.] *Pert.* brain and spinal cord.
- cerebrovisceral (ser'ebrovis'eral) a. [L. cerebrum, brain; viscera, viscera. Appl. connective joining cerebral and visceral ganglia in molluscs.
- cerebrum (ser'ebrum) n. [L. cerebrum, brain.] The fore-brain, arising from differentiation of first primary vesicle.
- cereous (sē'reŭs) a. waxen.] Wax-like. [L. cereus,
- ceriferous (sērif'ërŭs) a. [L. cera, wax; fcrre, to carry.] producing; appl. organs.
- cernuous (sĕr'nūŭs) a. [L. cernuus, with face turned downwards.] Drooping; pendulous.
- ceroma (sē'rōmă) n [Gk. keroma, ointment.] The cere of birds.
- cerous (sē'rŭs) a. [L. cera, wax.] Appl. structure resembling a cere.
- certation (sĕrtā'shun) n. [L. certatio, contest.] Competition in growth rate of pollen tubes of genetically different types.
- cerumen (serū'měn) n. [L. cera, Wax-like secretion from ceruminous glands of ear; wax secreted by scale insects; wax of nest of certain bees.
- cervical (ser'vikal, servi'kal) a. [L. cervix, neck.] Appl. or pert. structures connected with neck, as nerves, bones, blood-vessels, also to cervix or neck of an organ.
- cervicum (sěr'vikům) n. [L. cervix, neck.] The neck-region of Arthropoda.
- cervix (sěr'víks) n. [L. cervix, neck.] The neck or narrow mouth of an organ, as cervix uteri.

- cespitose, -caespitose, q.v.
- cetolith (se'tölith) n. [Gk. ketos, whale; lithos, stone.] The fused tympanic and petrosal of whales, found in deep-sea dredging.
- cevitamic acid,—ascorbic acid or vitamin C.
- chaeta (kē'tă) n. [Gk. chaite, hair.] A seta (q.v.), or bristle, as of certain worms.
- chaetic (kē'tik) a. [Gk. chaite, hair.] Bristle-like, appl. a type of tactile sensilla in insects.
- chaetiferous (kētīf'ērūs) a. [Gk. chaite, hair; L. ferre, to bear.] Bristle-bearing; chaetigerous, setigerous.
- chaetophorous (kētŏi'örŭs) a. [Gk. chaite, hair; pherein, to bear.] Bristle-bearing; appl. worms and certain insects.
- chaetosema (kē'tösē'mā) n. [Gk. chaite, hair; sema, sign.] One of two small sensory organs located on head of certain Lepidoptera, and provided with bristles and sensory cells connected by a sheathed nerve to brain; Jordan's organ.
- chaetotamy (kē'tötāk'sĭ) n. [Gk. chaite, hair; taxis, arrangement.]
 Bristle pattern or arrangement.
- chain behaviour,—a series of actions, each being induced by the antecedent action and being an integral part of a unified performance.
- chalaza (kălā'ză) n. [Gk. chalaza, hail.] One of two spiral bands attaching yolk to membrane of a bird's egg; base of nucellus of ovule, from which integuments arise.
- chalaziferous (kălăzif'ērūs) a. [Gk. chalaza, hail; L. ferre, to bear.]
 Appl. layer of albumen surrounding yolk and continuous with chalazae.
- chalazogamy (kălăzŏg'ămĭ) n. [Gk. chalaza, hail; gamos, marriage.] Fertilisation in which the pollentube pierces chalaza of ovule; cf. porogamy.
- chalice (chal'is) n. [L. calix, goblet.]

 Appl. simple gland cells or goblet cells; a modified columnar epithe-

- lial gland cell; arms and disc of a crinoid.
- chalones (kăl'ōnz) n. plu. [Gk. chalinos, curb.] Internal secretions which depress activity; opp. hormones.
- chalonic (kălŏn'īk) a. [Gk. chalinos, curb.] Depressor, inhibitory, or restraining; appl. internal secretions; opp. hormonic.
- chamaephyte (kămī'fit) n. [Gk. chamai, on the ground; phyton, plant.] A plant with shoots that bear dormant buds lying on or near the ground.
- chasmatoplasm (kāz'mătöplăzm) n. [Gk. chasma, expanse; plasma, mould.] An expanded form of plasson.
- chasmochomophyte (kăz'mökō' möfit) n. [Gk. chasma, opening; choma, mound; phyton, plant.] A plant growing on detritus in rock crevices.
- chasmogamy (kāzmōg'āmĭ) n. [Gk. chasma, opening; gamos, marriage.]
 Opening of a mature flower to ensure fertilisation, opp. cleistogamy.
- chasmophyte (kaz'möfit) n. [Gk. chasma, opening; phyton, plant.]
 A plant which grows in crevices of rocks; a chasmophilous plant.
- cheek (chēk) n. [A.Ś. céace, cheek.] The fleshy wall of mouth in mammals; side of face; in invertebrates the lateral portions of head, as fixed and free cheeks of trilobites.
- cheilocystidium (kī'lösīstĭd'ĭūm) n. [Gk. cheilos, edge; kystis, bag; idion, dim.] A cystidium in hymenium at edge of lamella; cf. pleurocystidium.
- cheiropterygium (kiröpterij'iŭm) n. [Gk. cheir, hand; pteryx, wing.] The pentadactyl limb typical of higher vertebrates.
- chela (kē'lā) n. [Gk. chele, claw.] The claw borne on certain limbs of Crustacea and Arachnoidea; a short sponge spicule with talon-like projections at one or each end.
- chelate (kē'lāt) a. [Gk. chele, claw.]
 Claw-like or pincer-like; cheliform; cheliferous, q.v.

- chelicerae (kēlis'ērē) n. plu. [Gk. chele, claw; keras, horn.] Anterior chelate or sub-chelate appendages of Arachnoidea; also cheliceres.
- cheliferous (kēlif'ërūs) a. [Gk. chele, claw; L. ferre, to bear.] Supplied with chelae or claws.
- cheliform (kē'lifôrm) a. [Gk. chele, claw; L. forma, shape.] Claw-like; appl. appendages.
- cheliped (kē'lipĕd) n. [Gk. chele, claw; L. pes, foot.] A claw-bearing appendage; forceps of decapod crustaceans.
- chelophores (kē'löfōrz) n. plu. [Gk. chele, claw; pherein, to bear.] First pair of appendages in Pycnogonida.
- chemiluminescence (kěm'ilūmĭněs' ens. -loo-) n. [Gk. chemeia, transmutation; L. luminescere, to grow light.] Light production at ordinary temperature during a chemical reaction, as bioluminescence, q.v.
- chemonasty (kem'önas'ti) n. [Gk. chemcia, transmutation; nastos, close pressed.] Response to diffuse or indirect chemical stimuli.
- chemoreceptor (kĕm'örësĕp'tŏr) n. |Gk. chemeia, transmutation; L. recipere, to receive.] A terminal organ receiving chemical stimuli.
- chemoreflex (kem'ore'fleks) n. [Gk. chemeia, transmutation; L. reflectere, to bend back.] A reflex caused by chemical stimulus.
- chemosynthesis (kěm'ösĭn'thesĭs) n. [Gk. chemeia, transmutation; syn, with; tithenai, to place.] The building up of chemical compounds in organisms.
- chemotaxis (kēm'ötāk'sīs) n. [Gk. chemeia, transmutation; taxis, arrangement.] The reaction of cells or freely motile organisms to chemical stimuli; also chemiotaxis.
- chemotrophic (kěmötrŏf'īk) a. [Gk. chemeia, transmutation; trophe, nourishment.] Deriving nourishment from certain inorganic substances, appl. certain organisms without chlorophyll, as iron bacteria and sulphur bacteria.

- chemotropism (kěmŏt'röpizm) n. [Gk. chemeia, transmutation; trope, turn.] Curvature of a plant or plant organ in response to chemical stimuli.
- chernozem (cher'nözem, chernözyom)
 n. [Russ. chernyi, black; zemlya, soil.] Black soil, characteristic of steppe and grass land and formed under continental climatic conditions; blackearth.
- chersophyte (ker'sofit) n. [Gk. chersa, waste places; phyton, plant.] A plant which grows on waste land.
- chestnut soils,—dark-brown soils of semi-arid steppe-lands, fertile under adequate rainfall or when irrigated.
- cheta,—chaeta, q.v.
- chevron (shĕv'rŏn) a. [F. chevron, rafter, from L. caper, goat.] Appl V-shaped bones articulating with ventral surface of spinal column in caudal region of many vertebrates.
- chiasma (kł̃áz'mā) n., chiasmata (kł̃az'mātā) plu. [Gk. chiasma, cross.] A decussation of fibres, as optic chiasma; in paired chromatids, an exchange of partners in meiosis.
- chiasmatypy (kłaś/mati'pi) n. [Gk. chiasma, cross; typos, character.] A form of recombination of chromosome material in synapsis; chiasmatype, appl. theory that chiasmata and crossing-over are causally correlated.
- chiastic (kšas'tik) a. |Gk. chiastos, diagonally arranged.| Decussating; crossing; obliquely or at right angles to axis; pert. chiasmata.
- chiastoneural (kłas'tönū'rāl) a. [Gk. chiastos, diagonally arranged; neuron, nerve.] Appl. certain gastropods in which visceral nerve cords cross and form a figure 8.
- chilaria (kīlā'rīā) n. plu. [Gk. cheilos, lip.] Pair of processes between sixth pair of appendages in Limulus. childium (kīlīd'iūm) n. [Gk. cheilos
- chilidium (kilid'iŭm) n. [Gk. cheilos, lip; idion, dim.] A shelly plate covering deltidial fissure in dorsal valve of certain Brachiopoda.

- chimaera (kimē'rā) n. [L. chimaera, monster.] A single organism developing from two fused rudiments from different individuals, or composed of tissues of two different genotypes; a mosaic.
- chimonophilous (kimönöf'ilüs) a. [Gk. cheimon, winter; philein, to love.] Thriving or growing during winter.
- chiropterophilous (kīrŏp'tërŏf'ĭlūs)
 a. [Gk. cheir, hand; pteron, wing; philos, loving.] Pollinated by agency of bats.
- chitin (kī'tīn) n. [Gk. chiton, tunic.] A nitrogenous carbohydrate derivative forming the skeletal substance in arthropods, also occurring in fungi.
- chitinase (kī'tīnās) n. [Gk. chiton, tunic.] An enzyme which hydrolyses chitin, in mould fungi and in digestive juice of snail.
- chlamydate (klăm'idāt) a. [Gk. chlamys, cloak.] Supplied with a mantle.
- chlamydeous (klămid'eŭs) a. [Gk. chlamys, cloak.] Pert. flower and envelope.
- chlamydospore (klăm'idöspōr) n. [Gk. chlamys, cloak; sporos, seed.] A thick-walled resting spore of certain fungi and protozoa.
- chloragen (klō'rājēn) a. [Gk. chloros, sandy yellow; genos, descent.] Appl. yellow cells found in connection with alimentary canal of annelids; also chloragogen.
- chloragocyte (klô'răgösīt) n. [Gk. chloros, sandy yellow; kytos, hollow.] A chloragogen cell. chloragogen,—chloragen, q.v.
- chloragosomes (klō'răgosoms) n.
 plu. [Gk. chloros, sandy yellow;
 soma, body.] Yellow or brownish
 globules formed in chloragogen cells.
- chloranthy (klōrăn'thĭ, klō'rānthĭ) n. [Gk. chloros, grass green; anthos, flower.] Reversion of floral leaves back to ordinary green leaves.
- chlorenchyma (kloreng'kıma) n. [Gk. chloros, grass green; engchyma, infusion.] Tissues collectively, or stem tissue, or mesophyll, containing chlorophyll.

- chlorocruorin (klō'rökroo'ŏrin) n. [Gk. chloros, grass green; L. cruor, blood.] A green respiratory pigment occurring in blood plasma of certain worms.
- chlorofucin (klōröfū'sin) n. [Gk. chloros, green; L. fucus, seaweed.] Chlorophyll c, in diatoms and brown algae; chlorophyll γ.
- chlorophane (klō'rofān) n. [Gk. chloros, grass green; phainein, to appear.] A green chromophane.
- chlorophore (klō'röfōr) n. [Gk. chloros, grass green; phora, carrying.] A chlorophyll granule in Protista.
- chlorophyll (klō'röfil) n. [Gk. chloros, grass green; phyllon, leaf.] The green colouring matter found in plants and in some animals; chlorophyll a, $C_{55}H_{72}O_5N_4Mg$; b, $C_{55}H_{70}O_6N_4Mg$; c, or γ or chlorofucin, q.v.; c formerly appl. a mixture of chlorophyll a and pheophytin a.
- chloroplast (klō'röpläst), chloroplastid (klō'röpläs'tid) n. [Gk. chloros, grass green; plastos, moulded.] A minute granule or plastid containing chlorophylls a and b, found in plant-cells exposed to light.
- chloroplast pigments, chlorophylls, carotene, and xanthophyll.
- chlorosis (klörö'sis) n. [Gk. chloros, pallid.] Abnormal condition characterised by absence of green pigments in plants, owing to lack of light, or to magnesium- or iron-deficiency; green-sickness in humans.
- chlorostatolith (klō'röstāt'ölīth) n.
 [Gk. chloros, grass green; statos, stationary; lithos, stone.] A chloroplast which moves under the influence of gravity in a statocyte; cf. statolith, amylostatolith.
- chlorotic (klōrŏt'ĭk) a. [Gk. chloros, pallid.] Pert. or affected by chlorosis.
- choana (kō'ănă) n. [Gk. choane, funnel.] A funnel-shaped opening; posterior naris.

- choanocyte (kō'ānösīt) n. [Gk. choane, funnel; kytos, hollow.] A cell with funnel-shaped rim or collar round the base of a flagellum.
- choanoid (kō'ănoid) a. [Gk. choane, funnel; eidos, like.] Funnel-shaped; appl. eye muscle, retractor bulbi, absent in snakes, birds, and higher primates.

choanosome (kō'anösōm) n. [Gk. choane, funnel; soma, body.] In sponges, the inner layer with flagellate cells.

cholecystokinin (köl'ësis'tokin'in) n. [Gk. chole, bile; kystis, bladder; kinein, to move.] A duodenal hormone which induces contraction of gall-bladder and relaxation of

Oddi's sphincter.

choledoch (kŏl'ĕdŏk) a. [Gk. chole, bile; dochos, containing.] Appl. common bile duct.

choleic (kölē'ik) a. [Gk. chole, bile.]

Pert. acid contained in ox bile.

- cholesterol (kölës'tërŏl) n. [Gk. chole, bile; stereos, solid.] Cholesterin, a white fatty alcohol found in protoplasm, nerve tissue, bile, yolk, and other animal substances; C₂₇H₄₆O.
- choline (kď/lĭn, -ēn) n. [Gk. chole, bile.] A crystalline base found in plants and animals, a decomposition product of lecithin; C₈H₁₈O₂N.
- cholinergic (köliner'jik) a. [Gk. chole, bile; ergon, work.] Appl. parasympathetic nerve fibres which liberate acetyl-choline from their terminations; cf. adrenergic.

cholochrome (kŏl'ökrōm) n. [Gk. chole, bile; chroma, colour.] A bile

pigment; biliphaein.

cholohaematin (kŏl'öhĕ'mătin) n. [Gk. chole, bile; haima, blood.] Phylloerythrin; cholehaematin.

- chomophyte (kō'möfīt) n. [Gk. choma, mound; phyton, plant.] A plant growing in detritus on rocks.
- chondral (kôn'drăl) n. [Gk. chondros, cartilage.] Pert. cartilage.
- chondrification (kôn'drĭfīkā'shūn) n. [Gk. chondros, cartilage; L. facere,

to make.] Conversion into cartilage.

chondrigen (kôn'drijěn) n. [Gk. chondros, cartilage; gennaein, to produce.] The base matrix of all cartilaginous substance, a collagen.

chondrin (kôn'drin) n. [Gk. chondros, cartilage.] A gelatinous substance obtained from cartilage.

- chondriocont (kôn'driökŏnt) n. [Gk. chondros, grain; kontos, pole.] A rod-like or fibrillar type of chondriosome.
- chondriodieresis (kôn'driödiĕr'ësïs)
 n. [Gk. chondros, grain; dieressein, to swing about.] Changes in mitochondria during cell division.
- chondriokinesis (kôn'driökinē'sīs) n. [Gk. chondros, grain; kinesis, movement.] Division of chondriosomes in mitosis and meiosis.
- chondrioma (kôndriō'mă) n. [Gk. chondros, grain.] The chondriosome content of a cell; chondriome.
- chondriomere (kôn'driömēr) n. [Gk. chondros, grain; meros, part.] Plastomere, q.v.; cytomere, q.v.

chondriomite (kôn'driömīt) n. [Gk. chondros, grain; mitos, thread.] A linear type of chondriosome.

chondrioplast (kôn'driöpläst) n. [Gk. chondros, grain; plastos, formed.] A rod-like formation of reticular

material; Golgi rod.

chondriosomes (kôn'driösōmz) n. plu. [Gk. chondros, grain; soma, body.] Mitochondria, q.v.; numerous synonyms: e.g. chondriomites, chondrioconts, chondriospheres, chondrioplasts.

chondriosphere (kôn'driösfēr) n. [Gk. chondros, grain; sphaira, globe.] A spherical type of chondriosome; mitochondria which have

coalesced.

chondroblast (kôn'dröblăst) n. [Gk. chondros, cartilage; blastos, bud.] A cartilage-producing cell.

chondroclast (kôn'drökläst) n. [Gk. chondros, cartilage; klastos, broken down.] A large multinucleate cell which destroys cartilage matrix; also chondrioclast.

- chondrocranium (kôn'drökrā'nĭŭm)
 n. [Gk. chondros, cartilage;
 kranion, skull.] The skull when in a cartilaginous condition, either temporarily as in embryos, or permanently as in some fishes.
- chondrocyte (kôn'drösīt) n. [Gk. chondros, cartilage; kytos, hollow.] A cartilage cell.
- chondrogenesis (kôn'dröjĕn'ësis) n. [Gk. chondros, cartilage; genesis, descent.] The production or formation of cartilage.
- chondroid (kôn'droid) a. [Gk. chondros, cartilage; eidos, shape.] Cartilage-like; appl. tissue, undeveloped cartilage or pseudocartilage serving as support in certain invertebrates and lower vertebrates; appl. vesicular supporting tissue of notochord; fibrohyaline.
- chondromucoid (kôn'drömū'koid) n. [Gk. chondros, cartilage; L. mucus, mucus; Gk. eidos, form.] A basophil protein which with collagen forms ground-substance of cartilage; chondromucin.
- chondrophore (kôn'dröfōr) n. [Gk. chondros, cartilage; pherein, to bear.] A structure which supports the inner hinge cartilage in a bivalve shell.
- chondroskeleton (kôn'dröskěl'ětŏn)
 n. [Gk. chondros, cartilage; skeleton, dried body.] A cartilaginous skeleton.
- chondrosteous (kôndrŏs'tëus) a. [Gk. chondros, cartilage; osteon, bone.] Having a cartilaginous skeleton.
- chondrosternal (kôn'dröster'năl) a. [Gk. chondros, cartilage; sternon, breast.] Pert. rib cartilages and sternum.
- chone (kō'nē) n. [Gk. chone, funnel.] A passage through cortex of sponges, with one or more external openings, and one internal opening.
- chorda (kôr'dă) n. [Gk. chorde, string.] The notochord; in plural, tendinous cords attached to valves of heart; any cord-like structure.

chordacentra (kôr'dăsĕn'tră) n. plu. [Gk. chorde, string; L. centrum, centre.] Centra formed by conversion of chordal sheath into a number of rings; cf. archentra.

chordate (kôr'dāt) a. [Gk. chorde, string.] Having a notochord.

- chordotonal (kôr'dötō'năl) a. [Gk. chorde, string; tonos, tone.] Appl. rod-like or bristle-like receptors for mechanical and sound vibrations, in various parts of body of insects.
- chore (kō'rē) n. [Gk. chore, place.] An area manifesting a unity of geographical or environmental conditions; cf. biochore, biotope.
- choreiathetose (kŏrīāth'ētōs) a. [Gk. choreia, dance; athetos, lawless.] Arhythmic and uncoordinated; appl. foetal movements.
- choriocapillaris (kör'iökäpil'äris, -käp'ilä'ris) n. [Gk. chorion, skin; L. capillaris, capillary.] The innermost vascular layer of choroid.

chorioid,—choroid, q.v.

- chorion (kő'riön) n. [Gk. chorion, skin.] An embryonic membrane external to and enclosing the amnion; allantochorion, q.v.; a hardened shell covering egg of insects; outer membrane of seed.
- chorion frondosum,—villous placental part of chorion.
- chorion laeve, smooth nonplacental part of chorion.
- chorionic (korion'ik) a. [Gk. chorion, skin.] Pert. the chorion; appl. gonadotrophic hormone or prolan.
- chorioretinal (kör'iörĕt'inăl) a. [Gk. chorion, skin; L. retina, retina.]
 Pert. choroid and retina.
- choripetalous (kō'rīpēt'ālūs) a. [Gk. choris, separate; petalon, leaf.]
 Having separate petals.
- choriphyllous (kō'rīfīl'ŭs) a. [Gk. choris, separate; phyllon, leaf.] Having perianth parts distinct.
- chorisepalous (kō'risĕp'ālūs) a. [Gk. choris, separate; F. sépale, sepal.] Having the sepals separate.

- chorisis (kō'rĭsīs) n. [Gk. choris, separate.] Increase in parts of floral whorl due to division of its primary members; deduplication.
- choroid (kōr'oid) a. [Gk. chorion, skin; eidos, form.] Appl. delicate and highly vascular membranes. n. Layer of eye between retina and sclera.
- choroidal (köroid'ăl) a. [Gk. chorion, skin; eidos, form.] Pert. choroid.
- chorology (köröl'öji) n. [Gk. choros, place; logos, discourse.] Biogeography; geographical distribution; biotopography; science of the distribution of organisms or of organs.
- choronomic (kōrönŏm'ĭk) a. [Gk. choros, place; nomos, law.] External, appl. influences of geographical or regional environment, opp. autonomic.
- chorotypes (kō'rötīps) n. plu. [Gk. choros, place; typos, pattern.] Local types.
- chresard (krēsârd') n. [Gk. chresis, use; ardo, I water.] Soil water available for plant growth; cf. echard, holard.
- chromaffin (krō'măfin) a. [Gk. chroma, colour: L. affinis, related.] Chromaphil.
- chromaphil (krō'māfil) a. [Gk. chroma, colour; philein, to love.] Stained by chromic acid or its salts when adrenaline is present; appl. cells forming medullary parts of suprarenal bodies; appl. bodies or paraganglia; chromophil, chromaffin.
- chromaphobe (krō'măfōb) a. [Gk. chroma, colour; phobos, fear.] Appl. non-stainable cells or tissues; chromophobe.
- chromasie (krō'māsī) n. [Gk. chroma, colour.] Increase of chromatin in nucleus and formation of nucleolus; cf. achromasie.
- chromatic (krōmăt'îk) a. [Gk. chroma, colour.] Colourable by means of staining reagents; pert. colour; having chromatophores.

- chromatic sphere, the sphere formed by coalescence of chromosomes after anaphase in mitosis.
- chromatic threshold,—the minimal stimulus, varying with wave length of light, which induces a colour sensation.
- chromaticity (krōmătis'īti) n. [Gk. chroma, colour.] Unlikeness to grey, or saturation of a colour.
- chromatid (krō'mătĭd) n. [Gk. chroma, colour.] A component of a tetrad in meiosis; a half-chromosome between early prophase and metaphase in mitosis, or between diplotene and second metaphase in meiosis.
- chromatin (krō'mătin) n. [Gk. chroma, colour.] A substance in the nucleus which contains nucleic acid proteids, and stains with basic dyes.
- chromatogen organ,—a brownish lobed body, the axial organ of certain echinoderms.
- chromatoid grains,—grains in cellprotoplasm, which stain similarly to chromatin.
- chromatolysis (krō'mătôl'īsīs) n.
 [Gk. chroma, colour; lysis, loosing.]
 Disintegration of Nissl granules, as in fatigued nerve-cells; tigrolysis.
- chromatophil (krō'mătöfil) a. [Gk. chroma, colour; philein, to love.] Staining easily.
- chromatophore (krō'mătöfōr) n.
 [Gk. chroma, colour; pherein, to bear.] A coloured plastid of plants and animals; a colourless body in cytoplasm and developing into a leucoplast, chloroplast, or chromoplast; a pigment cell, or group of cells, which under control of the sympathetic nervous system can be altered in shape to produce a colour change.
- chromatophorotropic (krō'mātöfō' rötrōp'ik) a. [Gk. chroma, colour; pherein, to bear; trope, turn.] Appl. a hormone, intermedin, secreted by pars intermedia of pituitary and causing expansion of chromatophores; appl. hormone of crustacean eye-stalk.

- chromatophyll (krē'mătöfil) n. [Gk. chroma, colour; phyllon, leaf.] The colouring matter of plant-like flagellates; also chromophyll.
- chromatoplasm (krō'mătöplăzm) n. [Gk. chroma, colour; plasma, mould.] The colour or pigment matter in cells.

chromatosome,-chromosome.

chromatospherite (krō'mătösfēr'īt)
n. [Gk. chroma, colour; sphaira, globe.] A nucleolus, q.v.

chromidia (krōmĭd'iā) n. plu. [Gk. chroma, colour; idion, dim.] Extra-nuclear particles of chromatin, which may replace or be re-formed into nuclei.

chromidial substance, — minute basophil granules containing iron, occurring in cytoplasm as chromophil or tigroid bodies.

chromidiogamy (krōmid'iŏg'āmi) n. [Gk. chroma, colour; idion, dim.; gamos, marriage.] The union of chromidia from two conjugants.

chromidiosomes (krōmid'iosomz) n.
plu. [Gk. chroma, colour; idion,
dim.; soma, body.] The smallest
chromatin particles of which the
chromidial mass is composed.

chromiole (krō'miōl) n. [Ġk. chroma, colour.] One of the minute granules of which a chromomere is composed.

chromo-argentaffin (krö'möârjĕn'tăfīn) a. [Gk. chroma, colour; L.
argentum, silver; affinis, related.]
Staining with bichromates and silver
nitrate; appl. flask-shaped cells in
epithelium of crypts of Lieberkühn.

chromoblast (krō'möblăst) n. [Gk. chroma, colour; blastos, bud.] An embryonic cell giving rise to a

pigment cell.

chromocentre (krōmösĕn'tër) n. [Gk. chroma, colour; kentron, centre.] The fused heterochromatic region around centromeres; fused prochromosomes.

chromocyte (krō'mösīt) n. [Gk. chroma, colour; kytos, hollow.]
Any pigmented cell.

chromogen (krō'möjĕn) n. [Gk. chroma, colour; genos, birth.] The substance which is converted into a

pigment, e.g. by oxidation; a chromogenic organism.

chromogenic (krō'möjěn'ik) a. [Gk. chroma, colour; genos, birth.] Colour - producing; appl. organisms, as bacteria.

chromolipides (krō'mölīp'fdz) n. plu. [Gk. chroma, colour; lipos, fat.] The carotenoids and related pigments.

- chromomere (krō'mömēr) n. [Gk. chroma, colour; meros, part.] One of the chromatin granules of which a chromosome is formed, and which corresponds to an id or a gene; granular part of blood platelet, opp. hyalomere.
- chromonema (krō'mönē'mā) n. [Gk. chroma, colour; nema, thread.] A coiled or convoluted thread in prophase of mitosis; central thread in chromosome. Plu. chromonemata.
- chromophanes (krō'möfanz) n. plu. [Gk. chroma, colour; phainein, to show.] Red, yellow, and green oil globules found in retina of birds, reptiles, fishes, marsupials; any retinal pigments.
- chromophil (krō'möfil) a. [Gk. chroma, colour; philein, to love.] Chromaphil, chromaffin, q.v.; chromophilic.

chromophilous (krömŏf'ilūs) a. [Gk. chroma, colour; philos, loving.]
Staining readily.

chromophobe (krō'möfōb) a. [Gk. chroma, colour; phobos, fear.] Non-stainable or staining slightly; appl. certain cells of pituitary gland; chromaphobe.

chromophore (krō'möfōr) n. [Gk. chroma, colour; pherein, to bear.] Any substance to whose presence colour in a compound is due.

chromophyll,—chromatophyll, q.v. chromoplast (krö'möpläst) n. [Gk. chroma, colour; plastos, moulded.] A coloured plastid or pigment body; coloured plastid other than a chloroplast; chromoplastid.

chromoproteins (krō'möprō'teinz) n.
plu. [Gk. chroma, colour; protos,
first.] Substances formed by combination of a protein with a pigment
or chromophore.

- chromosome (krō'mösōm) n. [Gk. chroma, colour; soma, body.] One of deeply staining bodies, the number of which is constant for the cells of a species, into which the chromatin resolves itself during karyokinesis and meiosis.
- chromosome-races,—races differing in number of chromosomes or of chromosome sets.
- chromosomin (krōmösō'min) n. [Gk. chroma, colour; soma, body.] One of the protein constituents of chromosomes.
- chromospire (krō'möspīr) n. [Gk. chroma, colour; speira, coil.] A spireme-like thread formed from nuclear granules in haplomitosis.
- chronaxie, chronaxy (krō'nāksī) n. [Gk. chronos, time; axia, value.] Latent period between electrical stimulus and muscular response; minimal excitation time required with a current of an intensity twice the threshold necessary for excitation when the duration of the stimulus is prolonged; chronaxia.
- chrysalis (kris'ălis) n. [Gk. chrysallis, gold, golden thing.] Pupa stage of certain insects.
- chrysocarpous (kris'okâr'pŭs) a.
 [Gk. chrysos, gold; karpos, fruit.]
 With golden fruit.
- chrysophanic (krisöfán'ík) a. [Gk. chrysos, gold; phainein, to show.] Having a golden or bright orange colour, appl. an acid formed in certain lichens.
- chrysophyll (kris'öfil) n. [Gk. chrysos, gold; phyllon, leaf.] A yellow colouring matter in plants, a decomposition product of chlorophyll.
- chylaceous (kīlā'sēus) a. [Gk. chylos, juice.] Of the nature of chyle.
- chyle (kil) n. [Gk. chylos, juice.] Lymph containing globules of emulsified fat, found in the lacteals during digestion.
- chylifaction (ki'lifăk'shun) n. [Gk. chylos, juice; L. facere, to make.] Formation of chyle; also chylification; chylopoiesis.
- chyliferous (kilif'ërus) a. [Gk. chylos, juice; L. ferre, to carry.]

- Chyle-conducting; *appl.* tubes or vessels; chylophoric.
- chylific (kilif'ik) a. [Gk. chylos, juice; L. facere, to make.] Chyleproducing; appl. ventricle or true stomach of insects.
- chylocaulous (kīlöcôl'ŭs) a. [Gk. chylos, juice; kaulos, stem.]
 With fleshy stems.
- chylocyst (kī'lösīst) n. [Gk. chylos, juice; kystis, bladder.] The chyle receptacle.
- chylomicrons (kīlömī'krönz) n. plu. [Gk. chylos, juice; mikros, small.] Minute fatty particles in plasma, plentiful during fat digestion.
- **chylophoric**,—chyliferous.
- chylophyllous (ki'lofil'ŭs) a. [Gk. chylos, juice; phyllon, leaf.] With fleshy leaves; appl. certain desert plants.
- chylopoiesis (kī'löpoiē'sīs) n. [Gk. chylos, juice; poiein, to produce.]
 The production of chyle.
- chyme (kim) n. [Gk. chymos, juice.] The partially digested food after leaving the stomach.
- chymification (kī'mĭfikā'shŭn) n. [Gk. chymos, juice; L. facere, to make.] The process of converting food into chyme.
- chymosin,--rennin, q.v.
- chytridium (kitrid'ium) n. [Gk. chytridion, little pot.] The spore vessel of certain fungi.
- cibarium (sǐbā'rĭŭm) n. [L. cibaria, victuals.] The part of the buccal cavity anterior to pharynx, in insects.
- cicatricial tissue, newly formed fibrillar connective tissue which closes and draws together wounds.
- cicatricle (sīkāt'rīkl), cicatricula (sīk'ātrīk'ūlā), cicatrix (sīk'ātrīks, sīkā'trīks) n. [L. cicatrix, scar.] The blastoderm in bird and reptile eggs; a small scar in place of previous attachment of an organ; a scar; the mark left after healing of a wound in plants.
- cicinnal (sīs'īnāl) a. [Gk. kikinnos, curled lock.] Appl. uniparous cymose branching in which daughter axes are developed right and left alternately; cincinnal.

- cilia (sil'ía) n. plu. [L. cilium, eyelid.] Hairlike vibratile outgrowths of ectoderm, or processes, of many cells; barbicels of a feather; eyelashes.
- ciliaris (sĭlĭa'rĭs) n. [L. cilium, eyelid.] Unstriped muscle forming a ring outside anterior part of choroid and, attached to ciliary processes, acting on convexity of lens.

ciliary (sĭl'íărī) a. [L. cilium, eyelid.]

Pert. cilia; appl certain structures in the eveball.

ciliate (sĭl'iāt) a. [L. cilium, eyelid.] Provided with cilia; ciliated.

ciliated epithelium,—an epithelium found lining various passages, usually with columnar cells provided with cilia on the free surface.

ciliograde (sĭl'iögrād) a. [L. cilium, eyelid; gradus, step.] Progressing by movement of cilia.

ciliospore (sīl'iöspōr) n. [L. cilium, eyelid; Gk. sporos, seed.] A ciliated protozoan swarm-spore.

cilium (sil'ium) n. [L. cilium, eyelid.] Sing. of cilia, q.v.

cinchonine (sin'könin) n. [After Countess de Chinchon.] Alkaloid found in various Rubiaceae.

cincinnus (sĭnsĭn'ŭs) n. [L. cincinnus, curl.] A scorpioid cyme.

cinclides (sing'klidez) n. plu. [Gk. kingklis, latticed gate.] Perforations, in body wall of certain Anthozoa, for extrusion of acontia. Sing. cinclis.

cinerea (sinë'rea) n. [L. cinereus, ashen.] The grey matter of the

nervous system.

cingulate (sing'gūlāt) a. [L. cingulum, girdle.] Having a girdle or cingulum; shaped like a girdle; appl. a gyrus and sulcus above corpus callosum.

cingulum (sīng'gūlūm) n. [L. cingulum, girdle.] Any structure which is like a girdle; part of plant between root and stem; part of diatom frustule uniting valves; a ridge round base of crown of a tooth; a tract of fibres connecting callosal and hippocampal convolutions of brain; outer ciliary zone on disc of rotifers; clitellum, q.v.

circinate (sĭr'sĭnāt) a. [L. circinus, compasses.] Rolled on the axis, so that apex is centre.

circulation (sër'kūlā'shūn) n. [L. circulatio, act of circulating.] The regular movement of any fluid within definite channels in the body; streaming movement of protoplasm of plant cells.

circulus (sīr'kūlūs) n. [L. circulus, circle.] Any ringlike arrangement, as of blood-vessels caused by branching or connection with one another, as circulus major of iris, or as of markings of fish scales.

circulus arteriosus,—a vascular ring at base of brain; circle of Willis.

circumduction (ser'kumduk'shun) n.
[L. circum, around; ductus, led.]
The form of motion exhibited
by a bone describing a conical
space with the articular cavity as
apex.

circumferential (sër'kumfërën'shăl)

a. [L. circum, around; ferre, to
bear.] Appl. cartilages which surround certain articulatory fossae;
appl. primary lamellae parallel to
circumference of bone.

circumfila (sërkŭm'filă) n. plu. [L. circum, around; filum, thread.]
Looped or wreathed filaments on antennal segments, as in gallmidges.

circumflex (sër'kŭmflěks) a. [L. circum, around; flectere, to bend.]
Bending round; appl. certain arteries, veins; appl. nerve, the axillary nerve.

circumfluence (sërkum'flooëns) n.

[L circum, around; fluens, flowing.] In Protozoa, ingestion by protoplasm flowing towards food and surrounding it after contact; cf. circumvallation.

circumgenital (sër'kŭmjën'îtăl) a. [L. circum, around; gignere, to beget.] Surrounding the genital pore; appl. glands secreting waxy powder in oviparous species of Coccidae.

- circumnutation (sër'kumnuta'shun)
 n. [L. circum, around; nutare, to nod.] The irregular elliptical or spiral movement exhibited by apex of a growing stem, shoot or tendril.
- circumoesophageal (sër'kŭmēsŏfăj'ēăl) a. [L. circum, around; Gk. oisophagos, gullet.] Appl. structures or organs surrounding or passing along the gullet.

circumpolar (sër'kŭmpō'lăr) a. [L. circum, around; polus, end of axle.] Appl. flora and fauna of Polar regions.

circumpulpar (sër'kumpul'păr) a. [L. circum, around; pulpa, fruit-pulp.] Appl. dentine forming layer around pulp cavity of teeth, as in fishes.

circumscissile (sër'kūmsīs'īl) a. [L. circum, around; scindere, to cut.]
Splitting along a circular line; appl. dehiscence exhibited by a pyxidium.

circumscript (sër'kumskript) a. [L. circumscribere, to draw line around.] Appl. marginal sphincter when sharply defined, in sea-anemones.

circumvallate (ser'kumvăl'āt) a. [L. circum, around; vallum, rampart.] Encircled by a wall, as of tissue; appl. certain tongue papillae.

circumvallation (sër'kŭmvălā'shŭn)
n. [L. circum, around; vallare, to wall.] Ingestion of food by extruded pseudopodia, as in protozoa or in phagocytes.

circumvascular (sër'kŭmvăs'kūlăr) a. [L. circum, around; vasculum, small vessel.] Appl. dentine lining vascular canals in pulp cavity of teeth, as in fishes.

cirrate (sĭr'āt) a. [L. cirratus, having curls.] Having cirri.

cirrhus and cirrhi, -- see cirrus.

cirrose (sĭr'ōs, sĭrōs') a. [L. cirrus, curl.] With cirri or tendrils.

cirrus (sīr'ūs) n., cirri (sīr'ī) plu.

[L. cirrus, curl.] Tendrils; appendages of barnacles; jointed filaments of axis or of aboral surface of crinoids; barbels of fishes; respiratory and tactile ap-

pendages of worms; organs of copulation in some molluscs and trematodes; hairlike structures on appendages of insects.

cisterna (sister'nă) n. [L. cisterna, cistern.] Closed space containing fluid, as any of the subarachnoid

spaces.

citrin (sĭt'rīn) n. [L.L. citrus, lemon.]

A factor in lemon juice which regulates capillary permeability; vitamin P.

citrulline (sitrul'in) n. [L. citrullus, water-melon.] An amino acid first obtained from water-melon, also occurring as intermediate product in formation of urea from ornithine; C₆H₁₈O₃N₃.

cladanthous (klădăn'thŭs) a. [Gk. klados, sprout; anthos, flower.] Having terminal archegonia on short lateral branches; cladocarpous.

cladautoicous (klad'ôtoik'ŭs) a. [Gk. klados, sprout; autos, self; oikos, house.] With antheridia on a special stalk as in mosses.

cladocarpous,—cladanthous, q.v.

cladode (klăd'ōd) n. [Gk. klados, sprout.] Branch arising from axil of leaf, or green flattened stem, resembling a foliage leaf; cladophyll; phylloclade.

cladodont (klad'ödönt) a. [Gk. klados, sprout; odous, tooth.] Having or appl. teeth with prominent central and small lateral cusps.

cladogenesis (klădöjěn'ësĭs) n. [Gk. klados, sprout; genesis, descent.] Branching of evolutionary lineages so as to produce new types.

cladogenous (klădŏj'ēnūs) a. [Gk. klados, sprout; gennaein, to produce.] Stem-borne; appl. certain roots; cladanthous, q.v.

cladome (klådōm') n. [Gk. klados, sprout.] The group of superficially situated rays in a triaene.

cladophyll (klad'öfil) n. [Gk. klados, sprout; phyllon, leaf.] Cladode.

cladoptosis (klåd'öptö'sis) n. [Gk. klados, sprout; ptosis, falling.]
Annual or other shedding of twigs.
cladose (klåd'ōs) a. [Gk. klados, sprout.] Branched.

- cladosiphonic (klăd'ösīfŏn'ĭk) a. [Gk. klados, sprout; siphon, tube.] With insertion of leaf-trace on periphery of the axial stele; opp. phyllosiphonic.
- cladotyle (kläd'ötil) n. [Gk. klados, sprout; tylos, knob.] A rhabdus with one actine branched, the other tylote.
- clamp-connections, swellings on certain dikaryotic hyphae, for passage of daughter nuclei to cell below, with subsequent septum formation; also occurring in whorls, for distribution of nuclei to hyphal branches.
- clandestine (klandes'tin) a. [L. clandestinus, from clam, secretly.]
 Appl. evolution which is not apparent in adult forms; or of adult characters from ancestral embryonic characters.
- clasmatocyte (klăz'măt'ösīt, klăsmăt'ösīt) n. [Gk. klasma, fragment; kytos, hollow.] A variable basiphil phagocyte or macrophage in areolar tissue; a histiocyte.
- claspers (klâs'përz) n. plu. [M.E. claspen, to hold.] Rod-like processes on pelvic fins of certain male elasmobranchs; outer gonapophyses of insects; valves or harpes of male Lepidoptera; any modification of an organ or part to enable the two sexes to clasp one another; tendrils or climbing shoots.
- claspettes,-harpagones, q.v.
- class (klâs) n. [L. classis, division.] A division of a phylum and divided into orders, in classification of plants or animals.
- clathrate (klăth'rāt) a. [Gk. klethra, lattice.] Lattice-like.
- claustrum (klôs'trum) n. [L. claustrum, bar.] In cerebral hemispheres, a thin layer of grey substance lateral to external capsule; one of the Weberian ossicles in Cyprinidae and Characinidae.
- clava (klā'vă) n. [L. clava, club.] The knob-like end of antenna of certain insects; swelling at end of fasciculus gracilis of medulla oblongata.
- clavate (klā'vāt) a. [L. clava, club.] Club-shaped; thickened at one end.

- clavicle (klăv'ikl) n. [L. clavicula, small key.] Collar-bone, forming anterior or ventral portion of shoulder girdle.
- clavicular (klăvĭk'ūlăr) a. [L. clavicula, small key.] Pert. clavicle. clavicularium (klăvik'ūlā'rīŭm) n. [L. clavicula, small key.] The epiplastron of Chelonia, probably corresponding to clavicles of other
- clavola (klăvō'lă) n. [L. clava, club.] The flagellar portion, or terminal joints, of insect antenna.
- clavula (kläv'ūlă) n. [L. clava, club.] A monactinal modification of triaxon spicule; a minute ciliated spine on fasciole of Spatangidae; a clavate sporophore of certain fungi.
- clavus (klā'vus) n. [L. clavus, nail.]
 The part of an hemelytron lying next scutellum in Hemiptera; a projection or crotchet from scape of spiders; ergot disease in grasses.
- claw (klô) n. [A.S. clawu, claw.] The unguis or stalk of a petal; a sharp curved nail on finger or toe; forceps of certain crustaceans; curved process on limb of insect.
- clearing foot,—filamentous process of exopodite of second maxilla in Phyllocarida.
- cleavage (klē'vēj) n. [A.S. cleofan, to cut.] The series of karyokinetic divisions which change the egg into a multicellular embryo.
- cleavage nucleus,—nucleus of fertilised egg or zygote produced by union of male and female pronuclei; the egg-nucleus of parthenogenetic eggs.
- cleidoic (klīdō'ik) a. [Gk. kleis, bar; oon, egg.] Having or pert. eggs enclosed within a shell or membrane.
- cleistocarp (klī'stökârp) n. [Gk. kleistos, closed; karpos, fruit.] Cleistothecium, q.v.
- cleistocarpous (klī'stökâr'pŭs) a. [Gk. kleistos, closed; karpos, fruit.] Appl. mosses with non-operculate capsules; cleistocarpic.

- cleistogamic (klīstögăm'īk) a. [Gk. kleistos, closed; gamos, marriage.] *Pert.* or possessed of characteristics of cleistogamy; cleistogamous.
- cleistogamy (klistog'ami) n. kleistos, closed; gamos, marriage.] State of having small inconspicuous self-fertilising flowers; fertilisation without opening of florets, opp. chasmogamy.
- cleistogene (klī'stöjēn) n. kleistos, closed; genos, descent.] A plant with cleistogamous flowers. cleistothecium (klīstöthē'sĭŭm) n. [Gk. kleistos, closed; theke, box.] An ascocarp which remains closed
- cleithrum (klī'thrūm) n. [Gk. kleithron, bar. The pair of additional clavicles in Stegocephalia; clavicular element of some fishes.

and produces its spores internally.

climacteric (klimakter'ik) n. klimakter, step of staircase.] critical phase, or period of change, in living organisms; appl. change associated with menopause; appl. phase of increased respiratory activity at ripening of fruit.

climatype (klī'mătīp) n. [Gk. klima, climate; typos, image.] A biotype resulting from selection in a particular climate; climatic ecotype.

climax (klī'māks) n. [Gk. klimax, ladder.] The mature or stabilised stage in a successional series of dominant communities, when species are completely adapted to environmental conditions; completion of development, appl. leaves.

clinandrium (klinăn'drium) n. [Gk. kline, bed; aner, man.] A cavity in the column between anthers in orchids.

cline (klīn) n. [Gk. klinein, to slant.] A series of form changes; gradient of biotypes; character-gradient.

clinging fibres,—tendril fibres, q.v. clinidium (klinid'iŭm) n. [Gk. klinidion, small couch.] A filament in a pycnidium, which produces spores.

clinoid (klī'noid) a. [Gk. kline, couch; eidos, form.] Appl. pro-

cesses of sella turcica.

clinosporangium (klī'nöspöran'jĭŭm) n. [Gk. kline, bed; spora, seed; anggeion, vessel.] Pycnidium, q.v.

clitellum (klitěl'um) n. [L. clitellae, pack-saddle.] The saddle or swollen glandular portion of skin of certain annelid worms.

clitoris (klī'tōris) n. [Gk. kleiein, to enclose.] An erectile organ, homologous with penis, at upper part of

clivus (klī'vŭs) n. [L. clivus, slope.] A shallow depression in sphenoid, behind dorsum sellae; posterior sloped part of the monticulus.

cloaca (kloā'kă) n. [L. cloaca, sewer.] The common chamber into which intestinal, genital, and urinary canals open, in vertebrates except most mammals.

clone (klon) n. [Gk. klon, twig.] An individual produced asexually; group of individuals propagated by mitosis from a single ancestor; an apomict strain.

clonus (klon'us) n. [Gk. klonos, violent motion.] A series of muscular contractions when individual contractions are discernible: incomplete tetanus.

cluster-crystals, - globular aggregates of calcium oxalate crystals in plant cells; sphaeraphides.

cluster-cup,—aecidium, q.v. clypeal (klip'eal) a. [L. clypeus,

shield.] *Pert.* clypeus of insects. clypeate (klip'eāt) a. [L. clypeus, Round or buckler-like; clypeiform; having a clypeus.

clypeola (klĭpē'ölă), clypeole (klĭp'-[L. clypeus, shield.] ëōl) n. sporophyll in the spike of an Equisetum.

clypeo-labral (klip'eölåb'rål) a. [L. clypeus, shield; labrum, lip.] Appl. suture between clypeus and labrum.

clypeus (klip'eus) n. [L. clypeus, shield.] A sclerite on anteromedian part of insect head; a band of tissue round mouth of perithecium of certain fungi.

cnemial (knē'mial, nē'mial) a. [Gk. knemis, legging.] Pert. tibia; appl. ridge along dorsal margin of tibia. cnemidium (knēmid'iŭm, nēmid'iŭm) n. [Gk. knemis, legging; idion, dim.] Lower part of bird's leg devoid of feathers, generally scaly.

cnemis (knē'mis, nē'mis) n. [Gk. knemis, legging.] Shin or tibia.

cnida (knī'dā, nī'dā) n. [Gk. knide, nettle.] A cnidoblast; a nematocyst. cnidoblast (knī'döblāst, nī'döblāst) n.

[Gk. knide, nettle; blastos, bud.] Stinging cell of Coelentera.

Stinging cell of Coelentera.

cnidocil (knī'dösīl, nī'dösīl) n. [Gk. knide, nettle; L. cilium, eyelid.] A minute process projecting externally from a cnidoblast.

cnidophore (knī'döfōr, nī'döfōr) n. [Gk. knide, nettle; pherein, to bear.] A modified zooid which

bears nematocysts.

cnidopod (knī'döpŏd, nī'döpŏd) n. [Gk. knide, nettle; pous, foot.] Drawn-out basal part of a nematocyst, embedded in mesogloea.

cnidosac (knī'dösāk, nī'dösāk) n. [Gk. knide, nettle; sakkos, bag.] A kidney-shaped swelling or battery, often protected by a hood, found on dactylozooids of Siphonophora.

coaction (kōăk'shŭn) n. [L. cum, with; actio, action.] The reciprocal activity of organisms within a community.

coadaptation (kō'ădăptā'shŭn)

[L. cum, with; ad, to; aptare, to fit.] The correlated variation in two mutually dependent organs.

coagulation (köägülä'shūn) n. [I... cum, with; agere, to drive.] Curdling or clotting; the changing from a liquid to a viscous or solid state by chemical reaction; appl. vitamin K, the antihaemorrhagic accessory food factor.

- coagulin (kōāg'ūlĭn) n. [L. coagulum, rennet.] Any agent capable of coagulating albuminous substances.
- coagulum (koăg'ūlūm) n. [L. coagulum, rennet.] Any coagulated mass; clot; curd.
- coarctate (kōârk'tāt) a. [L. coarc-tare, to press together.] Com-

pressed; closely connected; with abdomen separated from thorax by a constriction.

coarctate larva or pupa,—semipupa; pseudopupa; a larval stage

of certain Diptera.

cocci (kök'si) n. plu. [Gk. kokkos, berry.] Septicidal carpels; spore mother cells of certain hepatics; rounded cells, as certain bacteria.

coccogone (kök'ögön) n. [Gk. kokkos, berry; gonos, birth.] A reproductive cell in certain algae.

coccoid (kŏk'oid) a. [Gk. kokkos, berry; eidos, form.] Like or pert. a coccus; spherical or globose.

coccolith (kŏk'ölĭth) n. [Gk. kokkos, berry; lithos, stone.] A calcareous spicule in certain Flagellata.

coccospheres (kŏk'ösfērz) n. plu. [Gk. kokkos, berry; sphaira, globe.] Remains of hard parts of certain algae and radiolarians.

coccus (kŏk'ŭs) n. [Gk. kokkos,

berry.] Sing. of cocci, q.v.

coccygeal (köksīj'eăl) a. [Gk. kokkyx, cuckoo.] Pert. or in region of coccyx.

- coccyx (kök'sĭks) n. [Gk. kokkyx, cuckoo.] The terminal part of the vertebral column beyond the sacrum.
- cochlea (kŏk'lëă) n. [Gk. kochlias, snail.] Anterior part of labyrinth of the ear, spirally coiled like a snail's shell
- cochlear (kŏk'lëăr) a. [Gk. kochlias, snail.] Appl. aestivation when wholly internal leaf is next but one to wholly external leaf; pert. the cochlea.
- cochleariform (kök'lear'ifôrm) a. [Gk. kochlias, snail; L. forma, shape.] Screw- or spoon-shaped; pert. thin plate or process of bone separating tensor tympani canal from Eustachian tube.
- cochleate (kŏk'lëāt) a. [Gk. kochlias, snail.] Screw-like; spiral.
- cocoon (kökoon') n. [F. cocon, cocoon.] The protective case of many larval forms before they become pupae; silky or other covering formed by many animals for their eggs.

- coelarium,—coelomic epithelium; mesothelium.
- coelenteron (sēlĕn'tërŏn) n. [Gk. koilos, hollow; enteron, intestine.]
 Cavity in body of Coelenterata.
- coeliac (sē'līāk) a. [Gk. koilia, belly.]

 Pert. the abdominal cavity;

 appl. arteries, veins, nerves,

 plexus.
- coeloblast (sē'loblāst) n. [Gk. koilos, hollow; blastos, bud.] A division of the embryonic hypoblast.
- coeloconic (sē'lökŏn'īk) a. [Gk. koilos, hollow; konos, cone.] Having, or consisting of, a conical process situated in a pit; appl. sensillae.
- coelogastrula (sē'lögăs'troolă) n. [Gk. koilos, hollow; gaster, stomach.] A gastrula developed from a blastula with a segmentation cavity.
- **coelom** (sē'löm) n. [Gk. koiloma. hollow.] Body cavity, q.v.
- coelomate (sē'lömāt sēlō'māt,) a. [Gk. koiloma, hollow.] Having a coelom.
- coelomesoblast (sēloměs'öblăst) n. [Gk. koilos, hollow; mesos, middle; blastos, bud.] In segmentation, the mesoblastic bands destined to form wall of coelom and outgrowths.

coelomic (sēlōm'ĭk) a. [Ğk. koiloma, hollow.] Pert. a coelom.

- coelomocytes (sēlō'mösīts) n. plu. [Gk. koiloma, hollow; kytos, hollow vessel.] Coelomic corpuscles, including amoebocytes and eleocytes, in annelids; mesenchymatous cells in body cavity of nematodes.
- coelomoduct (sēlō'mödŭkt) n. [Gk. koiloma, hollow; L. ducere, to lead.] A channel leading from body cavity to exterior.
- coelomopores (sēlō'möpōrz) n. plu. [Gk. koiloma, hollow; poros, passage.] Ducts leading directly from pericardial cavity to exterior, peculiar to Nautilus.
- coelomostome (sēlō'möstōm) n. [Gk. koiloma, hollow; stoma, buth.]
 The external opening of a celomoduct.
- coelosperm (sē'löspērm) n. [Gk.

- koilos, hollow; sperma, seed] A carpel, hollow on its inner surface.
- coelozoic (sē'lözō'ik) a. [Gk. koilos, hollow; zoon, animal.] Appl. a trophozoite when situated in some cavity of the body.
- coenanthium (sēnān'thium) n. [Gk. koinos, common; anthos, flower.] Inflorescence with a nearly flat receptacle having upcurved margins.
- coenenchyma (sēněng'kimă), coenenchyme (sēněng'kim) n. [Gk. koinos, common; engchyma, infusion.] Common tissue which connects the polyps or zooids of a compound coral.
- coenobium (sēnō'bium) n. [Gk. koinos, common; bios, life.] A unicellular colony with no marked distinction between vegetative and reproductive units; colony or unit of undifferentiated cells.
- coenoblast (sē'nöblāst) n. [Gk. koinos, common; blastos, bud.] A germ-layer which gives origin to endoderm and mesoderm.
- coenocentre (sē'nosčn'tër) n. [Gk. koinos, common; kentron, centre.] A deeply-staining body accompanying the ovum in certain fungi.
- coenocyte (sē'nösīt) n. [Gk. koinos, common; kytos, hollow.] A plant body in which constituent protoplasts are not separated by cell walls.
- coenoecium (sēnē'siŭm) n. [Gk. koinos, common; oikos, house.]
 The common groundwork of a polyzoan colony.
- coenogamete (sē'nögămēt') n. [Gk. koinos, common; gametes, spouse.]
 A multinucleate gamete.
- coenogenesis (sē'nöjēn'ēsīs) n. [Gk. koinos, common; genesis, descent.]
 Common descent from the same ancestry; blood relationship.
- coenosarc (sē'nösârk) n. [Gk. koinos, common; sarx, flesh.] The common tissue uniting the polyps in a compound colony.
- coenosteum (sēnŏs'tēum) n. [Gk. koinos, common; osteon, bone.] The common colonial skeleton in corals.

- coenozygote (sē'nozī'gōt) n. [Gk. koinos, common; zygon, yoke.] A zygote formed by coenogametes.
- coenurus (sēnū'rūs) n. [Gk. koinos, common; oura, tail.] A metacestode with large bladder, from whose walls many daughter-cysts arise, each with one scolex.
- co-enzyme (kō'en'zīm) n. [L. cum, with; Gk. en, in; zyme, leaven.] A substance which activates an enzyme or accelerates its action; co-ferment.
- cog-tooth,—spur or projection of incudal facet of malleus.
- coherent (köhë'rënt) a. [L. cohaerere, to stick together.] With similar parts united; adherent.
- cohesion (köhē'zhǔn) n. [L. cohaerere, to stick together.] Condition of union of separate parts of floral whorl; cf. adhesion.
- cohort (kō'hôrt) n. [L. cohors, enclosure.] A group of related families; in earlier classifications a somewhat indefinitely limited group.
- coition (köĭsh'ŭn) n. [L. coire, to go together.] Sexual intercourse; coitus; copulation.
- colchicine (köl'kisin) n. [L. colchicum, meadow saffron, from Colchis, ancient Mingrelia.] An alkaloid obtained from meadow saffron, influencing mitosis and tissue metabolism; C₂₂H₂₅O₆N.
- coleogen (köl'eöjen) n. [Gk. koleos, sheath; gennaein, to produce.] Meristematic layer giving rise to endodermis.
- coleopterous (köl'ëŏp'tërūs) a. [Gk. koleos, sheath; pteron, wing.]
 Having the anterior wings hard and used as elytra; pert. beetles.
- coleoptile (kŏl'ĕŏptil) n. [Gk. koleos, sheath; ptilon, feather.] The first leaf in seedling of monocotyledons.
- coleorhiza (köl'ëöri'ză) n. [Gk. koleos, sheath; rhiza, root.] The layer surrounding the radicle.
- colic (köl'ik) a. [Gk. kolon, colon.]

 Pert. the colon.
- collagen (köl'ājēn) n. [Gk. kolla, glue; genos, descent.] A scleroprotein, occurring as chief constituent of white connective tissue

fibres and organic part of bone, also of some fish scales.

- collar (kŏl'ăr) n. [M.E. coler, collar.] The choana of a collared cell; a prominent fold behind the proboscis in Hemichorda; the fleshy rim projecting beyond the edge of a snail shell; any structure comparable with a collar; collum, q.v.; junction between root and stem; collet, q.v.
- collateral (kölät'eräl) a. [L. cum, with; latera, sides.] Side by side; appl. ovules; appl. bundles with xylem and phloem in the same radius; appl. fine lateral branches from the axon of a nerve cell; appl. prevertebral ganglia of sympathetic system; appl. inheritance of character from a common ancestor in individuals not lineally related; appl. circulation established through anastomosis with other parts when the chief vein is obstructed.
- collenchyma (kölěng'kimå) n. [Gk. kolla, glue; engcyhma, infusion.] Parenchymatous peripheral supporting tissue with cells more or less elongated and thickened, either at the angles (angular c.), or on walls adjoining intercellular spaces (lacunar c.), or tangentially (lamellar c.); the middle layer of sponges; collenchyme.
- collencyte (köl'ěnsīt) n. [Gk. kolla, glue; en, in; kytos, hollow.] A clear cell with thread-like pseudopodia found in sponges.
- collet (köl'ēt) n. [F. collet, collar.]
 Root zone, of hypocotyl, where cuticle is absent.
- colleterium (köl'ětē'rĭūm) n. [Gk. kolla, glue.] A colleterial or mucus-secreting gland in female reproductive system of insects.
- colleters (kŏlē'tërz) n. plu. [Gk. kolletos, glued.] The hairs, usually secreting a gluey substance, which cover many resting buds; multicellular glandular trichomes.
- colle ocystophore (köle'tösist'öför)

 n. Gk. kolletos, glued; kystis,
 biacder; pherein, to bear.] The
 statorhabd of Haliclystus.

- colliculus (kŏlĭk'ūlŭs) n. [L. colliculus, little hill.] A prominence of corpora quadrigemina; a rounded elevation near apex of antero-lateral surface of arytaenoid cartilages; slight elevation formed by optic nerve at entrance to retina; elevation of urethral crest, with openings of ejaculatory ducts and prostatic utricle.
- colloblast (kŏl'öblāst) n. [Gk. kolla, glue; blastos, bud.] A cell on tentacles and pinnae of ctenophores, which carries little globules of adhesive substance; lasso-cell.
- colloid (köl'oid) n. [Gk. kolla, glue; eidos, form.] A gelatinous substance which does not readily diffuse through an animal or vegetable membrane; opp. crystalloid; a substance composed of two homogeneous parts or phases, one of which is dispersed in the other.
- collophore (kŏl'ofōr) n. [Gk. kolla, glue; pherein, to bear.] The ventral tube of Collembola.
- collum (köl'ŭm) n. [L. collum, neck.] Neck; collar, q.v.; any collar-like structure; dorsal plate of first body-segment in Diplopoda; basal portion of sporogonium in mosses.
- colon (kō'lŏn) n. [Gk. kolon, colon.] The second portion of intestine of insects; part of the large intestine of vertebrates.
- colony (köl'öni) n. [L. colonia, farm.] Any collection of organisms living together, appl. ants, bees; a group of animals or plants living together and somewhat isolated, or established in a new area; a coenobium; a group of bacteria or of other micro-organisms in a culture.
- colostrum (kölös'trum) n. [L. colostrum.] Milk secreted at end of pregnancy and differing from that secreted later.
- colulus (köl'ūlūs) n. [Dim. of L. colus, distaff.] A small conical structure between anterior spinnerets of spiders.
- columella (köl'ūměl'ă) n. [L columella, small column.] A prolongation of stalk into sporangium;

- central core in root-cap; central pillar in skeleton of some corals; the central pillar in gasteropod shells; epipterygoid; the rod, partly bony, partly cartilaginous, connecting tympanum with inner ear in birds, reptiles, and amphibia; the axis of cochlea; lower part of nasal septum.
- columellar (köl'űmĕl'ár) a. [L. columella, small column.] Pert. columella.
- column (kŏl'ŭm), columna (kŏlŭm'nā) n. [L. columna, pıllar.] Any
 structure like a column, as spinal
 column; actinian body; stalk of a
 crinoid; longitudinal bundle of
 nerve fibres in white matter of
 spinal cord; nasal septum edge;
 thick muscular strands found in
 ventricle; stamens in mallows;
 united stamens and style in orchids.
 columnals (kŏlŭm'nālz) n. blu. [L.
- columnals (kŏlŭm'nălz) n. plu. [L. columna, pillar.] Stem ossicles in crinoids.
- columnar (kölüm'när) a. [L. columna, pillar.] Pert., or like, a column or columna; appl. cells longer than broad; appl. epithelium of columnar cells.
- coma (kō'mā) n. [Gk. kome, hair.] A terminal cluster of bracts, as in pine-apple; hair-tufts on certain seeds. [Gk. koma, deep sleep.] Stupor.
- comb-ribs, combs, see swimming-plates.
- comes (kō'mēz) n. [L. comes, companion.] A blood-vessel that runs alongside a nerve.
- comitalia (kŏmitā'lĭā) n. plu. [L. comitari, to accompany.] Small di- or tri-actine spicules in sponges.
- comma (kŏm'ā) n. [Gk. komma, short clause.] A sarcomere; inocomma; appl. tract, certain nerve fibres in dorsal or posterior column of spinal cord; appl. bacillus, the spirillum causing cholera.
- commensal (kŏměn'săl) n. [L. cum, with; mensa, table.] An organism living with another and sharing the food, both species as a rule benefiting by the association.

- comminator (köm'inātör) a. [L. cum. with; minari, to threaten.] Appl. muscles which connect adjacent jaws in Aristotle's lantern.
- commissure (kŏm'ísūr) n. [L. commissura, seam.] The union-line between two parts; inner side of mericarp; carpellary cohesion plane; a connecting band of nerve tissue.
- comose (kō'mōs) a. [L. comosus, hairy.] Hairy; having a tuft of hairs.
- companion cell,—a narrow cell, retaining its nucleus, derived from a cell giving rise also to a sievetube element, in phloem of angiosperms.
- compass (kŭm'păs) n. [L. cum, together; passus, pace.] A curved bifid ossicle, part of Aristotle's lantern.
- compass plants,—certain plants with permanent north and south direction of their leaf edges.
- compensation point,—incidence of balance between respiration and photosynthesis, as determined by intensity of light at a given temperature: compensation intensity; limit of sea depth below which plants lose more by respiration than they gain by photosynthesis: compensation depth.
- competence (kŏm'pëtëns) n. [L. competere, to suit.] Reactive state permitting directional development and differentiation in response to a stimulus, as of part of an embryo in response to an evocator or organiser stimulus.
- complement (kŏm'plëmënt) n. [L. complere, to fill up.] The substance in the blood-serum which when destroyed by heat acts with an amboceptor to produce lysis; alexin; a group composed of one, two, or more genomes or chromosome sets derived from a single nucleus.
- complemental air,—volume of air which can be taken in addition to that drawn in during normal breathing.
- complemental male,—a purely male

- form, usually small, found living in close proximity to the ordinary hermaphrodite form in certain animals, as barnacles.
- complementary (kōm'plėmčn'tari) n. [L. complere, to fill up.] The coronoid bone. a. Appl. non-suberised cells loosely arranged in cork tissue and forming air passages; appl. genes producing a similar effect when inherited separately but a different effect together.
- complexus (kömplěk'sŭs) n. [L. complexus, embrace.] An aggregate; appl. muscle, the semispinalis capitis.
- complicant (kŏm'plikănt) a. [L. cum, together; plicare, to fold.] Folding over one another; appl. elytra of certain insects.
- complicate (kŏm'plikāt) a. [L. cum, together; plicare, to fold.] Folded; conduplicate; appl. leaves folded longitudinally so that right and left halves are in contact; appl. insect wings.
- composite (kŏm'pösit) a. [L. cum, together; ponere, to place.] Closely-packed, as a capitulum; appl. fruits, as sorosis, syconus, strobilus.
- compound (kŏm'pownd) a. [L. cum, together; ponere, to place.] Made up of several elements; appl. flowers, pistils, leaves, medullary rays, eyes, etc.; appl. starch grains with two or more hila.
- compressor (kömpres'ör) n. [L. cum, together; premere, to press.] Something that serves to compress; appl. muscles, as compressor naris.
- conarium (kōnā'rīum) n. [Gk. konarion, little cone.] Transparent deep-sea larva of Velella; the pineal gland or epiphysis cerebri.
- concatenate (kŏnkăt'ēnāt) a. [L. cum, together; catenatus, chained.] Forming a chain, as spores.
- concentric (könsĕn'trik) a. [L. cum, together; centrum, centre.] Having a common centre; appl. vascular bundles with one kind of tissue surrounding another; appl. corpuscles of Hassall.

conceptacle (könsép'tákl) n. [L. concipere, to conceive.] A depression in thallus of certain algae in which gametangia are borne.

conceptive (kŏnsĕp'tĭv) a. [L. concipere, to conceive.] Capable of being fertilised and producing an

embryo.

concha (kŏng'kă) n. [Gk. kongche, shell.] The cavity of the external ear, which opens into the external acoustic meatus; a superior, middle, and inferior projection from lateral wall of nasal cavity; turbinal body; one of two curved plates of sphenoidal bone; a marine shell.

conchiform (kŏng'kĭfôrm) a. [L. concha, shell; forma, shape.]
Shaped like a concha; shell-

shaped; conchoid.

conchiolin (köngki'ölin) n. [Gk. kongche, shell.] The organic substance that forms the basis of shells of molluscs.

- conchology (kŏngkŏl'öji) n. [Gk. kongche, shell; logos, discourse.]
 The branch of zoology dealing with molluscs or their shells.
- conchula (kŏng'kūlā) n. [L. concha, shell.] The conspicuous protuberant lip of the modified sulcus in Peachia.
- concolorate (kŏnkŭl'orāt) a. [L. concolor, of the same colour.] Similarly coloured on both sides.

concrescence (könkrés'éns) n. [L. concrescere, to grow together.] The growing together of parts.

concrete (könkrēt) a. [L. concretus, grown together.] Grown together to form a single structure.

condensation (kŏndĕnsā'shŭn) n. [L. condensatio; from cum, together, densare, to make thick.] Process of making or becoming thick; contraction, thickening and spiralisation of chromatids during prophase.

condensed (kŏndĕn'sd) a. [L. condensare, to press close together.] Appl. inflorescence with shortstalked or sessile flowers closely crowded.

conditional,—appl. dominance owing to influence of modifying genes.

conditioned, appl. reflex depending on new functional connections in central nervous system; appl. stimulus inducing a conditioned reflex.

conducting (kŏndŭk'tĭng) a. [L. conducere, to lead together.] Conveying; appl. tissues, bundles.

conduction (kŏndūk'shūn) n. [L. conducere, to lead together.] The transference of soluble matter from one part of a plant to another; the transmission of an excitation, function of nervous system.

conductivity (kŏn'dŭktīv'iti) n. [L. conducere, to lead together.] Power

of transmitting an impulse.

conductor (kŏndŭk'tor) n. [L. conducere, to lead together.] That which can transmit; a projection at base of embolus in spiders.

conduplicate (köndű'plikāt) a. [L. conduplicare, to fold together.] Appl. cotyledons folded to embrace the radicle; appl. vernation when one half of the leaf is folded upon the other.

condylar (kŏn'dĭlăr) a. [Gk. kondy-los, knuckle.] Pert. a condyle.

condyle (kŏn'dīl) n. [Gk. kondylos, knuckle.] The antheridium of stoneworts; a process on a bone for purposes of articulation; a rounded structure adapted to fit into a socket.

condyloid (kŏn'dĭloid) n. [Gk. kon-dylos, knuckle; eidos, form.]
Shaped like, or situated near a condyle.

cone (kōn) n. [Gk. konos, cone.]
The female flower of Coniferae, with woody axis and spirally-arranged carpels; strobile; terminal spike or fructification in clubmosses and horsetails; a conical elevation on an egg just before fertilisation; a conical or flask-shaped cell of the retina.

cone of origin,—small clear area of nerve cell at the point of exit of the axon; implantation cone; axon hill.

cone of Wulzen [R. Wulzen, American physiologist]. A structure projecting forwards from pars intermedia into hypophysial cavity in pituitary region of ox and pig.

- cone-bipolars, bipolar cells whose inner ends ramify in contact with dendrites of ganglionic cells.
- confluence (kon'flooëns) n. [L. cum, with; fluere, to flow.] Angle of union of superior sagittal and transverse sinuses at occipital bone; torcular Herophili.

congeneric (kŏn'jenĕr'ik) a. [L. congener, of same race.] Belonging to the same genus.

- congenetic (kŏnjĕnĕt'îk) a. [L. cum, with; Gk. genesis, descent.]
 Having the same origin; alike in descent.
- congenital (kŏnjĕn'itāl) a. [L. cum, with; gignere, to beget.] Present at birth; born with.
- congestin (kŏnjĕs'tĭn) n. [L. congestus, heaped up.] A toxin of seaanemone tentacles.
- conglobate (kŏn-glō'bāt) a. [L. conglobatus, formed into a ball.] Ball-shaped; appl. gland on lower side of ductus ejaculatorius in insects.
- conglomerate (kŏn-glŏm'ërāt) a. [L. cum, together; glomerare, to wind.] Bunched or crowded together.
- congression (köngrësh'ŭn) n. [L. congressio, meeting.] Chromosome movement to equatorial plane of spindle at metaphase.
- coni (kō'nī) n. plu. [L. conus, cone.] Cones; coni vasculosi; lobules forming head of epididymis.
- conidial (könĭd'ĭăl)a. [Gk. konis, dust; idion, dim.] Pert. a conidium.
- conidiferous (könĭd'ĭif'ërŭs) a. [Gk. konis, dust; idion, dim.; I.. ferre, to bear.] Bearing conidia.
- conidiocarp (könid'iökârp) n. [Gk. konis, dust; idion, dim.; karpos, fruit.] A collection of conidiophores enclosed in a covering; a pycnidium.
- conidiophore (könīd'iöfōr) n. [Gk. konis, dust; idion, dim.; pherein, to bear.] A hypha with sterigmata which bear conidia.
- conidiospore (könĭd'iöspōr) n. [Gk. konis, dust; idion, dim.; sporos, seed.] Spore or conidium produced when dry conditions inhibit

reproduction by zoospores, in Phycomycetes.

- conidium (könid'ium) n. [Gk. konis, dust; idion, dim.] A fungal spore asexually produced by constriction of sterigma or of part of a hypha; gonidium.
- coniferous (könif'erus) a. [L. conus, cone; ferre, to bear.] Conebearing.
- conjugate (kŏn'joogāt) v. [L. conjugare, to join together.] To unite, as protozoa; to undergo conjugation. a. United in pairs; appl. pores united by a groove; appl. division in pairs of monoploid nuclei.
- conjugated (kŏn'joogātèd) a. [L. conjugare, to join together.] United; appl. protein, when molecule united to non-protein molecule.
- conjugation (kŏn'joogā'shŭn) n. [L. cum, together; jugare, to yoke.] The temporary union or complete fusion of two gametes or unicellular organisms; the pairing of chromosomes.
- conjunctiva (kŏn'jŭngktī'vă) n. [L. cum, together; jungere, to join.]
 Mucous membrane of eye, lining eyelids and reflected over fore part of sclera and constituting corneal epithelium.
- conjunctive (kŏnjŭngk'tĭv) a. [L. cum, together; jungere, to join.]
 Appl. parenchyma cells in interspaces of stelar elements.
- connate (kŏn'nāt, kŏnāt') a. [L. cum, together · gnatus, born.] Firmly joined together from birth; connate-perfoliate, joined together at base so as to surround stem, appl. opposite sessile leaves.
- connective (kŏnčk'tīv) n. [L. connectere, to bind together.] A connecting band of nerve tissue between two ganglia; tissue separating two lobes of anther.
- connective tissue,—a mesoblastic tissue with a large amount of intercellular substance, and usually connecting and supporting other tissues.

connexivum (köněk'sĭvům) n. [L. connectere, to fasten together.] Flattened lateral margin of abdomen in bugs.

connivent (kŏnī'vënt) a. [L. connivere, to close the eyes.] Converging; arching over so as to meet.

conoid (kō'noid) a. [Gk. konos, cone; eidos, form.] Cone-like, but not quite conical.

conoid ligament,—one of the fasciculi of the coraco-clavicular

ligament.

conoid tubercle,—coracoid tuberosity, a small rough eminence on posterior border of clavicle, serving for attachment of conoid ligament.

conotheca (konothe'ka) n. [Gk. konos, cone; theke, case.] Thin integument of phragmocone.

- consensual (kŏnsĕn'sūăl) a. [L. consensus, agreement.] Appl. involuntary action correlated with voluntary action; reacting to excitation of a corresponding organ; appl. contraction of both pupils when only one retina is directly stimulated.
- consociation (kŏn'sōsĭā'shŭn) n. [L. consociatio, partnership.] A unit of a plant association, characterised by a single dominant species.
- consocies (könsö'siēz) n. [L. cum, together; socius, fellow.] A consociation representing a stage in the process of succession.
- consortes (kŏnsôr'tēz) n. plu. [L. consortes, partners.] Associate organisms other than symbionts, commensals, or hosts and parasites. Sing. consors.
- consortium (kŏnsôr'tĭum) n. [L. consortium, partnership.] The compound thallus of lichens.
- constricted (könstrik'tëd) a. [L. constrictus, drawn together.] Narrowed; compressed at regular intervals.
- constriction (könstrik'shun) n. [L. constrictus, drawn together.] A constricted part or place, as a node of Ranvier; non-spiralising chromosome segment at metaphase,

- either associated with the centromere, or acentric, or controlled by the nucleolus.
- constrictor (könstrik'tör) n. [L. constrictus, drawn together.] A muscle which compresses or constricts, e.g., constrictor pharyngis, c. urethrae.
- consute (kŏn'sūt) a. [L. consuere, to sew together.] With stitch-like markings; appl. elytra of certain beetles.
- contabescence (kŏn'tăbĕs'ëns) n.
 [I.. contabescere, to waste away.]
 Abortion or atrophy of stamens.
- contact receptor,—a receptor in epidermis or in dermis.
- context (kön'těkst) n. [L. cum. together; texere, to weave.] The layers developed between hymenium and true mycelium in certain fungi.
- continuity (kŏntĭnū'ĭtĭ) n. [L. continuus, continuous.] Succession without a break, especially continuity of germ plasm.
- contorted (köntôr'têd) a. [L. contortus, twisted together.] Twisted; appl. aestivation in which one leaf overlaps the next with one margin, and is overlapped by the previous on the other.
- contortuplicate (kŏntŏr'tūplĭkāt a. [L. cum, with; torquere, to twist; plicare, to fold.] Appl. bud with contorted and plicate leaves.
- contour (kŏn'toor) n. [F. contourner, to twist.] Outline of a figure or body; appl. outermost feathers that cover the body of a bird.
- contractile (könträk'til) a. [L. cum, together; trahere, to draw.] Capable of contracting.
- contractile cell,—any cell in a sporangium or an anther wall which by hygroscopic contraction helps to open the organ.
- contractile fibre-cells, elongated, spindle-shaped, more or less polyhedral, nucleated muscle-cells, containing a central bundle of fibrillae.
- contractile vacuole,—a small spherical vesicle, found in cytoplasm of many Protista, with excretory or hydrostatic function.

- contractility (kön'trāktīl'ītī) n. [L. cum, together, trahere, to draw.]
 The power by which muscle-fibres are enabled to contract.
- contractin, presumable neurohumor inducing contraction of chromatophores in crustaceans. Opp. expantin.
- contracture (könträk'tür) n. [L. contractus, drawn together.] Contraction of muscles persisting after stimulus has been removed.
- contra-deciduate (kŏn'trădēsĭd'ūāt)
 a. [L. contra, opposite to; decidere,
 to fall off.] Appl. foetal placenta
 and distal part of allantois.
- contranatant (kŏn'trănā'tănt) a. [L. contra, against; natare, to swim.] Swimming or migrating against the current, opp. denatant.
- conuli (ko nuli) n. plu. [Dim. of L. conus, cone.] Tent-like projections on surface of certain sponges caused by principal skeletal elements.
- conus (kō'nūs) n. [L. conus, cone.]
 Any cone-shaped structure, as conus arteriosus, a structure between ventricle and aorta in fishes and amphibians; diverticulum of right ventricle from which pulmonary artery arises; conus medullaris, the tapering end of spinal cord.
- convergence (kŏnvĕr'jēns) n. [I.. convergere, to incline together.]

 The development of similar characters in organisms belonging to different groups; heterogenetic homoeomorphosis; homoplasty.
- convolute (kon'volut) a. [L. cum, together; volvere, to wind.] Rolled together; appl. leaves and cotyledons; appl. shells in which outer whorls overlap inner; coiled; convoluted, appl. parts of renal tubule.
- convolution (kŏn'völū'shūn) n. [L. cum, together; volvere, to wind.] A coiling or twisting, as of brain, intestine.
- coprodaeum (köp'rödē'um) n. [Gk. kopros, dung; odos, way.] The division of cloaca which receives rectum.
- coprolite (kop'rolit) n. [Gk. kopros,

- dung; lithos, stone.] Petrified faeces.
- coprophagous (köpröf'ágŭs) a. [Gk. kopros, dung; phagein, to eat.] Feeding on dung; appl. insects.
- coprophil (kŏp'röfil) a. [Gk. kopros, dung; philos, loving.] Appl. dung bacteria and flagellates; coprophytic.
- coprozoic (kŏp'rözō'ĭk) a. [Gk. kopros, dung; zoon, animal.] Inhabiting faeces, as some protozoa.
- copula (kŏp'ūlă) n. [L. copula, bond.] A ridge in development of the tongue, formed by union of ventral ends of second and third arches; basihyal or os interglossum in certain reptiles; any bridging or connecting structure.
- copularium (köpülä'riüm) n. [L. copula, bond.] A cyst formed around two associated gametocytes, in gregarines.
- copulation (kŏpūlā'shūn) n. [L. copula, bond.] Sexual union; coition; in protozoa, complete fusion of two individuals; conjugation, as in yeasts.
- coracoid (kör'ākoid) a. [Gk. korax, crow; eidos, form.] Appl. or pert. bone or part of the pectoral girdle between scapula and sternum; appl. ligament which stretches over the suprascapular notch.
- coracoid process, the rudimentary coracoid element fused to the scapula in most mammals.
- coralliferous (kŏrălif'ërŭs) a. [Gk. korallion, coral; L. ferre, to bear.]
 Coral-forming; containing coral.
- coralline (kŏr'alīn) a. [Gk. korallion, coral.] Resembling a coral; appl. Hydrozoa and Polyzoa; composed of coral; appl. certain Algae; appl. a Pliocene crag or deposit containing fossil Polyzoa and Mollusca.
- corallite (kŏr'ālīt) n. [Gk. korallion, coral.] Cup of a single polyp of coral.
- coralloid (kŏr'ăloid) a. [Gk. korallion, coral; eidos, form.] Resembling, or branching like a coral; appl. gleba, roots, etc.

- corallum (köräl'ŭm) n. [Gk. korallion, coral.] Skeleton of compound coral.
- corbiculum (kôrbĭk'ūlŭm) n. [L. dim. of corbis, basket.] Fringe of hair on insect tibia; the pollencollecting apparatus of a bee. Plu. corbicula.
- corbula (kôr'būlă) n. [L. corbis, basket.] The phyllactocarp of Aglaeophenia, etc., a stem with alternate branches rising upwards and forming a pod-like structure.
- cord (kôrd) n. [Gk. chorde, cord.] Any cord-like structure, as spinal cord, spermatic cord.
- cordate (kôr'dāt) a. [L. cor, heart.] Heart-shaped; cordiform.
- cordiform tendon,—the central aponeurosis of the diaphragm.
- cordylus (kôrdi'lus) n. [Gk. kordyle, swelling.] An intertentacular exumbral structure with core of vacuolated cells and flattened ectoderm.
- coremata (korē'mātā) n. plu. [Gk. korema, broom.] Paired sacs bearing hairs, on membrane between seventh and eighth abdominal segments, accessory copulatory organ in moths.
- coremiform (kŏrē'mĭfôrm) a. [Gk. korema, broom; L. forma, shape.] Formed like a broom or sheaf.
- coremium (körē'miŭm) n. [Gk. korema, broom.] A sheaf-like aggregation of conidiophores, or of hyphae.
- coriaceous (kōriā'shūs) a. [L. corium, leather.] Leathery; appl. leaves.
- corium (kō'riŭm) n. [L. corium, leather.] The middle division of an elytron; deeper-seated layer of the skin, consisting of a vascular connective tissue; cutis vera; derma.
- cork (kôrk) n. [Sp. alcorque, cork.] A tissue derived usually from outer layer of cortex in woody plants.
- cork-cambium,—phellogen, q.v.
 corm (kôrm), cormus (kôr'mŭs) n.
 [Gk. kormos, trunk.] An enlarged solid subterranean stem, rounded in shape, composed of two or more

- internodes and covered externally by a few thin membranous scales or cataphyllary leaves.
- cormel (kôr'měl) n. [Gk. kormos, trunk.] A secondary corm produced by an old corm.
- cormidium (kôrmid'ium) n. [Gk. kormos, trunk; idion, dim.] An aggregation of individuals in a siphonophore, borne on the coenosarc and capable of liberation therefrom.
- cormoid (kôr'moid) a. [Gk. kormos, trunk; eidos, form.] Like a corm.
- cormophylogeny (kôr'möfilŏj'ënī) n. [Gk. kormos, trunk; phyle, tribe; genos, offspring.] Development of families or races.
- cormophyte (kôr'möfit) n. [Gk. kormos, trunk; phyton, plant.] A plant which possesses stem and root. Opp. thallophyte.
- cormous (kôr'mŭs) a. [Gk. kormos, trunk.] Corm-producing.
- cornea (kôr'nëă) n. [L. corneus, horny.] The transparent covering on anterior surface of eycball; outer transparent part of each element of a compound eye.
- corneagen (kôr neajen) a. [L. cornu, horn; Gk. -genes, producing.] Cornea-producing; appl. cells immediately below cuticle, which secrete cuticular lens and are renewed on ecdysis.
- corneal (kôr'neăl) a. [L. corneus, horny.] Pert. the cornea.
- corneoscute (kôr'nëöskūt) n. [L. corneus, horny; scutum, shield.]
 An epidermal scale.
- corneous (kôr'nëŭs) a. [L. corneus, horny.] Horny; appl. sheath covering bill of birds.
- cornicle (kôr'nĭkl) n. [L. dim. of cornu, horn.] A wax-secreting organ of aphids; corniculum, q.v.
- corniculate (kôrnik'ūlāt) a. [L. dim. of cornu, horn.] Having small horns.
- corniculate cartilages,—two small, conical, elastic cartilages articulating with apices of arytaenoids; Santorini's cartilages.

- corniculum (kôrnĭk'ūlŭm) n. dim. of cornu, horn.] A small horn or horn-like process.
- cornification (kôr'nĭfĭkā'shŭn) n. [L. cornu, horn; facere, to make.] Formation of outer horny layer of epidermis.
- cornua (kôr'nūă) n. plu. [L. cornu, horn.] Horns; horn-like prolongations; appl. bones, nerve tissues, cavities, etc.; the dorsal, lateral, and ventral columns of grey substance in spinal cord. Sing. cornu. cornucopia (kôr'nūkō'piă) n. [L.

cornu, horn; copia, plenty.] horizontal portion of taeniae of fourth ventricle.

cornute (kôrnūt') a. [L. cornutus, horned.] With horn-like processes. corolla (köről'ă) n. [L. corolla, small crown.] The petals of a flower.

corollaceous (kor'ola'shus) a. [L. corolla, crown.] Pert. a corolla.

- corolliferous (kor'olif'erus) a. [L. corolla, small crown; ferre, to bear.] Having a corolla.
- corona (körö'nă) n. [L. corona, crown.] A cup-shaped body formed by union of scales on perianth leaves, as in daffodil; theca and arms of a crinoid; echinoid test excepting apical and antapical plates; ciliated disc or circular band of certain animals; head or upper portion of any structure.

corona radiata,—layer of cells surrounding mammalian egg; fibres of internal capsule of brain.

- coronal (kŏr'ŏnăl) a. [L. corona, crown.] Pert. corona; appl. suture between frontal and parietal bones; situated in the coronal sutural plane; appl. later roots of grasses, opp. seminal.
- coronary (kor'onari) a. [L. corona, crown.] Crown-shaped or crownlike; encircling; appl. arteries, bones, sinus, ligaments, plexus, vein.

coronary arteries,—arteries supplying tissue of heart; labial arteries. coronary bone,—a small conical

bone in mandible of reptiles; small pastern bone of horse.

coronary sinus,—channel receiving

- most cardiac veins and opening into right auricle.
- coronate (kŏr'önāt) a. [L. corona, crown.] Having a corona; having a row of tubercles encircling a structure, or mounted on whorls of spiral shells.
- coronet (kŏr'önĕt) n. [L. corona, crown.] The burr of an antler.
- coronoid (kŏrō'noid) a. [Gk. koronis, crook-beaked; eidos, form.] Shaped like a beak; appl. processes. [L. corona, crown.] n. Coronary bone of reptiles.
- coronula (kŏrŏn'ūlă) n. [Dim. of L. corona, crown.] A group of cells forming a crown on the oosphere, as in Charophyta.
- corpora (kôr'pöra) n. plu. corpus, body.] Bodies. See corpus. **corpora adiposa,**—fat-bodies, q.v.
- corpora albicantia,—white bodies or scars formed in ovarian follicle after disintegration of luteal cells; corpora mamillaria, ganglia below posterior part of third ventricle.
- corpora allata,—paired ovoid whitish endocrine glands in insects.
- corpora amylacea, brain sand.
- corpora bigemina,—the optic lobes of vertebrate brain, corresponding to the superior colliculi of corpora quadrigemina of mammals.
- corpora cardiaca,—neuroglandular bodies between cerebral ganglia and corpora allata, in some insects.
- corpora cavernosa,—erectile masses of tissue, forming anterior part of body of penis; erectile tissue of clitoris.
- corpora mamillaria, two white bodies enclosing grey matter in hypothalamus, beneath floor of third ventricle; corpora albicantia.
- quadrigemina, four corpora rounded eminences which form dorsal part of mesencephalon.
- corpus (kôr'pŭs) n. [L. corpus, body.] Body; any fairly homogeneous structure which forms part of an organ; plu. corpora.
- corpus albicans,-white tissue replacing corpus luteum in nonpregnancy.

- corpus callosum,—the broad transverse band of white substance connecting the cerebral hemispheres.
- corpus fibrosum,—fibrous tissue remaining after disintegration of corpus luteum.
- corpus geniculatum, geniculate body, q.v.
- corpus haemorrhagicum, body developed from ruptured Graafian follicle around blood clot, and later developing into corpus luteum.
- corpus luteum,—the glandular body developed from a Graafian follicle after extrusion of ovum; yellow body.
- corpus spongiosum,—a mass of erectile tissue forming posterior wall of penis; corpus cavernosum urethrae.
- corpus sterni,—sternebrae fused into a single mesosternal bone; the gladiolus.
- corpus striatum,—a mass of grey matter containing white nerve fibres and consisting of the caudate nucleus which projects into the lateral ventricle, and of the lenticular nucleus.
- corpuscle (kôr'pŭsl, kôrpŭs'l) n.
 [L. corpusculum, small body.] A
 protoplasmic cell, floating freely in a
 fluid, or embedded in a matrix; any
 minute particle, as in a cell.
- correlation (kŏr'ĕlā'shūn) n. [L.L. correlatio, relationship.] Mutual relationship; proportional growth; interdependence of characters, particularly of quantitative characters, measured by correlation coefficient which is plus or minus one if characters are exactly inter-related, and zero if entirely unrelated; combination of nervous impulses in sensory centres, resulting in adaptive reactions.
- correlator (kŏr'ēlātör) n. [L.L. correlatio, relationship.] A diffusible substance correlating activities of coleoptile tip and hypocotyl; auxin, q.v.
- corrugator (kor'oogātor) a. [L. cum, together; rugare, to wrinkle.]

- Wrinkled or wrinkling; appl. muscles.
- cortex (kôr'těks) n. [L. cortex, bark.] The extrastelar fundamental tissue of the sporophyte; outer or more superficial part of an organ.
- cortical (kôr'tĭkăl) a. [L. cortex, bark.] Pert. the cortex.
- corticate (kôr'tĭkāt) a. [L. cortex, bark.] Having a special outer covering.
- corticiferous (kôr'tisif'ërŭs) a. [L. cortex, bark; ferre, to carry.]
 Forming or having a bark-like cortex.
- corticolous (kôrtik'olŭs) a. [L. cortex, bark; colere, to inhabit.] Inhabiting, or growing on, bark.
- corticosterone,—the active constituent of adrenal cortical hormone; C₂₁H₃₀O₄.
- corticostriate (kôr'tiköstri'āt) a. [L. cortex, bark; stria, channel.] Appl. fibres which join corpus striatum to cerebral cortex.
- corticotrophic, corticotropic, adrenocorticotrophic, q.v.
- cortin (kôr'tǐn) n. [L. cortex, bark.]
 Adrenal cortex extract, containing cortical hormones.
- cortina (kôrtē'nă) n. [L. cortina, curtain.] The velum in some agarics.
- cortinate (kôr'tĭnāt) a. [L. cortina, curtain.] Having a velum; of a cobweb-like texture.
- Corti's organ [A. Corti, Italian histologist]. The organon spirale, on inner portion of membrana basilaris of ear.
- Corti's rods,—double row of arching rods based on basilar membrane and forming the spiral tunnel of Corti.
- coruscation (koruska'shun) n. [L. coruscatio, flash.] Twinkle, rapid fluctuation in a flash or oscillation in light emission, as of fire-flies.
- corymb (kôr'imb) n. [Gk. korymbos, cluster of flowers.] A raceme with lower pedicels elongated so that the top is nearly flat.
- corymbose (körīm'bōs) a. [Gk. korymbos, cluster of flowers.] Pert. or like a corymb; arranged in a corymb; corymbous.

- cosmine (kŏs'mĭn) n. [Gk. kosmios, regular.] The outer layer of dentine-like material in cosmoid and ganoid scales.
- cosmoid (kös'moid) a. [Gk. kosmios, regular; eidos, form.] Having an outer periodically resorbed layer of cosmine, appl. dermal bones, scales, and lepidotrichia in Crossopterygii.
- cosmopolitan (közmöpöl'ītān), cosmopolite (közmöp'ölīt) a. [Gk. kosmos, world; polites, citizen.] World-wide in distribution.
- costa (kŏs'tă) n. [L. costa, rib.] A rib; anything rib-like in shape, as a ridge on shell, coral, etc.; anterior vein, or margin, of insect wing; comb-rib or swimming-plate of Ctenophora; structure at base of undulating membrane in Trichomonadidae. Plu. costac.
- costaeform (kŏs'tēfôrm) a. [L. costa, rib; forma, shape.] Rib-like; appl. unbranched parallel leaf-veins.
- costal (kôs'tăl) a. [L. costa, rib.] Pert. ribs or rib-like structures; appl. bony shields of Chelonia; pert. costa of insect wing; pert. primary brachial series in Crinoids; pert. a main rib.
- costalia (köstä'liä) n. plu. [L. costa, rib.] The supporting plates in theca of Cladoidea.
- costate (kös'tāt) a. [L. costa, rib.] With one or more longitudinal ribs; with ridges or costae.
- coterminous (kōtēr'minus) a. [L. cum, with; terminus, end.] Of similar distribution; bordering on.
- cotyledon (kötilē'don) n. [Gk. kotyle, cup.] The seed-leaf, primary or first leaf of an embryonic sporophyte; a patch of villi on mammalian placenta.
- cotyledonary (kötile'dönari) a. [Gk. kotyle, cup.] Pert. cotyledons; with villi grouped in cotyledons, appl. placenta.
- cotyloid (köt'iloid) a. [Gk. kotyle, cup; eidos, form.] Cup-shaped; pert. the acetabular cavity.
- cotylophorous (kötilöf'örüs) a. [Gk. kotyle, cup; pherein, to bear.]
 With a cotyledonary placenta.

- cotype (kō'tīp) n. [L. cum, with; typus, image.] An additional type specimen, frequently collected in same place at same time, or a specimen from a description of which, along with others, the type is defined; syntype.
- covariation (kō'vārīā'shun) n. [L. con, with; varius, diverse.] Correlation, q.v.
- cover scales,—small scales arranged spirally and developed directly on the axis of a cone of Coniferae; bract scales.
- covert (kŭv'ërt) n. [F. couvrir, to cover.] Appl. feathers covering bases of quills in birds.
- cowled (kowld) a. [L. cucullus, hood.] Furnished with or shaped like a hood.
- Cowper's glands [W. Cowper, English surgeon]. Bulbo-urethral glands, q.v.
- coxa (kŏk'să) n. [L. coxa, hip.] Proximal joint of leg of an insect or arachnid.
- coxal (kök'săl) a. [L. coxa, hip.] Pert. the coxa; appl. glands; pert. the hip.
- coxite (kŏk'sit) n. [L. coxa, hip.]
 One of paired lateral plates in contiguity with insect sternum; limb base bearing stylus in Thysanura.
- coxopleurite, catapleurite, q.v.
- coxopodite (kŏks'öpodīt) n. [L. coxa, hip; Gk. pous, foot.] The proximal part of protopodite of crustacean limb; coxa of spiders.
- coxosternum (kök'söstĕr'nŭm) n. [L. coxa, hip; sternum, breast-bone.] Plate formed by fusion of coxites and sternum; vinculum, in Lepidoptera.
- crampon (krăm'pŏn) n. [O.H.G. chramph, crooked.] An aerial root, as in ivy.
- cranial (krā'nīāl) a. [Gk. kranion, skull.] Pert. skull, or that part which encloses the brain; appl. bones, fossae, nerves, muscles, blood-vessels, etc.
- craniate (krā'niāt) a. [Gk. kranion, skull.] Having a skull.

- cranidium (krănĭd'ĭŭm) n. [Gk. kranion, skull; idion, dim.] Glabella together with fixed genae, in trilobites.
- cranihaemal (krā'nĭhē'măl) a. [Gk. kranion, skull; haima, blood.] Appl. anterior lower portion of a sclerotome.
- cranineural (krā'nĭnū'răl) a. [Gk. kranion, skull; neuron, nerve.] Appl. anterior upper portion of a sclerotome.
- craniology (krā'niŏl'öji) n. [Gk. kranion, skull; logos, discourse.]
 The study of the skull.
- craniometry (krā'nĭōm'ëtrĭ) n. [Gk. kranion, skull; metron, measure.]

 The science of the measurement of skulls.
- craniosacral (krā'nĭösā'krăl) a. [Gk. kranion, skull; L. sacer, sacred.]

 Pert. skull and sacrum; appl.

 nerves, the parasympathetic system.
- cranium (krā'nĭŭm) n. [Gk. kranion; L. cranium, skull.] The skull of any craniate, or more particularly. that part enclosing the brain.
- craspedodromous (krās'pēdŏd'römūs) a. [Gk. kraspedon, edge; dromein, to run.] With nerves running directly from mid-rib to margin.
- craspedote (krăs'pëdōt) a. [Gk. kraspedon, edge.] Having a velum.
- craspedum (krăs'pëdum) n. [Gk. kraspedon, edge.] A mesenteric filament of sea-anemones.
- crassula (krăs'ūlă) n. [L. crassus, thick.] Thickened bar on middle lamella between two bordered pits in tracheids of wood of conifers; bar of Sanio. Plu. crassulae.
- craticular (krătĭk'ūlăr) a. [L. craticula, gridiron.] Crate-like; appl. stage in life-history of a diatom where new valves are formed before the old are lost.
- creatine (krē'ātīn) n. [Gk. kreas, flesh.] A nitrogenous substance found in muscles, brain, and blood of vertebrates; C₄H₉O₂N₃.
- creatinine (krē'ātinin) n. [Gk. kreas,

- flesh.] A katabolic product in muscle and other tissues, excreted in urine; C₄H₇ON₃.
- cremaster (krëmăs'tër) n. [Gk. kremastos, hung.] A thin muscle along the spermatic cord; a stout terminal abdominal spine in subterranean insect pupae; the anal hooks for suspension pupae.
- cremocarp (krěm'ökârp) n. [Gk. kremannumai, to hang; karpos, fruit.] An inferior, dry, indehiscent, bilocular, two-seeded fruit.
- crena (krē'nă) n. [L. crena, notch.] Notch in a crenate margin, as of leaf; cleft, as anal cleft; deep groove, as longitudinal sulcus of heart.
- crenate (krē'nāt) a. [L. crena, notch.] With scalloped margin.
- crenation (krěná'shun) n. [L. crenatus, notched.] A scalloped margin, or rounded tooth, as of leaf; crenature; notched or wrinkled appearance, as of erythrocytes exposed to hypertonic solutions.
- crenulate (krčn'ūlāt) a. [Dim. of L. crena, notch.] With margins minutely crenate; crenellated, crenulated.
- crepis (krē'pis) n. [Gk. krepis, foundation.] The fundamental spicule by deposition of silica upon which a desma is formed.
- crepitation (krěpita'shun) n. [L. crepitare, to crackle.] In insects, the discharge of a fluid with an explosive sound.
- crepuscular (krepus'kūlar) a. [L. crepusculum, dusk.] Pert. dusk; flying before sunrise or in twilight.
- crescent (krěs'ënt) n. [L. crescere, to grow.] A crescentic structure.
- crescentiform (krěsěn'tifôrm) a. [L. crescere, to grow; forma, shape.] Crescent-shaped; crescentic; appl. mouth of Sipunculoidea.
- crescents of Gianuzzi, small crescent-shaped bodies of the salivary glands; demilunes of Heidenhain.

- crest (krest) n. [L. crista, crest.] A ridge on a bone; a fleshy longitudinal ridge as in newts; crown or feather tuft on head of birds.
- Cretaceous (krētā'shŭs) a. [L. creta, chalk.] Appl., and pert., the last period of the Mesozoic era.
- cribellum (krībēl'ŭm) n. [L. dim. of cribrum, sieve.] A plate perforated by openings of silk ducts in certain spiders.
- cribriform (krib'rifôrm) a. [L. cribrum, sieve; forma, shape.] Sievelike.
- cribriform organ, folded membrane carrying papillae in interradial angles of certain starfishes.
- cribriform plate, the portion of ethmoid, or of mesethmoid, perforated by many foramina for exit of olfactory nerves; lamina cribrosa.
- cricoid (krī'koid) a. [Gk. krikos, ring; eidos, form.] Ring-like; appl. cartilage in larynx, articulating with thyroid and arytaenoid cartilages; appl. placenta lacking villi on central part of disc, as in certain Edentata.
- crinome (krīn'ōm) n. [L. crinis, hair.] Network formed in cytoplasm by basophil substances reacting to vital staining.
- criocone (krī'ökōn) a. [Gk. krios, ram; konos, cone.] With uncoiled spiral shaped like ram's horn; appl. shell of certain ammonites.
- crissal (kris'al) a. [L. crissare, to move haunches.] Pert. the crissum.
- criss-cross,—appl. inheritance when offspring resemble the parent of the opposite sex.
- crissum (krīs'ŭm) n. [L. crissare, to move haunches.] The circum-cloacal region of a bird; vent-feathers or lower tail-coverts.
- crista (krīs'tă) n. [L. crista, crest.]
 A crest or ridge; projection from ectoloph into median valley in lophodont molars; a fine membrane attached to body of certain spirochaetes; ligule of palm-leaves.
- crista acustica,—thickening, covered with neuroepithelium, of mem-

- brane lining ampullae of semicircular canals; a chordotonal structure in Orthoptera.
- crista galli,—anterior median process of cribriform plate.
- crista urethralis,—verumontanum, a.v.
- cristate (kris'tāt) a. [L. cristatus, crested.] Crested; cristiform, shaped like a crest.
- crochet (krŏshā') n. [F. crochet, small hook.] The projection of the protoloph in lophodont molars; a balancer in larval salamanders; a larval locomotory hook in insects.
- crop (kröp) n. [M.E. croppe, craw.] Sac-like dilatation of gullet of a bird; a similar structure in alimentary canal of insect or worm; ingluvies.
- crosier (krō'zhyër) n. [M.E. croce, crook.] Circinate young frond of fern; hook formed by terminal cells of ascogenous hyphae; flat spiral shell, as of Spirula.
- cross (krös) n. [M.E. crois, cross.]

 An organism produced by mating parents of different breeds. v. To hybridise.
- crossing over, interchange of corresponding chromosome segments by homologous pairs of chromosomes during maturation.
- crossover,—a chromatid formed as a result of crossing-over.
- crotaphite (krōt'áfīt) n. [Gk. kro-taphos, side of forehead.] The temporal fossa.
- crotchet (kroch'ët) n. [F. crochet,
 small hook.] A curved chaeta,
 notched at the end; uncinus;
 clavus, in spiders.
- crown (krown) n. [L. corona, crown.]

 The exposed part of a tooth, especially the grinding surface; distal part of antler; crest; head; cup and arms of a crinoid; corona, q.v.; leafy upper part of a tiee.
- crozier,—crosier, q.v.
- cruciate (kroo'shiāt, kroo'siāt) a. [L. crux, cross.] Cruciform; with leaves or petals in form of a cross; X-shaped or +-shaped, appl. muscles, ligaments; crucial.

- cruciform (kroos'ifôrm) a. [L. crux, cross; forma, shape.] Arranged like the points of a cross; appl. division, promitosis in Plasmo-diophorales.
- cruor (kroo'ŏr) n. [L. cruor, blood.]
 The clots in coagulated blood.

cruorin (kroo'ŏrĭn) n. [L. cruor, blood.] Haemoglobin.

- crura (kroo'rā) n. plu. [L. crura, legs.] The shanks; leg-like or columnar structures; lumbar part of diaphragm muscle fibres; proximal processes of corpora cavernosa penis; branches of incus and stapes; pillars of subcutaneous inguinal ring; posterior pillars of fornix; crura cerebri, q.v.
- crura cerebri,—the cerebral peduncles, two cylindrical masses forming the ventrolateral portion of midbrain.
- crural (kroo'răl) a. [L. crus, leg.] Pert. the thigh.
- crureus (kroorē'ŭs) n. [L. crus. leg.] Vastus intermedius muscle of thigh.
- crus (krŭs) n. [L. crus, leg.] The shank; any leg-like organ. Plu. crura.
- crusta (krūs'tā) n. [L. crusta, shell.]
 Ventral part or base or pes of cerebral peduncles; cement layer of teeth, crusta petrosa.
- crustaceous (krūstā'shūs) a. [L. crusta, shell.] With crustacean characteristics; crustose, q.v.
- crustose (krūs'tōs) a. [L. crusta, shell.] Forming crusts on substratum, aptl. lichens.
- cryophil (kri'öfil) a. [Gk. kryos, chill; philein, to love.] Thriving at a low temperature; cryophilic.
- cryoplankton (kri'öplängk'tön) n. [Gk. kryos, frost; plangktos, wandering.] Glacial and polar plankton; algal communities thriving on snow.
- cryoscopic (krī'ösköp'īk) a. [Gk. kryos, frost; skopein, to view.] Appl. method of determining osmotic pressure, using a freezing-point depression.
- crypt (kript) n. [Gk. kryptos, hidden.]

A simple glandular tube or cavity; pit of stoma; depression in uterine mucous membrane.

- cryptic (krip'tik) a. [Gk. kryptos, hidden.] Appl. protective coloration facilitating concealment; appl. polymorphism due to presence of recessive genes; appl. species extremely similar as to external appearance but which do not normally interbreed.
- cryptocarp (krip'tokârp) n. [Gk. kryptos, hidden; karpos, fruit.] A fruit-like structure, the sporophyte phase in red algae
- cryptogam (krip'tögäm) n. [Gk. kryptos, hidden; gamos, union.] A plant without apparent reproductive organs: a spore-plant; cf. phanerogam.
- cryptogene (krĭp'töjēn) a. [Gk. kryptos, hidden; genos, origin.] Of unknown descent; having an indeterminate phylogeny.
- cryptohaplomitosis (krīp'töhāp'lömītō'sīs) n. [Gk. kryptos, hidden; haploos, simple; mitos, thread.] Type of cell-division in some flagellates where chromatin divides into two masses which pass to opposite poles without spireme-formation.
- cryptomere (krip'tömēr) n. [Gk. kryptos, hidden; meros, part.] A hidden recessive hereditary factor.
- cryptomitosis (krīp'tömītō'sīs) n. [Gk. kryptos, hidden; mitos, thread.] Division of unicellular organisms, in which chromatin assembles in the equatorial region without apparent chromosome formation.
- cryptoneurous (krĭp'tönū'rūs) a. [Gk. kryptos, hidden; neuron, nerve.] With no definite or distinct nervous system.
- cryptophyte (krip'töfit) n. [Gk. kryptos, hidden; phyton, plant.]
 A plant perennating by means of rhizomes, corms, or bulbs under ground, or of under water buds.
- cryptoplasm (krĭp'töplăzm) n. [Gk. kryptos, hidden; plasma, form.] The non-granular portion of cytoplasm.

- cryptoptile (krīp'tötīl, -ptīl) n. [Gk. kryptos, hidden; ptilon, feather.]
 A feather filament, developed from papilla.
- cryptorchid (krĭptôr'kĭd) a. [Gk. kryptos, hidden; orchis, testis.]
 Having testes abdominal in position.
- cryptorhetic (krip'törĕt'ik) a. [Gk. kryptos, hidden; rhein. to flow.] Secreting internally; endocrine.
- cryptosolenial (krip'tösölë'niăl) a. [Gk. kryptos, hidden; solen, channel.] Appl. region of attachment of Malpighian vessels to hind-gut in certain Coleoptera.
- cryptostomata (krĭp'töstðm'átá) n. plu. [Gk. kryptos, hidden; stoma, mouth.] Non-sexual conceptacles in Fucaccae.
- cryptozoic (krip'tözō'ik) a. [Gk. kryptos, hidden; zoon, animal.] Appl. fauna dwelling in darkness, or under stones, bark, etc.
- cryptozoite (krip'töző'it) n. [Gk. kryptos, hidden; zoon, animal.] Stage of sporozoite when living in tissues before entering blood.
- crypts of Lieberkühn,—see Lieberkühn's crypts.
- crystallin (kris'tălin) n. [Gk. krystallos, ice.] A globulin which is the principal constituent of lens of eye.
- crystalline (krīs'tālīn) a. [Gk. krystallinos, crystalline.] Transparent; appl. various structures.
- crystalline style,—a proteid hyaline rod with amylolytic function, in alimentary canal of some molluscs.
- crystalloid (krīs'tăloid) n. [Gk. krystallos, ice; eidos, form.] A substance which in solution readily diffuses through an animal membrane; opp. colloid; a protein crystal found in certain plant cells.
- crystal-sand,—a deposit of minute crystals of calcium oxalate, as in Solanaceae.
- crystal-spore,—an isospore containing a crystal, of Radiolaria.
- cteinophyte (tin'öfit, ktin'öfit) n. [Gk. kteinein, to kill; phyton, plant.] A parasitic plant, e.g. fungus, which destroys its host.

- ctene (tēn, ktēn) n. [Gk. kteis, comb.] The swimming-plates of ctenophores.
- ctenidium (těnīd'ĭum, ktěnīd'ĭum) n. [Gk. kteis, comb; idion, dim.] The respiratory apparatus in molluscs, feather-like or comb-like in appearance; a row of spines forming a comb in some insects.
- ctenocyst (těn'ösist, ktěn'ösist) n.
 [Gk. kteis, comb; kystis, bladder.]
 Aboral sense organ of Ctenophora.
- ctenoid (těn'oid, ktěn'oid) a. [Gk. kteis, comb; eidos, form.] With comb-like margin, as scales.
- ctenophoral (těnŏf'örăl, ktěnŏf'örăl)
 a. [Gk. kteis, comb; pherein, to bear.] Supplied with swimming-plates.
- ctetology (tētŏl'öjĭ, ktē-) n. [Gk. ktetos, acquired; logos, discourse.]
 Aspect of biology concerned with acquired characters.
- ctetosome (tē'tösōm, ktē') n. [Gk. ktetos, acquired; soma, body.] A supernumerary chromosome associated with a sex chromosome during meiosis.
- cubical (kū'bĭkăl) a. [L. cubus, cube.]
 Appl. cells as long as broad.
- cubital (kū'bītăl) a. [L. cubitalis, of elbow.] Pert. the ulna or cubitus. n. A secondary wing-quill, connected with the ulna.
- cubitus (kū'bĭtŭs) n. [L. cubitum, elbow.] The ulna, forearm; primary vein in an insect wing.
- cuboid (kū'boid) a. [Gk. kyboeides, cube-like.] Nearly cubic in shape. n. Outermost of distal tarsal bones.
- cuboidal (kūboid'ăl) a. [Gk. kyboeides, cube-like.] Pert. the cuboid.
 - cucullate (kū'kŭlāt, kūkŭl'āt) a. [L. cucullus, hood.] With hood-like sepals or petals; with prothorax hood-shaped.
- cucullus (kūkūl'ūs) n. [L. cucullus, hood.] A hood-shaped structure; upper part of harpe, in Lepidoptera.
- cuiller (kwē'yā) n. [F. cuiller, spoon.] Spoon-like terminal portion of male insect clasper.

- cuirass (kwirăs') n. [F. cuirasse, leathern jacket.] Bony plates or scales arranged like a cuirass; a lorica, q.v.
- culm (kŭlm) n. [L. culmus, stalk.] The stem of grasses and sedges.
- culmen (kul'men) n. [L. culmen, summit.] Median longitudinal ridge of a bird's beak; anterior raised part of monticulus.

cultellus (kultel'us) n. [L. cultellus, A sharp knife-like little knife.] organ, one of mouth-parts of certain blood-sucking flies.

culture (kul'tur) n. [L. cultura; colere, to till.] The cultivation of micro-organisms or tissues in prepared media.

cumulose (kū'mūlōs) a. [L. cumulus, heap.] Appl. deposits consisting chiefly of plant remains, e.g. peat.

- cumulus (kū'mūlŭs) n. [L. cumulus, heap.] The mass of epithelial cells bulging into cavity of an ovarian follicle and in which ovum is embedded: cumulus oopherus. discus proligerus.
- cuneate (kū'nëāt) a. [L. cuneatus, Wedge-shaped; wedge-shaped.] appl. leaves with broad abruptlypointed apex and tapering to the base; appl. a fasciculus and tubercle formed by a grey nucleus at posterior end of rhomboid fossa of medulla oblongata.

cuneiform (kūnē'ĭfôrm) a. cuneus, wedge; forma, shape.] Wedge-shaped; appl. distal tarsal bones; appl. a carpal bone, os triquetrum; appl. two small cartilages of larynx.

cuneus (kū'nėŭs) n. [L. cuneus, wedge.] A division of elytron of certain insects; a wedge-shaped area of the occipital lobe between calcarine fissure and medial part of parieto-occipital fissure.

cup (kup) n. [A.S. cuppe, cup.] Any structure resembling a cup.

cupula (kū'pūlă) n. [L. cupula, little tub.] The bony apex of cochlea; the part of pleura over the apex of lung; cupule, q.v.

cupulate (kū'pūlāt) a. [L. cupula,

little tub.] Cup-shaped; certain aecidia.

cupule (kūp'ūl) n. [L. cupula, little tub.] The involucre of female flower of oak, etc.; the gemmae-bearing cup of Marchantia; a small sucker of various animals.

curviserial (kŭr'visē'riăl) a. curvus, curve; series, row.] Appl. phyllotaxis in which divergence is such that orthostichies themselves are slightly twisted spirally.

- cushion (koosh'ŭn) n. [M.E. cuischen, cushion.] The central thick region in prothallus of fern; appl. habitus of many plants, as in alpine species; certain torus tubarius, prominence pharyngeal opening of Eustachian tube; tubercle or elevation of laryngeal surface of epiglottis; embryonic endocardial thickening of wall of atrial canal.
- cusp (kŭsp) n. [L. cuspis, point.] A prominence, as on teeth; a sharp point. cuspidate (kus'pidat) a. [L. cuspidare, to make pointed.] Terminating in a point; appl. leaves, teeth. cutaneous (kūtā'neŭs) a. [L. cutis, skin.] Pert. the skin.

cuticle (kū'tĭkl) n. [L. cutis, skin.] An outer skin or pellicle; the epidermis; cuticula.

cuticular (kūtĭk'ūlăr) a. [L. cutis, skin.] *Pert*. the cuticle or external integument; appl. transpiration through the cuticle.

cuticularisation (kūtik'ūlarīzā'shun) [L. dim. of cutis, skin.] Cutinisation in external layers of epidermal cells.

cutin (kū'tĭn) n. [L. cutis, skin.] A substance allied to cellulose found in external layers of thickened epidermal cells.

cutinisation (kū'tĭnīzā'shŭn) n. [L. cutis, skin.] The deposition of cutin in cell-wall, thereby forming a cuticle. cutis (kū'tis) n. [L. cutis, skin.] The corium, or deeper layer of the skin.

cutocellulose (kū'tösěl'ūlōs) n. [L. cutis, skin; cellula, small cell.] Cellulose with cutin, as in plant epidermis.

- cutose (kū'tōs) n. [L. cutis, skin.] Cutin, q.v.
- Cuvier, ducts of [G.L.C.F.D. Cuvier, French comparative anatomist]. Short veins opening into sinus venosus, and formed by union of anterior and posterior cardinal veins.
- Cuvierian organs,—glandular tubes extending from cloaca of holothurians.
- cyanic (sīān'īk) a. [Gk. kyanos, dark blue.] Blue, bluish; appl. flowers, birds' eggs.
- cyanin (sī'ānīn) n. [Gk. kyanos, cornflower.] The blue pigment or anthocyanin of the cornflower.
- cyanogenesis (sī'ānöjēn'ēsīs) n. [Gk. kyanos, blue; genesis, origin.] The elaboration of hydrocyanic acid, prussic acid, as in certain plants.
- cyanophil (sīān'ŏfīl) a. [Gk. kyanos, blue; philein, to love.] With special affinity for blue or green stains; appl. cell structure.
- cyanophycin (sī'ānöfī'sīn) n. [Gk. kyanos, blue; phykos, seaweed.] Protein reserve forming granules in peripheral region of cells in bluegreen algae; β granules.
- cyanophyll (sīǎn'öfil) n. [Gk. kyanos, blue; phyllon, leaf.] A bluishgreen colouring matter in plants.
- cyathium (sī'āthīum) n. [Gk. kyathos, cup.] The peculiar inflorescence in Euphorbia, a cup-shaped involucre with stamens and stalked gynoecium, each stamen and the gynoecium being a separate flower.
- cyathozooid (sī'āthözō'oid) n. [Gk. kyathos, cup; zoon, animal; eidos, shape.] The primary zooid in certain tunicates.
- cyathus (sī'āthŭs) n. [Gk. kyathos, cup.] A small cup-shaped organ; the gemma-cup of Marchantia.
- cycle (sīkl) n. [Gk. kyklos, circle.] The circulation of a fluid through a definite series of vessels; recurrent series of phenomena, as life cycle, ovarian cycle, etc.
- cyclic (sl'klik) a. [Gk. kyklos, circle.] Having parts of flower arranged in whorls; cyclical; periodic.

- cyclocoelic (sī'klösē'lĭk) a. [Gk. kyklos, circle; koilia, intestines.]
 With the intestine coiled in one or more distinct spirals.
- cyclogenous (sikloj'enŭs) a. [Gk. kyklos, circle; gennaein, to produce.] Exogenous; appl. a stem growing in concentric circles.
- cyclogeny (siklöj'ëni) n. [Gk. kyklos, circle; genos, generation.] Production of a succession of different morphological types in a life cycle.
- cycloid (si kloid) a. [Gk. kyklos, circle; eidos, shape.] Appl. scales with evenly curved free border.
- cyclomorial (sī'klomŏr'iăl) a. [Gk. kyklos, circle; morion, constituent part.] Appl. scales, growing in area by apposition of marginal zones, as in Palaeozoic elasmobranchs.
- cyclomorphosis (sť/klömôr'fōsīs) n.
 [Gk. kyklos, circle; morphosis, form.] A cycle of changes in form, as seasonal changes in daphnids.
- cyclopean (sĭklöpē'án), cyclopic (sĭklöp'ik) a. [Gk. kyklos, circle; ops, eye.] Appl. single median eye developed under certain artificial conditions instead of the normal pair.
- cyclosis (sǐklō'sĭs) n. [Gk. kyklosis, whirling round.] Circulation, as of protoplasm within a cell.
- cyclospermous (st'klösper'mus) a.
 [Gk. kyklos, circle; sperma, seed.]
 With embryo coiled in a circle or spiral.
- cyclospondylic (st'klöspŏndil'ik) a. [Gk. kyklos, circle; sphondylos, vertebra.] Appl. centra in which the internal calcareous matter is confined to the middle zone.
- cyclospondylous (sl'klöspŏn'dĭlŭs) a. [Gk. kyklos, circle; sphondylos, vertebra.] Appl. vertebra formed of successive concentric layers of cartilage; also cyclospondylic.
- cyesis (siē'sis) n. [Gk. kyesis, conception.] Pregnancy.
- cylindrical (silin'drikăl) a. [Gk. kylindros, cylinder.] Appl. leaves rolled on themselves, or to solid cylinder-like leaves; appl. a type of silk gland in spiders; tubuliform.

- cymba (sim'ha) n. [L. cymba, boat.] Upper part of concha of ear.
- cymbiform (sim'bifòrm) a. [L. cymba, boat; forma, shape.] Boat-shaped.
- cymbium (simbī'úm) n. [Gk. kymbion, small boat.] Boat-shaped tarsus of pedipalpus in certain spiders.
- cyme (sīm) n. [L. cyma, young sprout.] Any determinate inflorescence.
- cymose (sī'mōs) a. [L. cyma, young sprout.] Sympodially branched, appl. inflorescence.
- cymotrichous (kīmŏt'rĭkŭs, sī-) a. [Gk. kyma, wave; thrix, hair.] Having wavy hair.
- cynarrhodium (sĭnărō'dĭŭm), cynarrhodon (sĭnărōd'ŏn) n. [Gk. kyon, dog; rhodon, rose.] An etaerio with achenes placed on concave thalamus.
- cynopodous (sĭnŏp'ödŭs) a. [Gk. kyon, dog; pous, foot.] With non-retractile claws.
- cyphella (sĭfĕl'ă) n. [Gk. kyphella, hollow of ear.] Small cavity on thallus of certain lichens.
- cyphonautes (sī'fönôt'ēz) n. [Gk. kyphos, bent; nautes, sailor.]
 Young free-swimming larva of certain Polyzoa.
- cypsela (sīpsēl'ā) n. [Gk. kypsele, hollow vessel.] An inferior bicarpellary achene, as in Compositae.
- cyst (sist) n. [Gk. kystis, bladder.] The enclosing membrane round a resting cell or apocyte; a bladder or air vesicle in certain seaweeds; abnormal sac containing fluid.
- cysteine (sis'tein) n. [Gk. kystis. bladder.] A reduction product of cystine, occurring in urinary bladder concretions.
- cystenchyma (sistěng'kimă) n. [Gk. kystis, bladder; engchyma, infusion.] A parenchyma in sponges with large vesicular cell-structure.
- cystencytes (sis'tensits) n. plu. [Gk. kystis, bladder; en, in; kytos, hollow.] In sponges, collencytes which have acquired a vesicular structure.

- cystic (sis'tik) a. [Gk. kystis, bladder.] Pert. a cyst; pert. gall-bladder or to urinary bladder.
- cysticercoid (sis'tiser'koid) a. [Gk. kystis, bladder; kerkos, tail; eidos, form.] Appl. the bladderworm stage of tapeworms.
- cysticercus (sīs'tīsēr'kūs) n. [Gk. kystis, bladder; kerkos, tail.] The larval form or bladderworm stage of certain tapeworms.
- cysticolous (sīstīk'ölŭs) a. [Gk. kystis, bladder; L. colere, to inhabit.] Living in a cyst.
- cystid (sis'tid) n. [Gk. kystis, bladder; idion, dim.] A fossil Cystoid.
- cystidiform (sis'tidifôrm) a. [Gk. kystis, bladder; idion, dim.; I. forma, form.] Appl. clavate cells on gill margins in agarics.
- cystidium (sistid'iŭm) n. [Gk. kystis, bladder; idion, dim.] A hair-like inflated cell in the hymenial layer of some fungi.
- cystine (sīs'tēn) n. [Gk. kystis, bladder.] Amino acid found in plants, egg albumin and keratin; dicysteine, C₆H₁₂O₄N₂S₂.
- cystoarian (sīstōā'rīān) a. [Gk kystis, bladder; oarton, small egg.] Appl. gonads when enclosed in coelomic sacs, as in most teleosts. Opp. gymnoarian.
- cystocarp (sis'tokârp) n. [Gk. kystis, bladder; karpos, fruit.] A cyst arising from carpogonial branch and containing spores, in certain Rhodophyceae; cryptocarp.
- cystocyte (sīs'tösīt) n. [Gk. kystis, bladder; kytos, hollow.] Cystencyte, q.v.
- cystogenous (sistŏj'čnŭs) a. [Gk. kystis, bladder; -genes, producing.] Cyst-forming; appl. large nucleated cells which secrete the cyst, in cercaria.
- cystolith (sis'tölith) n. [Gk. kystis, bladder; lithos, stone.] A mass of calcium carbonate, occasionally of silica, formed on ingrowths of epidermal cell walls in some plants; a vesical calculus.

- cyston (sĭs'tŏn) n. [Gk. kystis, bladder.] A dactylozooid modified for excretory purposes, in Siphonophora.
- (sĭs'töspōr) [Gk. cystospore n. kystis, bladder; sporos, seed.] A carpospore; a cyst containing zoospores.
- cystozooid (sis'töz \bar{o} 'oid) n. [Gk. kystis, bladder; zoon, animal; eidos, form.] The body portion of a metacestode, opp. acanthozooid.
- cytase (sī'tās) n. [Gk. kytos, hollow.] A cellulose digesting or hydrolysing enzyme; alexin.
- cytaster (sītăs'tër) n. [Gk. kytos, hollow; aster, star.] A star-shaped achromatinic figure consisting of attraction-sphere and aster rays; aster, opp. karyaster.
- cytes (sīts) n. plu. [Gk. kytos, hollow.] Spermatocyte and oocyte stages of germ-cell formation; auxocytes.
- cytoblast (sī'töblăst) n. [Gk. kytos, hollow; blastos, bud.] The cell nucleus; a hypothetical unit, q.v.
- cytoblastema (sī'töblās'tēmā) n. [Gk. kytos, hollow; blastema, growth.] formative material from which cells were supposed to
- cytocentrum (sī'tösĕn'trŭm) n. [Gk. kytos, hollow; kentron, centre.] Centrosome; idiozome.
- cytochrome (sī'tokrōm) n. [Gk. kytos, hollow; chroma, colour.] A chromoprotein essential for oxidation-reduction processes in plant and animal cells; histohaematin; myohaematin.
- cytochylema (sī'tökīlē'mă) n. [Gk. kytos, hollow; chylos, juice.] Cytolymph, q.v.
- cytoclesis (sī'töklē'sĭs) n. [Gk. kytos, hollow; klesis, summons.] influence of a cell group or placode upon development or differentiation of neighbouring cells; cf. organiser.
- cytococcus (sī'tökŏk'ŭs) n. [Gk. kytos, hollow; kokkos, kernel.] The nucleus of a fertilised egg. cytocyst (sī'tösĭst) n. [Gk. kytos,

kystis, bladder.] The hollow: envelope formed by remains of host-cell within which a protozoan parasite multiplies.

cytode (sī'tōd) n. [Gk. kytos, hollow; A non - nucleated eidos, form.]

protoplasmic mass.

cytoderm (sī'tödērm) n. [Gk. kytos, hollow; derma, skin.] A cell-wall. cytodiaeresis (sī'tödīē'rēsīs) n. [Gk. kytos, hollow; diairesis, division.]

Mitosis; karyokinesis.

- cytogamy (sītog'āmi) n. [Gk. kytos, hollow; gamos, marriage.] Cellconjugation.
- cytogenesis (sī'töjěn'ësĭs) n. kytos, hollow; genesis, descent.] Development or formation of cells.
- cytogenetic (sī'töjĕnĕt'īk) a. kytos, hollow; genesis, descent.] cytogenesis; pert. genetics; appl. map showing location of genes within a chromosome.
- cytogenetics (sī'töjēnēt'iks) n. [Gk. kytos, hollow; genesis, descent.] Genetics in relation to cytology; the cytological aspect of genetics.
- cytogenous (sītoj'ėnus) a. kytos, hollow; genos, offspring.] Producing cells; appl. lymphatic tissue.
- cytoglobin (sī'töglō'bĭn) n. kytos, hollow; L. globus, globe.] A protein which retards coagulation of blood.
- cytohyaloplasma (sī'töhī'ālöplāz'mā) [Gk. kytos, hollow; hyalos, glass; plasma, mould.] The substance of the cytomitome; hyaloplasm.
- cytokinesis (sī'tökīnē'sīs) n. kytos, hollow; kinesis, movement.] Changes attending general cytoplasm during karyokinesis; the separation of daughter-cells following division of parent cell.
- **cytology** (sītŏl'öjĭ) n. [Gk. kytos, hollow, hollow vessel; logos, discourse.] The science dealing with structure, functions, and life-history of cells.
- cytolymph (sī'tölimf) n. [Gk. kytos, hollow; L. lympha, water.] Cellsap; the fluid part of cytoplasm.

- cytolysin (sī'tŏl'īsīn) n. [Gk. kytos, hollow; lysis, loosing.] A substance inducing cytolysis.
- cytolysis (sītōl'īsīs) n. [Gk. kytos, hollow; lysis, loosing.] Cell-dissolution; cell-degeneration.
- cytome (sī'tōm) n. [Gk. kytos, hollow.] The microsome or cytosome system of a cell.
- cytomeres (sī'tömērz) n. plu. [Gk kytos, hollow; meros, part.] Cells in Caryotropha formed by division of schizont and giving rise to merozoites; agametoblasts; nonnuclear portions of sperms.
- cytomicrosome (sī'tömī'krösōm) n. [Gk. kytos, hollow; mikros, small; soma, body.] A microsome of cytoplasm, opp. karyomicrosome.

cytomitome (sī'tömī'tōm) n. [Gk kytos, hollow; mitos, thread.] The cytoplasmic thread-work.

- cytomorphosis (sī'tömôr'fōsīs) n. [Gk. kytos, hollow; morphosis, shaping.] The life-history of cells; the series of structural modifications of cells or successive generations of cells; cellular change, as in senescence.
- cyton (sī'tŏn) n. [Gk. kytos, hollow.] The body of a nerve cell; neurocyton.
- cytophan (sī'töfān) n. [Gk. kytos, hollow; phaneros, visible.] Ovoid matrix surrounding karyophans in spironeme and axoneme fibres in infusorian stalk.
- cytopharynx (sī'töfăr'ingks) n. [Gk. kytos, hollow; pharyngx, gullet.]
 A tube-like structure leading from mouth into endoplasm in certain protozoa.
- cytophil (sī'töfil) a. [Gk. kytos, hollow; philein, to love.] Pert. haptophorous groups; having an affinity for cells.
- cytophore (sī'töför) n. [Gk. kytos, hollow; phora, burden.] A cell regarded as bearer of parasitic Sporozoa; central non-nucleated protoplasm in sperm morula; blastophore.
- cytoplasm (sī'töplazm) n. [Gk. kytos, hollow; plasma, mould.] Sub-

- stance of cell-body exclusive of nucleus; cf. karyoplasm.
- cytoproct (sī'töprökt), cytopyge (sī'töpīj') n. [Gk. kytos, hollow; proktos, anus; pyge, rump.] A cell-anus.
- cytoreticulum (sī'törētīk'ūlūm) n. [Gk. kytos, hollow; L. reticulum, little net.] The cytoplasmic threadwork; cytomitome; spongioplasm.
- cytosine (sī'tösīn) n. [Gk. kytos, hollow.] A cleavage product of nucleic acid; C₄H₆N₃O.
- cytosome (sī'tösōm) n. [Gk. kytos, hollow; soma, body.] The cytoplasmic part of a cell; a microsome, q.v.
- cytostome (sī'töstōm) n. [Gk. kytos, hollow; stoma, mouth.] A cellmouth.
- cytotaxis (si'tötäk'sis) n. [Gk. kytos, hollow; taxis, arrangement.] Rearrangement of cells on stimulation cytothesis (si'tăth'ăi's) n. [Gk.
- cytothesis (sī'tŏth'ĕsĭs) n. [Gk. kytos, hollow; thesis, arranging.]
 Regenerative tendency of a cell.
- cytotoxin (sī'tötŏk'sĭn) n. [Gk. kytos, hollow; toxikon, poison.] A cellpoisoning substance formed in blood serum; cytolysin; enzymoid.
- cytotrophoblast (sī'tötröf'ŏblăst) n. [Gk. kytos, hollow; trophe, nourishment; blastos, bud.] Inner layer of trophoblast, layer of Langhans.
- cytotropism (sitŏt'röpizm) n. [Gk. kytos, hollow; trope, turning.] The mutual attraction of two or more cells.
- cytozoic (sī'tözō'ĭk) a. [Gk. kytos, hollow; zoon, animal.] Living within a cell; appl. sporozoan trophozoite.
- cytula (st'ūla) n. [Gk. kytos, hollow.] The fertilised ovum or parent cell.

D

- dacryocyst (dăk'riösist) n. [Gk. dakryon, tear; kystis, bladder.]
 Lacrimal sac; saccus lacrimalis.
- dacryon (dăk'riŏn) n. [Gk. dakryon, tear.] Point of junction of anterior border of lacrimal with frontal bone and frontal process of maxilla.

- dactyl (dăk'til) n. [Gk. daktylos, finger.] A digit; finger, or toe; terminal ventral projection of praetarsus in scorpions; dactylus.
- dactylar (dăk'tilăr) a. [Gk. daktylos, finger.] Pert. finger or digit.
- dactylognathite (däk'tilönåth'īt) n. [Gk. daktylos, finger; gnathos, jaw.] Terminal segment of a maxillipede.
- dactyloid (dăk'tiloid) a. [Gk. daktylos, finger; eidos, form.] Like a finger.
- dactylopatagium (dăk'tilöpătăjī'ŭm)

 n. [Gk. daktylos, finger; L.
 patagium, border.] Ectopatagium,
- dactylopodite (dăk'tilŏp'ödīt) n. [Gk. daktylos, finger; pous, foot.] Distal joint in certain limbs of Crustacea; metatarsus and tarsus, of spiders.
- dactylopore (dăk'tilöpōr') n. [Gk. daktylos, finger; poros, channel.]
 Opening in skeleton of Milleporina, for protrusion of a dactylozooid.
- dactylopterous (dăk'tilop'tërŭs) a. [Gk. daktylos, finger; pteron, wing.] With anterior rays of pectoral fins more or less free.
- dactylozooid (dăk'tilözō'oid) n. [Gk. daktylos, finger; zoon, animal; eidos, form.] A hydroid modified for catching prey, long, with tentacles or short knobs, with or without a mouth.
- dactylus (dăk'tīlūs) n. [Gk. daktylos, finger.] Part of tarsus of an insect; dactyl of scorpions.
- dart (dârt) n. [O.F. dart, dagger.] Any structure resembling a dart; appl. a crystalline structure in molluscs.
- dart sac,—a small sac, containing a limy dart, attached to vagina near its orifice in some gastropods.
- dartoid (dâr'toid) a. [Gk. dartos, flayed.] Pert. the dartos.
- dartos (dâr'tos) n. [Gk. dartos, flayed.] Tunica dartos, a thin layer of non-striped muscle united to skin of scrotum.
- Darwinian tubercle,—the slight prominence on helix, of external

- ear, near the point where it bends downwards.
- Darwinism (Dâr'wĭnīzm) n. [C. Darwin]. The theory of origin of species by natural selection working on slight variations that occur, thereby selecting those best adapted to survive.
- dasypaedes (dăs'îpē'dēz) n. plu. [Gk. dasys, hairy; pais, child.] Birds whose young are downy at hatching.
- dasyphyllous (dās'īfīl'us) a. [Gk. dasys, hairy; phyllon, leaf.] With thickly haired leaves.
- dauermodification (dow'ermodifikā'shūn) n. [Ger. Dauer, duration; L. modificatio, modification.] A change induced by environmental factors and persisting for several generations but not permanently, the organism eventually reverting to type.
- daughter (dô'ter) n. [A.S. dohtor, daughter.] Offspring of first generation with no reference to sex, as daughter-cell, daughter-nucleus, etc.; daughter-chromosome: a chromatid during anaphase.
- day-neutral,—appl. plants in which flowering can be induced by either a long or a short photoperiod. Cf. long-day, short-day.
- dealation (dēālā'shūn) n. [L. de, away; alatus, winged.] The removal of wings, as by female ants after fertilisation.
- deamination (deam'inashun) n. [L. de, down; Gk. ammoniacum, resinous gum.] The conversion of ammonium salts into urea, partly accomplished in the liver.
- death (děth) n. [A.S. deáth, death.] Complete and permanent cessation of vital functions in an organism.
- death-point,—temperature above or below which organisms cannot exist.
- **Débove's membrane** [M. G. Débove, French histologist]. Layer between tunica propria and epithelium of tracheal, bronchial, and intestinal mucous membranes; subepithelial endothelium.

decalcify (dēkāl'sīfī) v. [L. de, away; calx, lime; facere, to make.] To deprive of lime salts; to treat with acids for removal of calcareous part.

decamerous (děkăm'ërus) a. [Gk. deka, ten; meros, part.] With the various parts arranged in tens.

decandrous (děkăn drus) a. [Gk. deka, ten; aner, male.] Having ten stamens.

decaploid (děk'áploid) a. [Gk. deka, ten; aploos, onefold; eidos, form.] Having ten times the haploid number of chromosomes.

decapod (děk'ápŏd) a. [Gk. deka, ten; pous, foot.] Of Crustacea, with five pairs of legs on thorax; of Cephalopoda, with ten arms.

decapodiform (děk'ăpŏd'ĭfôrm) a. [Gk. deka, ten; pous, foot; L. forma, shape.] Resembling a decapod, appl. insect larvae.

decemfid (desem'fid) a. [L. decem, ten; findere, to cleave.] Cut into

ten segments.

decemfoliate (děs'ěmfō'liāt) a. [L. decem, ten; folium, leaf.] Tenleaved.

decemjugate (děsěm'joogāt) a. [L. decem, ten; jugare, to join.] With ten pairs of leaflets.

decempartite (děs'ěmpâr'tīt) a. [L. decem, ten; partiri, to divide.]

Ten-lobed; divided into to lobes.

decidua (dēsīd'ūā) n. [L. decidere, to fall off.] The mucous membrane lining the pregnant uterus, cast off

after parturition.
decidual (dēsĭd'ūăl) a. [L. decidere,

to fall off.] Pert. decidua. deciduate (dēsid'ūāt) a. [L. decidere, to fall off.] Characterised by having a decidua; partly formed by the decidua.

deciduous (dēsīd'ūūs) a. [L. decidere, to fall down.] Falling at end of growth period or at maturity.

declinate (děk lināt) a. [L. de, away; clinare, to bend.] Bending aside in a curve, as anther filament in horse-chestnut.

decollated (dēkŏl'ātëd) a. [L. de, away from; collum, neck.] With apex of spire wanting.

decomposed (dē'kŏmpōzd') a. [L. de, away; cum, with; pausare, to rest.] Not in contact; not adhering, said of barbs of feather when separate; decayed.

decompound (de 'kompound') a. [L. de, away; cum, with; ponere, to place.] With monopodial branching very complete, and ultimate wings little developed; appl. leaf.

deconjugation (dē'kŏnjoogā'shūn) n.
[L. de, away from; conjugare, to join together.] Separation of paired chromosomes, as before end of meiotic prophase.

decorticate (dēkôr'tīkāt) v. [L. decorticare, to peel.] To remove bark or cortex. a. With cortex

ablated

decumbent (dēkŭm'bënt) a. [L. decumbere, to lie down.] Appl. stems which trail on ground, but rise at apex.

decurrent (dēkur'ënt) a. [L. de, away; currere, to run.] Having leaf base prolonged down stem as a winged expansion or rib; prolonged down stipe, as gills of agaric.

decussate (dëkŭs'āt) a. [L. decussare, to cross.] Crossed; having paired leaves, succeeding pairs crossing at right angles.

decussation (děk'úsā'shǔn) n. [L. decussare, to cross.] Decussate condition of leaves; crossing of nerves with interchange of fibres, as in optic and pyramidal tracts.

dedifferentiation (dēdif'erenshīa'shūn) n. [L. de, away from; differentia, difference.] The losing of characteristics of specialised cells and regression to a more simple state.

dédoublement (dā'dooblëmông') n.
[F. dédoublement, dividing into two.] Chorisis; deduplication.

deduplication (dēdū'plikā'shūn) n.
[L. de, by reason of; duplicare, to double.] Chorisis, q.v.

- defaecation (dē'fēkā'shun) n. [L. defaecatio, voiding of excrement.]
 The expulsion of faeces.
- defensive (dëfën'siv) a. [L. defendere, to defend.] Protective; appl. proteid substances which destroy toxic substances of bacteria; appl. numerous organs or parts of organs in various animals or plants.

deferent (def'erent) a. [L. deferre, to carry away.] Conveying away; appl. ducts, vasa deferentia, q.v.

deferred (deferd') a. [L. deferre, to carry off.] Appl. shoots arising from dormant buds.

deficiency,—inactivation or absence of a chromosomal segment or gene.

- deficiency diseases, pathological conditions in plants and animals, due to lack of certain necessary nutritive substances; e.g. crown rot in sugar beet due to boron deficiency; diseases in mammals due to vitamin deficiency, absence of vitamin A causing poor growth and xerophthalmia,—of B₁, beriberi,—of B₂, retardation of growth,—of C, scurvy,—of D, rickets,—of E, infertility and paralysis,—of K, bleeding,—etc.
- definite (def'init) a. [L. definire, to limit.] Fixed, constant; cymose, appl. inflorescences with primary axis terminating early in a flower; appl. stamens limited to twenty in number.
- definitive (dëfin'îtĭv) a. [L. definire, to limit.] Defining or limiting; complete, fully developed; final, appl. host of adult parasite.

defoliate (dēfo'liāt) a. [L. defoliare, to strip of leaves.] Bared at the annual fall. v. To deprive of leaves.

- degeneration (dējěněrā'shun) n. [L. degenerare, to degenerate.] Change to a less specialised or functionally less active form; retrogressive evolution.
- deglutition (dēglootish'ŭn) n. [L.
 de, down; glutire, to swallow.]
 The process of swallowing.

dehiscence (dēhīs'ens) n. [L. de-

- hiscere, to gape.] The spontaneous opening of an organ or structure along certain lines or in a definite direction.
- deinopore (dī'nöpōr) n. [Gk. deinos, urn; poros, channel.] A cell bridge.
- deirids (dī'rīdz) n. plu. [Gk. deiras, chain of hills.] Cervical papillae in Nematoda.
- Deiters' cells [O. F. C. Deiters, German anatomist]. Supporting cells between rows of outer haircells in organ of Corti; outer phalangeal cells.

delamination (dēlām'ĭnā'shŭn) n. [L. de, down; lamina, layer.] The dividing off of cells to form new layers.

deletion (dēlē'shun) n. [L. delere, to efface.] A deficiency of an acentric part of chromosome; absence of a chromosome segment and of the genes involved.

deliquescent (děl'íkwěs'ënt] a. [L. deliquescere, to become fluid.] Having lateral buds the more vigorously developed, so that the main stem seems to divide into a number of irregular branches; becoming fluid.

delitescence (dělitěs'ëns) n. [L. delitescere, to lie hidden.] The latent period of a poison; incubation period of a pathogenic organism.

delomorphic (děl'ömôr'fĭk) a. [Gk. delos, visible; morphe, shape.] With definite form, appl. oxyntic cells of the gastric glands; delomorphous.

delthyrium (dělthī'rĭŭm) n. [Gk. delos, visible; thyrion, little door.] The opening, between hinge and beak, for peduncle exit in many Brachiopoda.

deltidium (děltĭd'ĭŭm) n. [Gk. Δ, delta; idion, dim.] A plate covering the delthyrium.

deltoid (děl'toid) a. [Gk. Δ, delta; eidos, form.] More or less triangular in shape, appl. muscle, etc.; appl. oral plates on calyx of Blastoidea.

- demanian (deman'ian) a. [J. G. de Man, French zoologist]. Appl. a complex system of paired efferent tubes connecting with intestine and uteri in Nematoda, and associated with gelatinous secretion for protection of eggs.
- deme (dēm) n. [Gk. demos, people.] Assemblage of taxonomically closely related individuals; aggregate of single cells.
- demersal (dēmĕr'săl) a. [L. demergere, to plunge into.] Living on or near bottom of sea or lake; sunk.
- demersed (dēměr'sd) a. [L. demergere, to plunge into.] Growing under water; appl. parts of plants. demibranch,—hemibranch, q.v.
- demifacet (demifas'et) n. [L. dimidius, half; facies, face.] Part of parapophysis facet when divided between centra of two adjacent vertebrae.
- demilunes,—crescentic cells; crescentic bodies of cells of some salivary gland alveoli, crescents of Gianuzzi or demilunes of Heidenhain.
- demiplate (dem'iplat) n. [L. dimidius, half; platus, flat.] Plate cut off by fusion of adjoining plates behind it from central suture line of ambulacral area in echinoderms.
- demisheath (dēm'īshēth) n. [L. dimidius, half; A.S. sceath, sheath.]
 One of paired protecting covers of insect ovipositor.
- demoid (de'moid) a. [Gk. demos, the commons.] Abundant.
- denatant (dēnā'tānt) a. [L. de, down from; natare, to swim.] Swimming, drifting, or migrating with the current, opp. contranatant.
- dendriform (děn'drifôrm) a. [Gk. dendron, tree; L. forma, shape.]
 Dendroid; tree-like.
- dendrite (děn'drīt) n. [Gk. dendron, tree.] A fine branch of a dendron; a dendron.
- dendritic (děndrit'ik) a. [Gk. dendron, tree.] Dendroid; appl. tree-like structures or markings; like, pert., or having, dendrites or dendrons.
- dendrochronology (děn'drökrönŏl'

- oji) n. [Gk. dendron, tree; chronos, time; logos, discourse.] Determination of age of trees or timber; dating by comparative study of tree rings; science of tree-ring analysis and its implications.
- Dendrogaea (děn'drojē'á) n. [Gk. dendron, tree; gaia, earth.] A zoogeographical region including all the neotropical region except temperate South America.
- dendroid (děn'droid) a. [Gk. dendron, tree; eidos, form.] Tree-like; much branched.
- dendrology (děndrol'oji) n. [Gk. dendron, tree; logos, discourse.]
 The study of trees.
- dendron (děn'dron) n. [Gk. dendron, tree.] A protoplasmic process of the nerve cell, which conducts impulses towards the cell body.
- denitrification (dē'nītrīfīkā'shūn) n. [L. de, away; Gk. mtron, soda; L. facere, to make.] Reduction of nitrates, to nitrites and ammonia, as in plant tissues, or to molecular nitrogen, as by certain soil bacteria.
- dens (děnz) n. [L. dens, tooth.]
 Tooth, or tooth-like process; odontoid process of axis or epistropheus.
 dens serotinus a the third molar
- dens serotinus, the third molar or wisdom-tooth.
- dental (děn'tăl) a. [L. dens, tooth.] Pert. teeth; appl. nerves, bloodvessels, canals, furrows, papillae, sac, tissue, etc.
- dentary (děn'tări) a. [L. dens, tooth.]

 Pert. dentaries, membrane bones in lower jaw of many vertebrates. n.

 Dentary bone or os dentale.
- dentate (den'tat) a. [L. dens, tooth.]
 Toothed; with sharp saw-like teeth
 on the margin.
- dentate ciliate,—with teeth and hairs on the margins; appl. leaves.
- dentate crenate,—with marginal teeth somewhat rounded.
- denticles (den'tiklz) n. plu. [L. denticulus, little tooth.] Small tooth-like processes; the paragnaths of certain Polychaeta; the teeth within the secondary orifice in Polyzoa; the scales of certain Elasmobranchii.

- denticulate (děntík'ūlāt) a. [I.. denticulus, little tooth.] Having denticles; with minute marginal teeth.
- dentin,—dentine, q.v.
- dentinal (děn'tǐnăl) a. [L. dens, tooth.] Pert. dentine; appl. tubules, i.e. canaliculi dentales.
- dentine (děn'tǐn) n. [L. dens, tooth.]
 A hard, elastic substance, chemically resembling bone, composing the greater part of teeth and denticles; dentin.
- dentition (děntish'ŭn) n. [L. dens, tooth.] The number, arrangement, and kind of teeth; teething.
- deperulation (dē'pērūlā'shun) n. [L. de, away; dim. of pera, wallet.] The pushing apart or throwing off, of bud scales.
- depigmentation (dēpig'mëntā'shūn)
 n. [L. de, away; pingere, to paint.]
 The destruction of colour in a cell, by natural or experimental physiological processes.
- depilation (dĕp'ilā'shŭn) n. [L. de, away; pilus, hair.] Loss of hairy covering, as of plants when maturing; removal of hair.
- deplanate (děp'lănāt) a. [L. deplanare, to level.] Levelled, flattened.
- depressant (dēpres'ant) n. [L. deprimere, to keep down.] Anything that lowers vital activity.
- depressomotor (dēpres'omō'tor) n.
 [L. deprimere, to keep down;
 movere, to move.] Any nerve which
 lowers muscular activity.
- depressor (dēprēs'ŏr) n. [L. deprimere, to keep down.] Any muscle which lowers or depresses any structure; appl. a nerve which lowers the activity of an organ.
- depula (děp'ūlă) n. [Gk. depas, goblet.] Invaginated blastula preceding gastrula stage in development of embryo.
- deric (děr'îk) a. [Gk. deros, skin.] Dermic; appl. epithelium, synonym of epidermis.
- derm (derm), derma (der'mă) n. [Gk. derma, skin.] The layers of integument below the epidermis; dermis.

- dermal (děr'măl) a. [Gk. derma, skin.] Pert. derma, or skin.
- dermalia (děrma'liă) n. plu. [Gk. derma, skin.] Microscleres in the dermal membrane of sponges.
- dermarticulare (děr'mârtik'ūlā'rē) n.
 [Gk. derma, skin; L. articulus, joint.] The goniale, q.v.
- dermatogen (der'matöjen) n. [Gk. derma, skin; genos, birth.] The young or embryonic epidermis in plants; antigen of skin disease.
- dermatoglyphics (děr'mătöglif'iks)

 n. [Gk. derma, skin; glyphein, to carve.] Skin, palm, finger, sole, toe prints; print formulae.
- dermatomes (děr'mătōmz) n. plu. [Gk. derma, skin; tome, cutting.] Lateral parts of segmental mesoderm, which develop into connective tissue of corium or dermis; dermatomere.
- dermatophyte (der'mătöfit) n. [Gk. derma, skin; phyton, plant.] Any fungous parasite of skin; dermatophyton, dermophyte, epidermophyte.
- dermatoplasm (dër mătöplăzm) n.
 [Gk. derma, skin; plasma, mould.]
 Cell-wall protoplasm.
- dermatoplast (děr'mătöplăst') n. [Gk. derma, skin; plastos, moulded.]
 A protoplast with a cell wall.
- dermatopsy (děr'mătŏp'si') n. [Gk. derma, skin; opsis, sight.] Condition of seeing with the skin, i.e. with a skin sensitive to light.
- dermatoskeleton,-exoskeleton.
- dermatosome (děr'mătösōm') n. [Gk. derma, skin; soma, body.] One of vital units forming a cell-membrane.
- dermatozoon (děr'mătözō'ŏn) n. [Gk. derma, skin; zoon, animal.] Any animal parasite of the skin; ectozoon.
- dermentoglossum (děrm'ěntöglős'ům) n. [Gk. derma, skin; entos, within; glossa, tongue.] A bone arising by fusion of dentinal bases, covering entoglossum, in some fishes.
- dermethmoid (děrměth'moid) n. [Gk. derma, skin; ethmos, sieve; ethmos, form.] Supra-ethmoid, q.v.
- dermic (děr'mik) a. [Gk. derma, skin.] Pert., or derived from, skin.

dermis (děr'mis) n. [Gk. derma, skin.] Derm, q.v.; corium.

DER-

- dermoblast (der'möbläst') n. [Gk. derma, skin; blastos, bud.] The layer of mesoblast which gives rise to the derma.
- dermoccipitals (derm'oksip'italz) n. plu. [Gk. derma, skin; L. occiput, back of head.] Two bones taking the place of interparietal in some lower forms and in development of higher.

dermo-ossification (der'möös'ifikā'-shun) n. [Gk. derma, skin; L. os, bone; fieri, to become.] A bone formed in the skin.

dermopharyngeal (der'möfarin'jeäl)

n. [Gk. derma, skin; pharyngx,
gullet.] Superior or inferior plate
of membrane bone supporting
pharyngeal teeth in some fishes.

dermophyte,—dermatophyte, q.v. dermosclerites (der'möskler'īts) n. plu. [Gk. derma, skin; skleros, hard.] Masses of spicules found in tissues of Alcyonidae.

dermoskeleton (děr'möskěl'etőn) n. [Gk. derma, skin; skeletos, dried.] Exoskeleton, q.v.

dermotrichia (děr'mötrĭk'ĭă) n. plu. [Gk. derma, skin; thrix, hair.] Dermal fin-rays.

dermozoon,—dermatozoon, q.v.

derotreme (der'ötrēm) n. [Gk. deros, skin; trema, aperture.] Skin forming an operculum, as in Megalobatrachus.

dertrotheca (děr'trothē'ka) n. [Gk. dertron, beak; theke, cup.] The horny casing of bird maxilla.

dertrum (der trum) n. [Gk. dertron, beak.] Any modification of the casing of maxilla in birds.

Descemet's membrane [J. Descemet, French anatomist]. The posterior elastic lamina of cornea; Demour's membrane.

descending (dēsěn'ding) a. [L. de, down; scandere, to climb.]
Directed downwards, or towards caudal region; appl. blood-vessels, nerves, etc.

desegmentation (dēsĕg'mëntā'shŭn)
n. [L. de, from; segmentum, piece

cut off.] Fusion of segments originally separate.

deserticolous (děz'ertik'ölüs) a. [L. desertus, waste; colere, to inhabit.] Desert-inhabiting.

desma (dĕs'mă, dĕz-) n. [Gk. desma, bond.] Megasclere which forms characteristic skeletal network of Lithistida.

desmactinic (děs'măktin'ik,) a. [Gk. desma, bond; aktis, ray.] With podia continued upwards to apical plate, appl. Stelleroidea; cf. lysactinic.

desmergate (des'mergat) n. [Gk. desma, bond; ergates, worker.] A type of ant intermediate between worker and soldier.

desmocyte (dĕs'mösīt) n. [Gk. desmos, bond; kytos, hollow.] A connective tissue cell; fibroblast.

desmogen (dës'möjën,) n. [Gk. desmos, bond; genos, descent.]
Merismatic or growing tissue.

desmognathous (děsmog'năthūs) a.

[Gk. desmos, bond; gnathos, jaw.]

Having maxillopalatines fused in middle line owing to other peculiarities in skull; appl. certain birds.

desmology (děsmŏl'öjĭ) n. [Gk. desmos, bond; logos, discourse.]
The anatomy of ligaments; cf. syndesmology.

desmones (des'monz) n. plu. [Gk. desmos, bond.] Chemical substances exchanged by way of protoplasmic bridges between cells; amboceptors, q.v.

desmose (děs'mōs) n. [Gk. desmos, bond.] A strand connecting ble-pharoplasts at mitosis.

desmosome (des'mösöm) n. [Gk. desmos, bond; soma, body.] A thickening of intercellular connections in epithelium; bridge corpuscle.

desquamation (děs'kwămā'shun) n. [L. de, away; squama, scale.] Shedding of cuticle or epidermis in flakes.

determinant (dētēr'mīnānt) n. [L. determinare, to limit.] A hypothetical unit, being an aggregation of biophores determining the

- development of a cell or of an independently variable group of cells; hereditary factor.
- determinate (dētěr'mīnāt) a. [L. determinare, to limit.] With certain limits; appl. inflorescence with primary axis terminated early with a flower-bud; appl. cleavage.
- determination (determina'shun) n.

 [L. determinatio, boundary.] The process adjusting regional development according to relative location of region and organisation centre.
- detorsion (dētôr'shun) n. [L. de, away; torquere, to twist.] Torsion in an opposite direction to that of original, resulting in a more or less posterior position of anus and circumanal complex.
- detoxication (dē'tōksīkā'shūn) n. [L. de, away; Gk. toxikon, poison.] The inhibition of effects of toxins in the body, either by a protective synthesis of comparatively harmless substances, or by means of antibodies.
- detrusor (dētroo'sŏr) n. [L. detrudere, to thrust from.] The outer of three layers of the muscular coat of the urinary bladder; physiologically, all three layers; detrusor urinae.
- deuter cell,—eurycyst, q.v. deutero-,—also see deuto-.
- deuterocerebrum(dū'teroser'ebrum)
 n. [Gk. deuteros, second; L. cerebrum, brain.] That portion of crustacean brain from which antennular nerves arise. Cf. deutocerebrum.
- deuterocoele (dū'tērösēl) n. [Gk. deuteros, second; koilos, hollow.]
 The coelom.
- deuterocone (dū'tërökōn') n. [Gk. deuteros, second; konos, cone.] Mammalian premolar cusp corresponding to molar protocone.
- deuterogenesis (dū'tēröjēn'ēsis) n. [Gk. deuteros, second; genesis, origin.] Second phase of embryonic development, involving growth in length and consequent bilateral symmetry; cf. protogenesis.

- deuteroplasm,—deutoplasm, q.v.
- deuteropolydesmic (dū'teröpolides' mik) a. [Gk. deuteros, second; desmos, bond.] Appl. cyclomorial scales composed mainly of synpolydesmic scales.
- deuteroproteose (dū'tēröpro'tēōs) n, [Gk. deuteros, second; protos, first.] A secondary product from digestion of proteids.
- deuterostoma (dū'teros'toma) n.
 [Gk. deuteros, second; stoma, mouth.] A mouth formed secondarily, as distinct from gastrula mouth.
- deuterotoky (dū'tērŏt'ökī) n. [Gk. deuteros, second; tokos, birth.] Reproduction of both sexes from parthenogenetic eggs; cf. arrhenotoky and thelyotoky.
- Deuterozoic (dū'terözō'ĭk) a. [Gk. deuteros, second; zoe, life.]
 Appl. and pert. the newer Palacozoic faunal epoch, the age of fishes, also of pteridosperms.
- deuterozooid (dū'tērözō'oid) n.
 [Gk. deuteros, second; zoon, animal; eidos, form.] A zooid produced by budding from a primary zooid.
- deuthyalosome (dūthī'ălösōm) n. [Gk. deuteros, second; hyalos, glass; soma, body.] The nucleus remaining in ovum after formation of first polar body.
- deuto-,-also see deutero-.
- deutoblasts (dū'töblăsts) n. plu. [Gk. deuteros, second; blastos, bud.] The amoeba-like bodies formed from protoblasts in zygote of Microclossia, and liberated to multiply in the blood.
- deutobroch (dū'töbrök) a. [Gk. deuteros, second; brochos, mesh.] Appl. nuclei of gonia preparing for leptotene stage; cf. protobroch.
- deutocerebrum (dū'tösĕr'ĕbrum) n. [Gk. deuteros, second; L. cerebrum, brain.] Portion of insect brain derived from fused ganglia of antennary segment of head; deutocerebron. Cf. deuterocerebrum.

deutomalae (dū'tömā'lē) n. plu. [Gk. deuteros, second; malon, cheek.] The broad plate in Chaetognatha, formed by fusion of second pair of mouth appendages.

deutomerite (dūtŏm'ērīt) n. [Gk. deuteros, second; meros, part.]

The posterior division of certain

gregarines; cf. primite.

deutonephros (dū'tonĕf'rŏs) n. [Gk. deuteros, second; nephros, kidney.]

Mesonephros.

- deutoplasm (dū'töplāzm) n. [Gk. deuteros, second; plasma, mould.] Yolk or food material in cytoplasm of ovum or other cell.
- deutoscolex (dū'töskō'lĕks) n. [Gk. deuteros, second; skolex, worm.] A secondary scolex produced by budding, in bladderworm stage of certain tape-worms.
- deutosomes (dū'tosōmz) n. plu. [Gk. deuteros, second; soma, body.]
 Granules of nucleolus cast out into cytoplasm, from which yolk is said to arise.
- deutovum (dūtō'vŭm) n. [Gk. deuteros, second; L. ovum, egg.]
 A stage in the metamorphosis of certain mites.
- development (dēvěl'ŏpmënt) n. [F. développer, to unfold.] The changes undergone by an organism from its beginning to maturity.

deviation (devia'shun) n. [L. de, away from; via, way.] Divergence from corresponding developmental

stages.

Devonian (dēvō'nĭān) a. [Devon, where strata were first studied.] Pert. or appl. Palaeozoic geological period preceding Carboniferous.

- dexiotropic (děk'siötrop'ik) a. [Gk. dexios, right; trope, turn.] Turning from left to right, as whorls; appl. shells; appl. spiral cleavage of cells; appl. movement of Volvox. dextral (děk'strál) a. [L. dexter,
- right-hand.] Dexiotropic, q.v. dextrin (děk'strin) n. [L. dexter, right-hand.] A soluble substance
- derived from starch by exposure to high temperature for a short time. dextrorse (dekstrôrs') a. [L. dexter,

- right; vertere, to turn.] Growing in a spiral which twines from left to right; clockwise. Opp. sinistrorse.
- dextrose (děk'strōs) n. [L. dexter, right.] Grape sugar or glucose, the end product of starch digestion, C₆H₁₂O₆.
- diabetogenic (dīābē'töjēn'īk) a. [Gk. diabainein, to cross over; gignesthai, to beget.] Causing diabetes; appl. a prepituitary hormone antagonistic to insulin, affecting carbohydrate metabolism; appl. a hormone of sinus gland of eye stalk in crustaceans.
- diachaenium (dī'ākē'nĭŭm) n. [Gk. dis, twice; a, not; chainein, to gape.] Each part of a cremocarp.
- diachronous (dīāk'rönūs) a. [Gk. dia, asunder; chronos, time.] Dating from different periods; appl. fossils occurring in the same geological formation, though in different areas, due, e.g., to changes in sea-level.
- diachyma (diāk'imā) n. [Gk. dia, throughout; chymos, juice.] Leaf parenchyma.
- diacoel (dī'āsēl) n. [Gk. dia, through; koilos, hollow.] Third ventricle of brain.
- diacranteric (di'ákrăntěr'ík) a. [Gk. dia, asunder; kranteres, wisdom teeth.] With diastema between front and back teeth, as in snakes.
- diactinal (diāk'tināl) a. [Gk. dis, twice; aktis, ray.] With two rays pointed at ends.
- diadelphous (di'adel'fus) a. [Gk. dis, twice; adelphos, brother.] Having stamens in two bundles owing to fusion of filaments.
- diadematoid (dī'ādēm'ātoid) a. [Gk. diadema, crown; eidos, shape.] Of Echinoidea, having three primary pore plates with occasionally a secondary between aboral and middle primary; as opp. arbacioid, one primary, with secondary on each side, and triplechinoid, two primaries, with one or more secondaries between.

- diadromous (dīād'römus) a. [Gk. diadromos, wandering.] Having nerves or veins radiating in fan-like manner; appl. leaves.
- diaene (di'ēn) n. [Gk. dis, twice; an analogy of triaene, from Gk. triaina, trident.] A form of triaene, with one of the cladi reduced or absent.
- diageotropism (dī'ājēŏt'röpĭzm) n. [Gk. dia, through; ge, earth; trope, turn.] Tendency in certain parts of plants to assume position at right angles to direction of gravity.
- diagnosis (dī'agnō'sīs) n. [Gk. diagnosis, discrimination.] A concise description of an organism with full distinctive characters; discrimination of a physiological or pathological condition by its distinctive signs.
- diagnostic (dī'āgnŏs'tīk) a. [Gk. diagnosis, discrimination.] Distinguishing; differentiating the species or genus, etc., from others similar.
- diaheliotropism (dī'āhēliŏt'röpizm)
 n. [Gk. dia, through; helios, sun;
 trope, turn.] Diaphototropism.
- diakinesis (dī'ākinē'sīs) n. [Gk. dia, through; kinesis, movement.] The later prophase stage of meiosis, between diplotene and prometaphase; movement of chromosomes between metaphase and telophase.
- dialyneury (dī'ālinū'rī) n. [Gk. dia-lyein, to reconcile; neuron. nerve.] In certain gastropods, condition of having pleural ganglia united to opposite visceral nerve by anastomosis with pallial nerve.
- dialypetalous (dī'ălīpēt'ălŭs) a. [Gk. dia, asunder; lyein, to loose; petalon, petal.] Polypetalous.
- dialyphyllous (di'ālifil'us) a. [Gk. dia, asunder; lyein, to loose; phyllon, leaf.] With separate leaves. dialysepalous (di'ālīsēp'ālus) a. [Gk. dia, asunder; lyein, to loose; F. sépale, sepal.] Polysepalous.
- dialysis (dīāl'īsis) n. [Gk. dia, asunder; lysis, loosing.] Separation of dissolved crystalloids and colloids through semipermeable

membrane, crystalloids passing more readily; permeation.

- dialystely (dī'ālīstē'lĭ) n. [Gk. dia, asunder; lyein, to loose; stele, post.] A condition in which the steles in the stem remain more or less separate.
- diancistron (di'ănsis'tron) n. [Gk. dis, twice; angkistron, hook.] A spicule resembling a stout sigma, but the inner margin of both hook and shaft thins out to a knife edge and is notched. Plu. diancistra.
- diandrous (dian'drus) a. [Gk. dis, twice; aner, man.] Having two free stamens.
- diapause (dī'āpôz) n. [Gk. diapauein, to make to cease.] A spontaneous state of dormancy during development, as of insects; resting stage between anatrepsis and katatrepsis in blastokinesis; sexual rest period, appl. annelids; cf. quiescence.
- diapedesis (dī'āpēdē'sīs) n. [Gk. diapedesis, leaping through.] Emigration of white blood corpuscles through walls of capillaries into surrounding tissue; migration of cells to exterior, in certain larval sponges.
- diaphototropism (di'afōtōt'röpizm)
 n. [Gk. dia, through; phos, light;
 trope, turn.] Tendency of plant
 organs to assume a position at right
 angles to rays of light; diaheliotropism.
- diaphragm (dī'afram), diaphragma (dī'āfrāg'mā) n. [Gk. diaphragma, midriff.] The wall which separates the small cell, the prothallus, from rest of macrospore in Hydropterideae; a septum at nodes in Equisetum; a sheet of muscular tissue attached to introvert in worms; single strongly developed septum in Terebelliformia; perforated tissue that subdivides tentacle cavity in Polyzoa; fibro-muscular abdominal septum enclosing perineural sinus in certain insects; the transverse septum separating cephalothorax from abdomen in certain Arachnida; a

- special fan-shaped muscle spreading from anterior end of ilia to oesophagus and base of lungs in Anura; a partition partly muscular, partly tendinous, separating cavity of chest from abdominal cavity in mammals; fold of dura mater on sella turcica.
- diaphysis (dīăf'isis) n. [Gk. dia, through; phyein, to bring forth.] Shaft of limb bone, opp. epiphysis; abnormal growth of an axis or shoot.
- **diapophysis** ($d\bar{i}'\bar{a}p\bar{o}f'\bar{i}s\bar{i}s$) n. [Gk. dia, through; apo, from; phyein, to produce.] Lateral or transverse process of neural arch.

diapsid (dīăp'sid) a. [Gk. dis, twice; apsis, arch.] Appl. skulls with supra- and infra-temporal fossae distinct; cf. synapsid.

- diarch (dī'ark) a. [Gk. dis, twice; arche, origin.] With two xylem and two phloem bundles; appl. root in which protoxylem bundles meet and form a plate of tissue across cylinder with phloem bundle on each side; appl. a bipolar type of spindle.
- diarthric (diar'thrik) a. [Gk. dis, twice; arthron, joint.] Pert. two joints; diarticular.
- diarthrosis (dī'arthro'sĭs) n. dis, twice; arthron, joint.] articulation allowing considerable movement.
- diaschistic (dī'āskīs'tīk) a. dia, through; schistos, split.] Appl. type of tetrads which divide once transversely and once longitudinally in meiosis; cf. anaschistic.
- diaspore (dī'aspor) n. [Gk. diaspora, dispersion (dia, asunder; spora, seed.)] Any spore, seed, fruit, or other portion of a plant when being dispersed and able to produce a new plant; disseminule, propagule.
- diastase (dī'āstās) n. [Gk. diastanai, to separate.] An enzyme which acts principally in converting starch into sugar.
- diastasis (dias'tasis) n. [Gk. dia, through; stasis, halt.] Rest period preceding systole.

diastatic (dī'āstāt'īk) a. [Gk. dia, through; histanai, to set.] Pert. diastase, or having similar properties; pert. diastasis.

diastem (dī'āstěm), diastema (dīăs'tēmă) [Gk. diastema. n. A toothless space usually space.] between two types of teeth; an equatorial modification of protoplasm preceding cell division.

diaster (dīās'ter) n. [Gk. dis, twice; aster, star.] The stage in mitosis where daughter chromosomes are grouped near spindle poles ready to form a new nucleus.

diastole (dīăs'tölē) n. [Gk. diastole,

difference.] Rhythmical relaxation of heart; rhythmical expansion of a contractile vacuole. Opp. systole.

diastomatic (dī'āstömāt'īk) a. [Gk. through; stoma, mouth.] Through stomata or pores; giving off gases from spongy parenchyma through stomata.

diathesis (dīāth'ēsīs) n. [Gk. diathesis, disposition.] A constitutional predisposition to a type of reaction, disease, or development.

- diatom (dī'ātŏm) n. [Gk. dia, through; temnein, to cut.] A unicellular form of alga with walls impregnated with silica.
- diatomin (diat'ömin) n. [Gk. diatemnein, to cut through.] A yellow pigment resembling fucoxanthin, in plastids of diatoms.
- diatropism (dīāt'röpizm) n. [Gk. dia, through; trope, turn.] tendency of organs or organisms to place themselves at right angles to line of action of stimulus.
- diaxon (dīāk'sŏn) a. [Gk. dis, twice; axon, axis.] With two axes, as certain sponge spicules.
- diaxone (diăk'son) n. [Gk. dis, twice; axon, axis.] A nerve-cell with two axis-cylinder processes.
- diblastula (dīblăs'tūlă) n. [Gk. dis, twice; blastos, bud.] A coelenterate embryo consisting of two layers arranged round a central cavity.
- dibranchiate (dibrang'kiāt) a. [Gk. dis, twice; brangchia, gills.] With two gills.

dicaryon, -dikaryon.

dicellate (di'sĕlāt) a. [Gk. dikella, two-pronged hoe.] With two prongs; appl. sponge spicules.

dicentral (disen'tral) a. [Gk. dia, through; kentron, centre.] Appl. canal in fish vertebral centrum.

- dicentric (dīsĕn'trīk) a. [Gk. dis, twice; kentron, centre.] Having two centromeres; appl. chromatids, chromosomes.
- dichasium (dīkā'zĭŭm) n. [Gk. dichazein, to divide in two.] A cymose inflorescence in which two lateral branches occur about same level.
- dichlamydeous (dī'klāmīd'eŭs) a. [Gk. dis, twice; chlamys, cloak.] Having both calyx and corolla.
- dichogamy (dikög'āmī) n. [Gk. dicha, in two; gamos, marriage.]
 Maturing of sexual elements at different times, ensuring crossfertilisation; cf. protandry, protogyny.

dichoptic (dǐkŏp'tĭk) a. [Gk. dicha, in two; opsis, sight.] With eyes

quite separate.

dichotomous (dikŏt'ömŭs) a. [Gk. dicha, in two; temnein, to cut.] Pert., characterised by dichotomy.

- dichotomy (dǐkot'omi) n. [Gk. dicha, in two; temnein, to cut.] Branching which results from division of growing point into two equal parts.
- dichroic (dikrō'ik) a. [Gk. dis, twice; chros, colour.] Exhibiting dichroism, as chlorophyll solution; cf. dichromatic.
- dichroism (dī'krōĭzm) n. [Gk. dis, twice; chros, colour.] Property of showing two colours, as one colour by transmitted and the other by reflected light.

dichromatic (di'krōmāt'ik) a. [Gk. di-, two; chroma, colour.] With two colour varieties; seeing only two colours.

dichromophil (dikrōm'öfil) a. [Gk. di-, two; chroma, colour; philein, to love.] Staining with both acid and basic dyes.

dichthadiigyne (dikthăd'iijīnē) n.

[Gk. dichthadios, double; gyne, female.] A gynaecoid ant with voluminous ovaries, and without eyes and wings; a dichthadiiform female.

diclinous (dī'klĭnūs, dīklī'nūs) a. [Gk. di-, asunder; kline, bed.] With stamens and pistils on separate flowers; with staminate and pistillate flowers on same plant.

dicoccous (dīkŏk'ūs) a. [Gk. di-, two; kokkos, kernel.] Having two one-seeded coherent capsules.

dicostalia (dī'kŏstā'lĭā) n. [Gk. di-, two; L. costa, rib.] The secundibrachs or second brachial series in a crinoid.

dicotyledon (dikŏtĭlē'dŏn) n. [Gk. di-, two; kotyledon, cup-shaped hollow.] A plant with two seed-leaves.

dictyodromous (dĭk'tiŏd'römŭs) a. [Gk. diktyon, net; dromein, to run.] Net-veined, when the smaller veins branch and anastomose freely.

dictyogen (dĭk'tĭöjĕn) n. [Gk. diktyon, net; -genes, producing.] A net-

leaved plant.

dictyokinesis (dik'tiökine'sis) n.
[Gk. diktyon, net; kinesis, movement.] The breaking-up of the Golgi-apparatus at mitosis and segregation of dictyosomes to daughter-cells.

dictyonalia (dik'tiŏnā'liā) n. [Gk. diktyon, net.] The principal parenchyma spicules of Dictyonina

and of many Lyssacina.

dictyosome (dik'tiösōm) n. [Gk. diktyon, net; soma, body.] An element of the Golgi-apparatus, q.v.

- dictyospore (dĭk'tiöspōr) n. [Gk. diktyon, net; sporos, seed.] A spore. with transverse and longitudinal septa, of reticular appearance.
- dictyostele (dik'tiöstē'lē] n. [Gk. diktyon, net; stele, post.] A network formed by meristeles.
- dictyotic (dik'tiōt'ik) a. [Gk. diktyon, net.] Appl. moment of shell or skeleton formation, or lorication moment, as of siliceous skeleton of radiolarians; appl. stage in cell growth where chromosomes are lost to view in nuclear reticulum.

- dicyclic (dīsšk'līk) a. [Gk. di-, two; kyklos, circle.] Having a row of perradial infrabasals, appl. theca of Crinoidea; with two whorls; biennial, appl. herbs.
- didactyl (dīdāk'til) a. [Gk. di-, two; daktylos, digit.] Having two fingers or two toes.
- didelphic (diděl'fik) a [Gk. di-, double; delphys, womb.] Having two uteri, as marsupials; amphidelphic.
- didymospore (did'imospor) n. [Gk. didymos, twin; sporos, seed.] A two-celled spore.
- didymous (did'imus) a. [Gk. didymos, twin.] Growing in pairs.
- didynamous (didin'amus) a. [Gk. di-, two; dynamis, power.] With four stamens, two long, two short.

diecious,—dioecious, q.v.

diencephalon. (di'enket'alon, -set'-)
n. [Gk. dia, between; engkephalos, brain.] Part of the fore-brain, comprising thalamencephalon, pars mamillaris hypothalami, and posterior part of third ventricle; between-brain, 'tween-brain, interbrain.

diestrum,- dioestrus, q.v.

- differentiation (dif'ërën'shiā'shūn) n.
 [L. differre, to differ.] Modification in structure and function of the parts of an organism, owing to division of labour.
- diffluence (dif'looens) n. [L. dis, away; fluere, to flow.] Disintegration by vacuolisation.
- diffuse (difūs') a. [L. diffundere, to pour.] Widely spread; not localised; not sharply defined at margin; appl. placenta with villi on all parts except poles.
- diffuse-porous, appl. wood in which vessels of approximately the same diameter tend to be evenly distributed in a growth ring; cf. ring-porous.
- digametic (dīgāmět'īk) a. [Gk. dis, twice; gametes, spouse.] Exhibiting digamety; having two types of gametes, one producing males, the other females; heterogametic.

digastric (dīgăs'trik) a. [Gk. di-,

two; gaster, belly.] Two-bellied, appl. muscles fleshy at ends, tendinous in middle; biventral; appl. one of the suprahyoid muscles; appl. a branch of facial nerve.

digenesis (dījěn'ësis) n. [Gk. dis, twice; genesis, descent.] Alterna-

tion of generations, q.v.

digenetic (dī'jēnēt'ik) a. [Gk. dis, twice; genesis, descent.] Pert. digenesis; requiring an alternation of hosts, appl. certain parasites.

digenoporous (dī'jēnŏp'orūs) a. [Gk. dis, twice; genos, birth; poros, pore.] With two genital pores,

appl. many Turbellaria.

digestion (dijěs'chůn) n. [L. digestio, digestion.] The process by which nutrient materials are rendered absorbable by action of various juices.

digestive (dijěs'tīv) a. [L. digestio, digestion.] Pert. digestion, or having power of aiding in digestion.

- digit (dij'īt) n. [L. digitus, finger.]
 Terminal division of limb in any
 vertebrate above fishes; toe or
 finger; distal part of chelae and
 chelicerae.
- digital (dij'ităl) a. [L. digitus, finger.] Pert. finger or digit; also appl. structures resembling a digit. n. Distal joint of spider's pedipalp.

digitaliform (dij'ītāl'īfôrm) a. [L. digitus, finger; forma, shape.] Finger-shaped, appl. corollae which are like the finger of a glove.

digitate (dij'îtât) a. [L. digitus, finger.] Having parts arranged like the fingers in a hand; with fingers.

digitiform (dij'itifôrm) a. [L. digitus, finger; forma, shape.] Finger-shaped; appl. roots.

digitigrade (dij'itigrād') a. [L. digitus, finger; gradus, step.]
Walking with only the digits touching the ground.

digitinervate (dij'itiner'vat) a. [L. digitus, finger; nervus, sinew.] Having veins radiating out from base like fingers of a hand, with usually five or seven veins; appl. leaves.

digitipartite (dij'îtĭpâr'tīt) a. [L. digitus, finger; partire, to divide.] Having leaves divided up in a hand-like pattern.

digitipinnate (dij'itipin'āt) a. [L. digitus, finger; pinna, feather.] Having digitate leaves of which the

leaflets are pinnate.

digitule (dĭj'ĭtūl) n. [L. digitulus, little finger.] Any small finger-like process; small process on insect tarsi.

digoneutic (dī'gönū'tīk) a. [Gk. dis, twice; goneuein, to produce.]

Breeding twice a year.

- digonic (dīgŏn'īk) a. [Gk. dis, twice; gone, seed.] Producing male and female gametes in separate gones in the same individual; cf. amphigonic.
- digonoporous (di'gönŏp'örŭs) a. [Gk. dis, twice; gone, seed; poros, pore.] With two distinct genital apertures, male and female.
- digynous (dĭj'inŭs) a. [Gk. di-, two; gyne, woman.] Having two carpels.
 diheliotropism, diaheliotropism,

q.v.

diheterozygote (dī'hĕtërözī'gōt) n. [Gk. dts, twice; heteros, other; zygotos, yoked together.] A dihybrid.

dihybrid (dihī'brīd) n. [Gk. dis, twice; L. hibrida, mixed offspring.] A cross whose parents differ in two distinct characters; an organism heterozygous regarding two pairs of alleles

dikaryon (dīkā'rĭŏn) n. [Gk. dis, twice; karyon, nucleus.] A pair of nuclei, as in cells of ascogenous hyphae.

dilatator,-dilator, q.v.

dilated (dilā'tēd) a. [L. dilatare, to flatten.] Expanded, or flattened; appl. parts of insects, etc., with a wide margin.

dilator (dīlā'tŏr) n. [L. dilatare, to expand.] Name appl. any muscle that expands or dilates an organ.

dilophous (dilöf'ūs) a. [Gk. di, two; lophos, crest.] Appl. a tetractinal spicule with two rays forked like a crest. diluvial (dilū'viăl) a. [L. diluvium, deluge.] *Pert*. the present, in geological reckoning.

dimegaly (dīmeg'ālī) n. [Gk. dis, twice; megalos, great.] Condition of having two sizes or a bimodal size frequency; appl. spermatozoa, ova.

dimerous (dim'ërus) a. [Gk. dis, twice; meros, part.] In two parts; having each whorl of two parts; with a two-jointed tarsus.

dimidiate (dimid'iāt) a. [L. dimidius, half.] Having only one-half developed; having capsule split on one side.

dimorphism (dīmôr'fīzm) n. [Gk. dis, twice; morphe, shape.] Condition of having stamens of two different lengths, of having two different kinds of leaves, flowers, etc.; state of having two different forms according to sex, or of one sex, two different kinds of zooids, or of offspring; of broods which, owing to differing conditions, differ in size or colouring.

dimyaric (dimiar'ik) a. [Gk. dis, twice; mys, muscle.] Having two adductor muscles; dimyarian.

dinergate (diner'gāt) n. [Gk. dinein, to roam; ergates, worker.] A soldier ant.

dineuronic (dīnūrŏn'īk) a. [Gk. dis, twice; neuron, nerve.] With double innervation; appl. chromatophores with concentrating and dispersing nerve fibres.

dinomic (dinom'ik) a. [Gk. dis, twice; nomos, district.] Appl. an organism restricted to two of the biogeographical divisions of the globe.

dioecious (diē'siūs) a. [Gk. dis, twice; oikos, house.] Having sexes separate; having male and female flowers on different individuals; gonochoristic; exhibiting dioecism or gonochorism.

dioestrus (dīē'strŭs) n. [Gk. dia, between; oistros, gadfly.] The quiescent period between heat periods in polyoestrous animals; dioestrum.

- dionychous (di'ŏnĭk'ŭs) a. [Gk. di-, two; onyx, nail.] Having two claws, as on tarsi of certain spiders.
- dioptrate (dīŏp'trāt) a. [Gk. dis, twice; ops, eye.] Having eyes or ocelli separated by a narrow line.
- dioptric (diöp'trik) a. [Gk. dioptron, spying-glass.] Pert. transmission and refraction of light; appl. structures, as cornea, lens, aqueous and vitreous humors.
- diorchic (diôr'kik) a. [Gk. dis, twice; orchis, testis.] Having two testes.
- dipetalous (dīpēt'ālūs) a. [Gk. dis, twice; petalon, petal.] Having two petals.
- diphasic (difā'zik) a. [Gk. dis, twice; phainein, to appear.] Appl. extended life cycle of some protozoa, including the active stage; cf. monophasic; periodically changing two states or appearances, as of winter and summer pelage or plumage.
- diphycercal (dif'isĕr'kăl) a. [Ğk. diphyes, twofold; kerkos, tail.] With a tail in which vertebral column runs straight to tip, thereby dividing the fin symmetrically.
- diphygenetic (dĭf'ĭjënĕt'ĭk) a. [Gk. diphyes, twofold; genetes, begotten.] Producing embryos of two different types, as Dicyemida.
- diphygenic (dǐf'ijěn'ik) a. [Gk. diphyes, twofold; genos, descent.] With two types of development.
- diphyletic (dī'fīlĕt'īk) a. [Gk. dis, twice; phylon, race.] Pert. or having origin in two lines of descent.
- diphyodont (dif'iödönt') a. [Gk. diphyes, twofold; odous, tooth.] With deciduous and permanent sets of teeth.
- diplanetary (diplăn'ētărĭ), diplanetic (di'plănět'îk) a. [Gk. dis, twice; planetikos, wandering.]
 With two distinct types of zoospores.
- diplanetism (diplăn'ētīzm) n. [Gk. dis, twice; planetikos, wandering.] Condition of having two periods of motility in one life history, as of zoospores in some fungi.
- diplarthrous (diplar'thrus) a. [Gk. diploos, double; arthron, joint.]
 With tarsal or carpal bones of one

- row articulating with two bones in the other.
- dipleurula (dīploor'ūlă) n. [Gk. dis, twice; pleuron, side.] A bilaterally symmetrical larva of echinoderms; an echinopaedium.
- diplobiont (dĭp'löbī'ŏnt) n. [Gk. diploos, double; bion, living.] An organism characterised by two kinds of individuals, asexual and sexual.
- diploblastic (dĭp'löbläs'tĭk) a. [Gk. diploos, double; blastos, bud.]
 Having two distinct germ layers.
- diplocardiac (dĭp'lökâr'dĭāk) a. [Gk. diploos, double; kardia, heart.]
 With the two sides of the heart quite distinct.
- diplocaulescent (dip'lökôlĕs'ent) a. [Gk. diploos, double; kaulos, stem.] With secondary stems.
- diplochromosome (dǐp'lokrō'mosōm)

 n. [Gk. diploos, double; chroma, colour; soma, body.] Anomalous chromosome having four chromatids, instead of two, attached to centromere.
- diplodal (dĭp'lōdăl) a. [Gk. diploos, double; hodos, way.] Having both prosodus and aphodus; appl. Porifera.
- diploe (dĭp'lōē) n. [Gk. diploe, double.] The cancellous tissue between outer and inner lamellae of certain skull bones; tail of scorpion; mesophyll.
- diplogangliate (dǐp'lögăng'glĭāt) a. [Gk. diploos, double; ganglion, ganglion.] With ganglia in pairs.
- diplogenesis (dip'löjën'ësis) n. [Gk. diploos, double; genesis, descent.]
 Supposed change in germ plasm that accompanies 'use and disuse' changes occurring in body tissues; development of two parts instead of usual single part.
- diploic (diplo'ik) a. [Gk. diploos, double.] Occupying channels in cancellous tissue of bones.
- diploid (dǐp'loid) a. [Gk. diploos, double; eidos, form.] Having a double set of chromosomes; appl. typical or zygotic somatic number of chromosomes of a species. n. A diploid organism; cf. haploid.

- diploidisation,—doubling of number of chromosomes in hapleid cells or hyphae.
- diplonema (dip'lönē'mā) n. [Gk. diploos, double; nema, thread] Double thread of diplotene stage in meiosis.
- diplonephridia (dĭp'lönĕfrĭd'ĭā) n. plu. [Gk. diploos, double; nephros, kidney; idion, dim.] Nephridia derived partly from ectoderm, partly from mesoderm.
- diploneural (dĭp'lönū'răl) a. [Gk. diploos, double; ncuron, nerve.]
 Supplied with two nerves.
- diplont (dĭp'lŏnt) n. [Gk. diploos, double; on, being.] An organism having diploid somatic nuclei, opp. haplont.
- diploperistomous (dip'löperis'tomus)

 a. [(k. diploos, double; peri, around; stoma, mouth.] Having a double projection or peristome.
- diplophase (dip'löfāz) n. [Gk. diploos, double; phasis, aspect.] Stage in life history of an organism when nuclei are diploid; sporophyte phase; diplotene stage in meiosis.
- diplophyll (dĭp'löfil) n. [Gk. diploos, double; phyllon, leaf.] A leaf having palisade tissue on upper and lower side with intermedial spongy parenchyma.
- diploplacula (dip'löpläk'ūlä) n. [Gk. diploos, double; plakoeis, flat cake.] A flattened blastula consisting of two layers of cells.
- diplopore (dīp'löpōr) n. [Gk. diploos, double; poros, passage.] Respiratory organ in Cystoidea.
- diploptile (dip'lotil, -ptīl) a. [Gk. diploos, double; ptilon, feather.]
 Double neossoptile, without rachis, formed by precocious development of the barbs of the teleoptile.
- diplosis (dĭplō'sĭs) n. [Gk. diploos, double.] Doubling of the chromosome number, in syngamy.
- diplosome (dip'lösom) n. [Gk. diploos, double; soma, body.] A double centrosome lying outside the nuclear membrane; a paired heterochromosome.

- diplosomite (diplosomit) n. [Gk. diploso, double; soma, body.] Body segment consisting of two annular parts, prozonite and metazonite, in Diplopoda.
- diplosphene (dĭp'lösfēn) n. [Gk. diploos, double; sphen, wedge.] Wedge-shaped process on neural arch of certain fossil reptiles.
- diplospondylic (dip'löspŏndĭl'īk) a. [Gk. diploos, double; sphondylos, vertebra.] With two centra to each myotome, or with one centrum and well-developed intercentrum; exhibiting diplospondyly.
- diplostemonous (dĭp'löstěm'önŭs) a. [Gk. dɪploos, double; stemon, warp.] With two whorls of stamens in regular alternation with perianth leaves; with stamens double the number of petals.
- diplostichous (dǐplos'tǐkūs) a. [Gk. diploos, double; stichos, row.]
 Arranged in two rows or series.
- diplotegia (dǐp'lötē'jīă) n. [Gk. diploos, double; tegos, roof.] An inferior fruit with dry dehiscent pericarp.
- diplotene (dip'loten) a. [Gk. diploos, double; tainia, band.] Appl. stage in meiosis at which bivalent chromosomes split longitudinally.
- diploxylic (dip'lozil'ik) a. [Gk. diploos, double; xylon, wood.] Appl. leaf-trace bundles with inner and outer strands of wood, in certain extinct plants.
- diplozoic (dĭp'lozō'ĭk) a. [Gk. diploos, double; zoon, animal.] Bilaterally symmetrical.
- dipnoan (dǐp'nōǎn) a. [Gk. dis, twice; pnein, to breathe.] Breathing by gills and lungs.
- diprotodont (diprō'tödont) a. [Gk. dis, twice; protos, first; odous, tooth.] Having two anterior incisors large and prominent, the rest of incisors and canines being smaller or absent.
- n. [Gk. dis, twice; pteron, wing; kekis, gall nut; idion, dim.] Gall caused by a dipterous insect.

dipterous (dip'terus) a. Gk. dis, twice; pteron, wing. With two wings or wing-like expansions.

directive mesenteries,-in Zoantharia, the dorsal and ventral pairs of mesenteries.

disc (disk) n. [L. discus, disc.] Any flattened portion like a disc in shape; middle part of capitulum in Compositae; adhesive tip of tendril; base of sea-weed thallus; circumoral area in many animals; circular areas at opposite poles of many animals; any modification of thalamus; area marking entrance of optic nerve into eye; cup-shaped tactile structures in skin; mass of cells of membrana granulosa which projects into cavity of egg follicle; anisotropic and isotropic parts of contractile fibrils of muscular tissue; disk.

disc-florets,-inner florets borne on abbreviated and reduced peduncle

in many inflorescences.

discal (dĭs'kăl) a. [L. discus, disc.] *Pert.* any disc-like structure; appl. cross-vein between third and fourth longitudinal veins of insect wing. A large cell at base of wing of Lepidoptera completely enclosed by wing-nervures, also in some Dip-

disciflorous (dis'kiflo'rus, dis'iflo'rus) a. [L. discus, disc; flos, flower.] With flowers in which receptacle is

large and disc-like.

disclimax (dis'klī'māks) n. [Gk. dis, double; klimax, ladder.] Disturbance climax, stage in plant succession replacing or modifying true climax, usually due to animal and human agency; e.g. cultivated crops.

discoblastic (dĭs'köblăs'tĭk) a. [Gk. diskos, disc; blastos, bud.] Pert. meroblastic eggs in which area of segmentation is disc-shaped.

discoblastula (dis'köblăs'tūlă) n. [Gk. diskos, disc; blastos, bud.] blastula formed from a meroblastic egg with disc-like blastoderm.

discocarp (dis'kökârp) n. [Gk. diskos, disc; karpos, fruit.] Special enlargement of thalamus below calyx; apothecium.

discoctasters (disk'oktăs'terz) n. plu. [Gk. diskos, disc; okto, eight; aster, star.] Sponge spicules with eight rays terminating in discs, each disc corresponding in position to corners of a cube; modified hexactines.

discodactylous (dĭs'ködăk'tĭlŭs) a. [Gk. diskos, disc; daktylos, finger.]

With sucker at end of digit.

discohexactine (dĭs'köhĕksāk'tĭn) n. [Gk. diskos, disc; hex, six; aktis, ray.] A sponge spicule with six equal rays meeting at angles.

discohexaster (dis'köhčksäs'tër) n. [Gk. diskos, disc; hex, six; aster, star.] A hexactine with rays ending in discs.

discoid (dis'koid) a. [Gk. diskos, disc; eidos, form.] Flat and circular; disc-shaped.

discoidal (diskoi'dal) a. [Gk. diskos, disc; eidos, form.] Disc-like; appl. segmentation in which blastoderm forms a one-layered disc or cap which spreads over yolk; appl. placenta.

(dĭs'kŏntĭnū'ĭtĭ) discontinuity [O.F. discontinuer; from L. dis-, asunder; continuare, to continue.] Occurrence in two or more separate areas or geographical regions; disjunction.

discontinuous variation, - mutation, q.v.

disconula (diskon'ūlă) n. [Gk. diskos, disc.] Eight-rayed stage in larval development of certain Coelentera.

discooctaster, —discoctaster, q.v. discoplacenta (dis'köpläsĕn'tă) [L. discus, disc; placenta, placenta.]

A placenta with villi on a circular

cake-like disc.

discoplasm (dĭs'köplăzm) n. diskos, disc; plasma, form.] Colourless framework or stroma of a red blood corpuscle.

discorhabd (dis'köråbd) n. diskos, disc; rhabdos, rod.] linear sponge spicule with disc-like outgrowths or whorls of spines.

- discus proligerus,-in a Graafian follicle, the mass of cells of membrana granulosa in which the ovum is embedded.
- disjunct (disjungkt') a. [L. disiunctus, separated.] With body regions separated by deep constrictions.
- disjunction (disjungk'shun) n. [L. disiunctus, separated.] Divergence of paired chromosomes at anaphase; geographical distribution in discontinuous areas.
- disjunctive symbiosis,—a mutually condition of symbiosis helpful although there is no direct connection between the partners.
- disjunctor (disjungk'tor) n. disiunctus, separated.] Weak connective structure and zone of separation between successive conidia; 'bridge'.

disk (disk) n. [Gk. diskos, disc.] See disc.

- **disomic** $(d\bar{\imath}'s\bar{o}m\bar{\imath}k)$ a. [Gk. dis, twice: soma, body.] Pert. or having two homologous chromosomes, or genes.
- disoperation (dĭs'ŏpërā'shŭn) n. [L. dis, asunder; opus, work.] Coactions resulting in disadvantage to individual or to group; indirectly harmful influence of organisms upon each other.
- dispermic (dīspēr'mik) a. [Gk. dis, twice; sperma, seed. | Pert., or by, two spermatozoa; appl. fertilisation of an ovum.
- dispermous (dīspēr'mūs) a. ſGk. dis, twice; sperma, seed.] Having two seeds.
- dispermy (dīspēr'mi) n. [Gk. dis, twice; sperma, seed.] The entrance of two spermatozoa into
- dispersal (disper'sal) n. [L. dispergere, to disperse.] The actual scattering or distributing of organisms on earth's surface; transport of diaspores.

disphotic,—dysphotic.

dispireme (dīspī'rēm) n. [Gk. dis, twice; speirema, skein.] The stage of karyokinesis in which each daughter nucleus has given rise to a spireme.

displacement (dĭsplās'mënt) [O.F. desplacier, to displace.] abnormal position of any part of a plant due to its shifting from its normal place of insertion.

disporocystid (dī'spōrösĭs'tĭd) [Gk. dis, twice; sporos, seed; kystis, bladder.] Appl. oocyst of Sporozoa when two sporocysts are present.

disporous (dīspō'rŭs) a. [Gk. dis, twice; sporos, seed.] With two spores.

dissected (dĭsĕk'tëd) a. [L. dissecare] to cut open.] Having lamina cut into lobes, incisions reaching nearly to midrib; with parts displayed.

dissepiment (dis'sep'iment) n. [L. dissaepire, to separate.] The partition found in some compound ovaries; in corals, one of oblique calcareous partitions stretching from septum to septum and closing interseptal loculi below.

dissilient (disil'ient) a. [L. dissilire, to burst asunder.] Springing open; appl. capsules of various plants which dehisce explosively.

dissimilation (dis'simila'shun) n. [L. dissimilis, unlike.] Katabolism,

dissoconch (dis'okongk') n. dissos, double; kongche, shell.] The shell of a veliger larva.

- dissogeny (disŏj'ëni) n. [Gk. dissos, double; genos, descent.] Condition of having two sexually mature periods in the same animal-one in larva, one in adult; also dissogony.
- distad (dĭs'tăd) adv. [L. distare, to stand apart; ad, to.] Towards or at a position away from centre or from point of attachment; in a distal direction.
- distal (dĭs'tăl) a. [L. distare, to stand apart.] Standing far apart, distant, appl. bristles, etc.; pert. end of any structure farthest from middle line of organism or from of attachment. point proximal.

- distalia (distā'liā) n. plu. [L. distare, to stand apart.] The distal or third row of carpal or of tarsal bones.
- distance receptor,—a sense-organ which reacts to stimuli emanating from distant objects; an olfactory, visual, or auditory receptor; disticeptor, distoceptor, teleceptor.

distichalia (distikā'liā) n. plu. [Gk. distichos, with two rows.] In Crinoidea, the secondary brachialia.

- distichous (dis'tikus) a. [Gk. distichos, with two rows.] Two-ranked; appl. alternate leaves, so arranged that first is directly below third.
- distipharynx (dis'tifār'ingks) n. [L. distans, standing apart; Gk. pharyngx, gullet.] A short tube formed by union of epi- and hypopharynx in some insects.

distiproboscis (dĭs'tiprōbŏs'is) n. [L. distans, standing apart; Gk. proboskis, trunk.] Distal portion of insect proboscis, part of ligula.

- dististyle (dis'tistil) n. [L. distans, standing apart; Gk. stylos, pillar.] Distal part or style borne on basistyle, q.v., of gonostyle in mosquitoes.
- distoceptor (dĭs'tósĕp'tŏr) n. [L. distare, to stand apart; recipere, to receive.] A distance receptor, q.v.; teleceptor.
- distractile (distrăk'tīl) a. [L. distractus, pulled asunder.] Widely separate; appl. usually to longstalked anthers.
- distribution (dis'tribū'shūn) n. [L. distributus, divided.] Range of an organism or group in biogeographical divisions of globe.

disymmetrical (dīsĭmĕt'rīkăl) a. [Gk. dis, twice; syn, with; metron, measure.] Biradial, q.v.

- dithecal (dithē'kăl) a. [Gk. dis, twice; theke, box.] Two-celled, as anthers.
- ditokous (dĭt'ökūs) a. [Gk. dis, twice; tokos, birth.] Producing two at a time, either eggs or young.
- ditrematous (dîtrē'mātūs) a. [Gk. dis, twice; trema, opening.] With separate genital openings; with anus and genital openings separate.

- ditrochous (dit'rökŭs) a. [Gk. dis, twice; trochos, runner.] With a divided trochanter.
- diuresis (diūrē'sis) n. [Gk. dia, through; ouron, urine.] Increased or excessive secretion of urine.
- diurnal (diūr'nāl) a. [L. dies, day.]
 Opening during the day only;
 active in the day-time.
- divaricate (dīvār'īkāt) a. [L. divaricatus, stretched apart.] Widely divergent; bifid; forked.
- divaricators (dīvăr'īkātŏrz) n. plu. [L. divaricatus, stretched apart.] Muscles stretching from ventral valve to cardinal process, in brachiopods; muscles in avicularia.
- divergency (diver'jensi) n. [L. divergere, to bend away.] The fraction of a stem circumference, usually constant for a species, which separates two consecutive leaves in a spiral.
- divergent (diver'jent) a. [L. divergere, to bend away.] Separated from one another; appl. leaves.
- diverticulum (di'vertik'ūlūm) n. [I.. de, away; vertere, to turn.] A tube or sac, blind at distal end, branching off from a canal or cavity.
- divided (divi'dëd) a. [L. dividere, to divide.] With lamina cut by incisions reaching midrib; appl. leaves.
- dizoic (dīzō'ĭk) a. [Gk. dis, twice; zoon, animal.] Pert. spore containing two sporozoites.
- dizygotic (dī'zīgŏt'īk) a. [Gk. dis, twice; zygotes, yoked.] Originating from two fertilised ova; appl. twins; dizygous; binovular.
- **Dobie's line,**—Z-disc or telophragma.
- dodecagynous (döděkăj'ínŭs) a. [Gk. dodeka, twelve; gyne, woman.]
 Having twelve pistils.
- dodecamerous (dōděkăm'ërŭs) a. [Gk. dodeka, twelve; meros, part.] Having each whorl composed of twelve parts.
- dodecandrous (dōděkăn'drūs) a. [Gk. dodeka, twelve; aner, man.] Having at least twelve stamens.

dolabriform (dölā'brifôrm) a. [L. dolabra, mattock; forma, shape.]
Axe-shaped.

dolichocephalic (döl'ikökĕfāl'ík,-sĕf-)
a. [Gk. dolichos, long; kephale, head.]
Long-headed; with cephalic index of under 75; cf. brachycephalic.

dolichohieric (dŏl'ĭköhī'erīk) a. [Gk. dolichos, long; hieros, sacred.] With sacral index below 100; cf. platyhieric.

dolichostylous (döl'iköstî'lŭs) a. [Gk. dolichos, long; stylos, pillar.] Pert. long-styled anthers in dimorphic flowers.

dolioform (dō'liöfôrm) a. [L. dolium, wine-cask; forma, shape.] Barrel-

shaped.

- (dom'inant) a. dominant dominans, ruling.] Appl. plants which by their extent determine biotic conditions in a given area; appl. species prevalent in a particular community, or at a given period; appl. character possessed by one parent which in a hybrid masks the corresponding alternative character derived from the other parent: appl. the parental allele manifested in the F_1 heterozygote; recessive; appl. stimulated part of brain when excitation is increased by stimuli usually inducing other reflexes; appl. parts of body controlling less active parts.
- dominator (dom'inātor) n. [L. dominator, ruler.] A broad band of the spectrum which evokes sensation of luminosity in light-adapted eye; cf. modulator.
- dopa (dō'pa) n. [Dihydroxyphenylalanine.] An amino-acid formed from tyrosine by action of ultraviolet rays, and oxidised by dopaoxidase or dopase to a red precursor of melanin, as in basal layers of epidermis; C₉H₁₁O₄N.
- dormancy (dôr'mănsi) n. [F. dormir, from L. dormire, to sleep.] A resting or quiescent condition; reduction in protoplasmic activity due to carbon dioxide concentration, appl. seeds; hibernation and aestivation.

dormancy callus,—callose deposited on sieve areas at the onset of winter.

dorsad (dôr'săd) adv. [L. dorsum, back; ad, to.] Towards back or dorsal surface; opp. ventrad.

dorsal (dôr'săl) a. ÎL. dorsum, back.] Pert. or lying near back, opp. ventral surface; pert. surface farthest from axis; upper surface of thallus or prothallus of ferns, etc.

dorsalis (dôrsā'lĭs) n. [L. dorsum, back.] The artery which supplies

the back of any organ.

dorsiferous (dôrsif'ērŭs) a. [L. dorsum, back; ferre, to carry.] With sori on back of leaf; carrying the young on the back.

dorsifixed (dôr'sifik'st) a. [L. dorsum, back; fingere, to fix.] Having filament attached to back of anther.

dorsigrade (dôr'sigrād) a. [L. dorsum, back; gradus, step.] Having back of digit on the ground when walking.

dorsispinal (dôr'sispi'năl) a. [L. dorsum, back; spina, spine.] Pert. or referring to back and spine.

- dorsiventral (dôr'siven'trăl) a. [L. dorsum, back; venter, belly.] With upper and lower surfaces distinct; bifacial; cf. dorsoventral.
- dorsocentral (dôr'sösĕn'trăl) a. [L. dorsum, back; centrum, centre.] Pert. mid-dorsal surface; pert. aboral surface of echinoderms.

dorsolumbar (dôr'sölüm'băr) a. [L. dorsum, back; lumbus, loin.] Pert. lumbar region of back.

dorsoumbonal (dôr'söŭm'bōnāl) a. [L. dorsum, back; umbo, shieldboss.] Lying on the back near the umbo.

- dorsoventral (dôr'sövěn'trăl) a. [L. dorsum, back; venter, belly.] Pert. structures which stretch from dorsal to ventral surface; cf. dorsiventral.
- dorsulum (dôr'sūlŭm) n. [Dim. of L. dorsum, back.] Upper surface lying between collar and scutellum; mesonotum.
- dorsum (dôr'sūm) n. [L. dorsum, back.] The sulcular surface of Anthozoa; tergum or notum of insects and crustaceans; inner

- margin of insect wing; the back of higher animals.
- dorylaner (dŏr'ilāner) n. [Gk. dory, spear; aner, male.] An exceptionally large male ant of driver-ant group.
- doublure (dooblūr') n. [F. doublure, lining.] The reflected margin of carapace in Trilobita and Xiphosura.
- **Doyère's cone** ([L. Doyère, French physiologist]. End-plate or eminence where nerve fibre branches and enters sarcolemma.
- drepanium (drepā'nĭŭm) n. [Gk. drepane, sickle.] A helicoid cyme with secondary axes developed in a plane parallel to that of main peduncle and its first branch.
- drift (drift) n. [A.S. drifan, to drive.]
 Transported, opp. bed-rock, soils;
 process of change in gene
 frequencies in a population of
 breeding individuals; Sewall
 Wright effect; genetico-automatic
 process.
- dromaeognathous (drom'eog'năthus) a. [Gk. dromein, to run;
 gnathos, jaw.] Having a palate
 in which palatines and pterygoids
 do not articulate, owing to intervention of vomer.
- dromotropic (drŏm'ötrŏp'îk) a. [Gk. dromos, course; trope, turn.] Bent in a spiral; influencing nerve conductivity.

drone (dron) n. [A.S. dran.] The male bee.

dropper (drŏp'ër) n. [A.S. dreópan, to drop.] Rhizomatous downward outgrowth of a bulb, which may form a new bulb.

drupaceous (droopā'shŭs) a. [Gk. dryppa, olive.] Pert. drupe; bearing drupes; drupe-like.

drupe (droop) n. [Gk. dryppa, olive.] A superior, one-celled fruit with one or two seeds and the pericarp differentiated into a thin epicarp, a fleshy sarcocarp, and a hard endocarp, as of plum.

drupel (droop'ël) n. [Gk. dryppa, olive.] An individual component of aggregate fruit, as of raspberry;

also drupelet.

- dry (dri) a. [A.S. dryge, dry.] Appl. achenial, capsular, and schizocarpic fruits. Opp. succulent.
- drymophytes (drī'mofīts) n. plu. [Gk. drymos, coppice; phyton, plant.] Small trees, bushes, and shrubs.
- duct (dūkt) n. [L. ducere, to lead.]
 Any tube which conveys fluid or other substance; a tube formed by a series of cells which have lost their walls at the points of contact; ductus.
- ductless glands,—glands which do not communicate with any organ directly by means of a duct; endocrine organs.
- ductule (duk'tūl) n. [L. ducere, to lead.] A minute duct; fine thread-like terminal portion of a duct.
- ductus (dŭk'tŭs) n. [L. ducere, to lead.] Duct, q.v.

ductus deferens,—vas deferens.

- ductus ejaculatorius,—a narrow muscular tube at end of vas deferens in various invertebrates.
- Dufour's gland [L. Dufour, French entomologist]. An alkaline gland with duct leading to terebra or sting of certain Hymenoptera.
- dulosis (dū'lōsis) n. [Gk. doulosis, subjugation.] Slavery, among ants. dumb-bell bone,—prevomer, q.v.
- duodenal (dū'ödē'năl) a. [L. duodeni, twelve each.] Pert. duodenum.
- duodenum (dū'ödē'nŭm) n. [L. duodeni, twelve each.] That portion of small intestine next to pyloric end of stomach.
- duplex (dū'plēks) a. [L. duplex, twofold.] Double; compound, appl. flowers; diploid; having two dominant genes, in polyploidy; consisting of two distinct structures; having two distinct parts.
- duplication (dū'plikā'shūn) n. [L. duplex, double.] Chorisis, q.v.; a translocated chromosome fragment attached to one of normal set.
- duplicature (dū'plikătūr) n. [L duplex, double.] A circular fold near base of protrusible portion of a polyzoan polypide.

duplicident (dūplis'iděnt) a. duplex, double; dens, tooth.] With two pairs of incisors in upper jaw,

one behind the other.

duplicity (dūplis'īti) n. [L. dupliciter, doubly.] Condition of being twofold; appl. theory that cones are the photopic, or colour, receptors, and rods the scotopic, or brightness, receptors.

duplicodentate (dū'pliköděn'tāt) a. [L. duplex, double; dens, tooth.] With marginal teeth on leaf bearing smaller teeth-like structures.

dura mater (dū'ră mā'tër) n. dura, hard; mater, mother.] The tough membrane lining the whole cerebro-spinal cavity.

dura spinalis,—the tough membrane lining the spinal canal.

dural (dū'răl) a. [L. dura, hard.] *Pert.* dura mater; appl. sheath of optic nerve.

duramen (dūrā'mën) n. [L. durare, to harden.] The hard, darker central region of a tree-stem; the heartwood.

duvet (dūvā') n. [F. duvet, down.] Downy coating, as soft matted coat-

ing by certain fungi.

dwarf male,-small three- or fourcelled plant formed from androspore of Oedogonium; a small, usually simply formed, individual in many classes of animals, either free or carried by the female.

dyad (dī'ad) n. [Gk. dyas, two.] The half of a tetrad group; a bivalent

chromosome.

dynamic (dinăm'ik) a. [Gk. dynamis, power.] Producing or manifesting activity, opp. static; appl. specific dynamic action, the calorigenic action of food, increasing metabolism above basal rate.

dynamoplastic (di'nămoplas'tik) a. [Gk. dynamis, power; plastos, formed.] Appl. active type of energid-product; opp. paraplastic.

dysgenic (disjěn'ik) a. [Gk. dysgeneia, low birth.] Pert. tending towards, or productive of, racial degeneration; kakogenic, •ugenic.

dysmerism (dis'mërizm) n. [Gk. dys-, mis-; meros, part.] An aggregate of unlike parts.

dysmerogenesis (dis'merojen'esis) n. [Gk. dys-, mis-; meros, part; genesis, descent.] Segmentation

resulting in unlike parts.

dysphotic (dĭs'fōtĭk) a. [Gk. dys-, mis-; phos, light.] Dim; appl. zone, waters at depths between 80 and 600 metres, between euphotic and aphotic zones, q.v.; layer of photic zone.

dysploid,—aneuploid, q.v.

dyspnoea (dispnē'a) n. [Gk. dyspnoos, breathless.] Difficulty in breathing.

- dysteleology (dĭs'tělëŏl'öjĭ) n. [Gk. dys-, mis-; teleos, ended; logos, discourse.] Haeckel's doctrine of purposelessness in Nature; appearance of uselessness, as of certain organs or other structures; frustration of function.
- dystrophic (distrof'ik) a. [Gk. dys-, mis-; trephein, nourish.] to Wrongly or inadequately nourished; inhibiting adequate nutrition; pert. faulty nutrition.
- theory [J. Dzierzon, Dzierzon Silesian apiculturist]. Belief that males of honey-bee are always produced from unfertilised eggs.

ear (ēr) n. [A.S. éare.] The auditory organ; among invertebrates, the various structures supposed to have an auditory function; specialised tufts of hair or feathers which are close to, or similar to an external ear or pinna; an earshaped structure; the spike of grasses, usually of cereals.

eared (ērd) n. [A.S. éare.] Having external ears or pinnae; with tufts of feathers resembling ears; having long bristles or processes, as in

grains of corn; auriculate.

ebracteate (ēbrăk'tëāt), ebracteolate (ēbrāk'tëölāt) a. [L. ex, out of; bractea, thin plate.] Without bracts; without bracteoles.

ecad,—oecad, q.v.

- ecalcarate (ēkàl'kārāt) a. [L. ex, out of; calcar, spur.] Having no spur or spur-like process.
- ecardinal (ēkâr'dĭnăl) a. [L. ex, out of; cardo, hinge.] Having no hinge; also ecardinate.
- ecarinate (ēkăr'ināt) a. [L. ex, out of; carina, keel.] Not furnished with keel or keel-like ridge.
- ecaudate (ēkô'dāt) a. [L. ex, out of; cauda, tail.] Without a tail.
- ecblastesis (ĕk'blāstē'sis) n. [Gk. ek, out of; blastos, bud.] Proliferation of main axis of inflorescence.
- eccritic (ekrit'ik) a. [Gk. ekkrinein, to expel, to select.] Causing or pert. excretion; preferred, appl. temperature, etc.
- ecdemic (ěkděm'ĭk) a. [Gk. ek, out of; demos, people.] Not native.
- ecderon (ĕk'derŏn) n. [Gk. ek, out; deros, skin.] The outer or epidermal layer of skin.
- ecderonic (ěkdërŏn'ĭk) a. [Gk. ek, out; deros, skin.] Ectodermic; epiblastic.
- ecdysial (ĕkdīs'īāl) a. [Gk. ekdysai, to strip.] Pert. ecdysis; appl. fluid between old and new cuticle which aids in disintegration of old cuticle, moulting fluid; appl. line along which cuticle splits in moulting; appl. glands, Verson's glands, secreting moulting fluid.
- ecdysis (ĕk'dĬsĭs) n. [Gk. ekdysai, to strip.] The act of moulting a cuticular layer or structure; cf. endysis.
- ece, oike. o.v.
- ecesis,—oikesis, q.v.
- echard (ékârd') n. [Gk. echein, to keep; ardo, I water.] Soil water not available for plant growth; cf. chresard, holard.
- echinate (ěk'ĭnāt) a. [Gk. echinos, hedgehog.] Furnished with spines or bristles.
- echinenone (ĕkǐn'ēnōn) n. [Gk. echinos, sea-urchin.] A carotenoid pigment of sea-urchin gonads, a provitamin A.
- echinochrome (ëkī'nökrōm) n. [Gk. echinos, sea urchin; chroma,

- colour.] A red-brown respiratory pigment of echinoderms; $C_{12}H_{11}O_2$.
- echinococcus (ëkī'nökŏk'ŭs) n. [Gk. echinos, spine; kokkos, berry.] A vesicular metacestode developing a number of daughter cysts, each with many heads.
- echinoid (ěk'ínoid) a. [Gk. echinos, sea-urchin; eidos, form.] Pert. or like sea-urchins.
- echinopaedium (ėkī'nopē'dīŭm) n. [Gk. echinos, sea-urchin; paidion, young child.] Dipleurula, q.v.
- echinopluteus (eki'nöploo'teŭs) n.
 [L. echinus, sea-urchin; pluteus, shed.] Larva or pluteus of echinoids, from supposed resemblance to an upturned easel.
- echinulate (ėkin'ūlāt) a. [Gk. echinos, spine.] Having small spines; having pointed outgrowths, appl. bacterial cultures.
- echolocation (ĕk'ōlōkā'shǔn) n. [L. echo, echo; locare, to place.] Location of objects by means of echos, as of supersonic sounds emitted by animals, e.g. by bats.
- eclipse (ĕklĭps') n. [Gk. ekleipein, to leave incomplete.] Plumage assumed after spring moult, as in drake.
- eclosion (ëklō'zhūn) n. [L. e, out; clausus, shut.] Hatching from an egg or imago.
- eco-, -see also occo-, oiko-.
- ecobiotic (ē'kobīŏt'ĭk) a. [Gk. oikos, household; biosis, manner of life.] Appl. adaptation to particular mode of life within a habitat.
- ecodeme (ē'ködēm) n. [Gk. oikos, household; demos, people.] A deme occupying a particular ecological habitat.
- **ecoid**,—oecoid, q.v.
- ecology (ēkŏl'öji) n. [Gk. oikos, household; logos, discourse.] That part of biology which deals with relationship between organisms and their surroundings; bionomics.
- economic density,—of a population, the number of individuals per unit of the inhabited area, opp. population density in an area only partly inhabited.

- ecostate (ēkŏs'tāt) a. [L. e, out; costa, rib.] Without costae; not costate.
- ecosystem (ē'kösīs'tem) n. [Gk. oikos, household; systema, composite whole.] Ecological system formed by the interaction of coacting organisms and their environment.
- ecotone (ē'kötōn) n. [Gk. oikos, household; tonos, brace.] A transitional species in intermediate area between two associations; the boundary line or transitional area between two communities.
- ecotope (ē'kötōp) n. [Gk. oikos, household; topos, place.] A particular kind of habitat within a region.
- ecotype (ē'kötīp) n. [Gk. oikos, household; typos, pattern] A biotype resulting from selection in a particular habitat; habitat type.

ecphoria (ěkfő'rĭă) n. [Gk. ekphorion, produce.] The revival of a latent memory pattern or engram.

- ecsoma (ĕksō'mă) n. [Gk. ek, from out of; soma, body.] Retractile posterior part of body in certain trematodes.
- ectad (ĕk'tăd) adv. [Gk. ektos, outside; L. ad, towards.] Towards the exterior; outwards externally. Opp. entad.
- ectadenia (ĕk'tădē'nĭā) n. plu. [Gk. ektos, outside; aden, gland.] Ectodermal accessory genital glands in insects; cf. mesadenia.
- ectal (ĕk'tăl) a. [Gk. ektos, outside.]
 Outer; external; appl. layer or
 membrane on margin of exciple.
 Opp. ental.
- ectamnion (ěktăm'niŏn) n. [Gk. ektos, outside; amnion, foetal membrane]. Ectodermal thickening in proamnion, beginning of head-fold.
- ectendotrophic (ĕk'těndötrŏf'īk) a. [Gk. ectos, without; endon, within; trophe, nourishment.] Partly ectotrophic and partly endotrophic, appl. mycorhizic fungus.
- ectental line,—the line where ecto-

derm and endoderm meet at blastopore of a gastrula.

- ectepicondylar (ĕkt'ĕpikŏn'dilär) a. [Gk. ektos, outside; epi, upon; kondylos, knob.] Appl. radial foramen of humerus.
- ectethmoid (ěktěth'moid) n. [Gk. ektos, outside; ethmos, sieve; eidos, form.] Lateral ethmoid bone.
- ecthoraeum (ěk'thŏrē'ŭm) n. [Gk. ekthroskein, to leap out.] The thread of a nematocyst.
- ectoblast (ĕk'toblăst) n. [Gk. ektos, outside; blastos, bud.] Epiblast, q.v.
- ectobronchus (ĕk'töbrŏng'kŭs) n.
 [Gk. ektos, outside; brongchos,
 windpipe.] Lateral branch of
 main bronchus in birds; also
 ectobronchium.
- ectocarpous (ĕk'tökâr'pŭs) a. [Gk. ektos, outside; karpos, fruit.] Having gonads of ectodermal origin.
- ectochondrostosis (ĕk'tökôndrōstō'sīs) n. [Gk. ektos, outside; chondros, cartilage; osteon, bone.] Deposition of lime-salts beginning in perichondrium and gradually invading cartilage.
- ectochone (ĕk'tökōnē) n. [Gk. ektos, outside; choane, funnel.] A funnel-shaped chamber into which lead the ostia in certain sponges.
- ectocoelic (ĕk'tösē'lĭk)a. [Gk. ektos, outside; koilos, hollow.] Pert. structures situated outside the enteron of coelenterates.
- ectocondyle (ĕk'tökŏn'dĬl) n. [Gk. ektos, outside; kondylos, knob.]
 The outer condyle of a bone.
- ectocranial (ĕk'tökrā'niāl) a. [Gk. ektos, outside; kranion, skull.]

 Pert. outside of skull.
- ectocuneiform (ĕk'tökūnē'ifôrm) n.
 [Gk. ektos, outside; L. cuneus, wedge; forma, shape.] A bone in distal row of tarsus; third cuneiform.
- ectocyst (ek'tosist) n. [Gk. ektos, outside; kystis, bladder.] Outer layer of zooecium in Polyzoa; outer covering of encysted Protozoa; epicyst.
- ectoderm (čk'töděrm) n. [Gk. ektos, outside; derma, skin.] The outer

layer of a multicellular animal; the epidermis in higher mammals.

ecto-ethmoid, -ectethmoid, q.v.

- ectogenesis (ĕk'töjĕn'esīs) n. [Gk. ektos, outside; genesis, descent.] Embryonic development outside the maternal organism; development in an artificial environment.
- ectogenous (ĕktŏj'ēnŭs) a. [Gk. ektos, outside; genos, birth.] Able to live an independent life; originating outside the organism.
- ectoglia (ĕktoglī'ā) n. [Gk. ektos, outside; glia, glue.] An outer layer in central nervous system.
- ectolecithal (ĕk'tölĕs'ĭthăl) a. [Gk. ektos, outside; lekithos, yolk of egg.] Having yolk surrounding formative protoplasm.
- ectoloph (ĕk'tölŏf) n. [Gk. ektos. outside; lophos, crest.] The ridge stretching from paracone to metacone in a lophodont molar.
- ectomere (ĕk'tömēr) n. [Gk. ektos, outside; meros, part.] An epiblast cell which gives rise to ectoderm.
- -ectomy (ĕk'tömĭ). [Gk. ek, out; temnein, to cut.] Suffix signifying an excision, e.g. thyroidectomy, gonadectomy, etc.
- ectoneural (ĕk'tönū'ral) a. [Gk. ektos, outside; neuron, nerve.] Appl. system of oral ring, radial, and subepidermal nerves in echinoderms.
- ectoparasite (ĕk'topăr'ăsīt) n. [Gk. ektos, outside; para, beside; sitos, food.] A parasite that lives on the exterior of an organism.
- ectopatagium (ĕk'töpătājī'ŭm) n. [Gk. ektos, outside; L. patagium, border.] The part of the wing-like membrane of bats which is carried on metacarpals and phalanges.
- ectophyte (ĕk'töfit) n. [Gk. ektos, outside; phyton, plant.] An external plant parasite.
- ectopic (ěktőp'ík) a. [Gk. ek, out of; topos, place.] Not in normal position; appl. organs, gestation, etc. Opp. entopic.
- ectopic pairing,—pairing between bands located in different regions of a chromosome.

- ectoplasm (ek'toplazm) n. [Gk. ektos, outside; plasma, mould.] The external layer of protoplasm in a cell, usually modified; ectosarc of protozoan cell; layer next cellwall.
- ectoplast (ěk'töplāst) n. [Gk. ektos, outside; plastos, formed.] The protoplasmic film or plasma-membrane just within the true wall of a cell.
- ectopterygoid (ĕk'tötĕr'igoid,-ptĕr-)

 n. [Gk. ektos, outside; pteryx, wing; eidos, form.] A ventral membrane bone behind palatine and extending to quadrate; mesopterygoid; os transversum between pterygoid and maxilla in many reptiles and in some fishes; cf. entopterygoid.
- ectosarc (čk'tösârk) n. [Gk. ektos, outside; sarx, flesh.] The external layer of protoplasm in a protozoon.
- ectosome (ĕk'tosōm) n. [Gk. ektos, outside; soma, body.] The enveloping portion of a sponge containing no flagellated chambers; a type of cell granule.
- ectosphere (ěk'tösfēr) n. [Gk. ektos, outside; sphaira, globe.] The outer zone of attraction sphere.
- ectospore (ĕk'töspōr) n. [Gk. ektos, outside; sporos, seed.] The spore formed at end of each sterigma in Basidiomycetes.
- ectostosis (čk'tŏstō'sis) n. [Gk. ektos, outside; osteon, bone.] Formation of bone in which ossification begins under the perichondrium and either surrounds or replaces the cartilage.
- ectotheca (ĕk'töthē'kā) n. [Gk. ektos, outside; theke, cup.] Outer coating of gonotheca in certain hydroids.
- ectotrachea (ĕk'tötrākē'ā) n. [Gk. ektos, outside; L. trachea, windpipe.] An epithelial layer on outer side of insect tracheae.
- ectotrophic (ĕk'tötröf'îk) a. [Gk. ektos, outside; trephein, to nourish.] Finding nourishment from outside; appl. fungi which surround roots of host with hyphae. Opp. endotrophic.

- ectoturbinal (ěk'tötűr'binál) n. [Gk. ektos, outside; L. turbo, whirl.] A division of the ethmoturbinal.
- ectozoon (ĕk'tözō'ŏn) n. [Gk. ektos, outside; zoon, animal.] An external animal parasite; epizoon.
- ecumene (ěk'ūměn'ē) n. [Gk. oikoumene, habitable world.] Any inhabited region; the biosphere.
- edaphic (ědăf'îk) a. [Gk. edaphos, ground.] Pert. or influenced by conditions of soil or substratum.
- edaphology (ĕdāfŏl'öjī) n. [Gk. edaphos, ground; logos, discourse.] Soil science; particularly the study of the influence of soil on living organisms; cf. pedology.
- edaphon (ed'afon) n. [Gk. edaphos, ground.] The organisms living within the soil; soil flora and fauna.
- edentate (ēděn'tāt) a. [L. ex, without; dens, tooth.] Without teeth or tooth-like projections.
- edestin (čděs'tǐn) n. [Gk. edestos, eatable.] A plant globulin, main protein of sunflower and certain other seeds.
- edriophthalmic (ĕd'rĭŏfthăl'mĭk) a. [Gk. edra, seat; ophthalmos, eye.] Having sessile eyes; appl. certain Crustacea.
- effector (éfék'tőr) n. [L. efficere, to carry out.] An organ which reacts to stimulus by producing work or substance, as muscle, electric and luminous organs, glands; a motor end-organ in muscle.
- efferent (čf'ërënt) a. [L. ex, out; ferre, to carry.] Conveying from, appl. vessels, lymphatics, etc.; carrying outwards, appl. impulses carried outwards by motor nerves.
- efflorescence (éflörés'ens) n. [L. efflorescere, to blossom.] Blossoming; time of flowering; bloom.
- effoliation (ĕfföliā'shūn) n. [L. ex, out of; folium, leaf.] Shedding or removal of leaves.
- effuse (ĕfūs') a. [L. effusus, poured out.] Spreading loosely, appl. inflorescence; spreading thinly, appl. bacterial cultures.

- egest (ējĕst') v. [L. ex, out; gerere.
 to carry.] To throw out; to void;
 to excrete.
- egesta (ējēs'tā) n. plu. [L. egestus, discharged.] The sum-total of substances and fluids discharged from body.
- egg (ĕg) n. [A.S. æg, Icel. egg.] The matured female germ-cell; ovule.
- egg-albumin,—the chief constituent of white of egg, a mixture of glucoproteins.
- egg-apparatus,—the two synergids and ovum proper, near micropyle in embryo-sac of seed plants.
- egg-case,—a protective covering for eggs.
- egg-cell,—the ovum proper apart from any layer of cells derived from it or from other cells.
- egg-membrane,—the layer of tough tissue lining an egg shell.
- egg-nucleus, the female pro-
- egg-tooth,—a small structure on tip of upper jaw, or of beak, by which the embryo breaks its shell.
- eiloid (i'loid) a. [Gk. eilein, to roll up; eidos, form.] Shaped like a coil.
- ejaculate (ëjăk'ūlāt) n. [L. ejaculatus, thrown out.] The emitted seminal fluid.
- ejaculatory (ëjăk'ūlătörĭ) a. [L. ejaculare, to throw out.] Throwing out; appl. certain ducts.
- ejaculatory sac,—organ pumping ejaculate from vas deferens through ejaculatory duct to penis, in certain insects.
- elaborate (ëlăb'örāt) v. [I.. elaborare, to work out.] To change from a crude state to a state capable of assimilation; to form complex organic substances from simple materials.
- elaeoblast (ĕlē'öblāst) n. [Gk. elaion, oil; blastos, bud.] A mass of nutrient material at posterior end of body in certain tunicates.
- elaeocyte (ĕlē'ösīt) n. [Gk. elaion, oil; kytos, hollow.] A cell containing fatty droplets, found in coelomic fluid of annelids.

- elaeodochon (ĕl'ĕŏd'okŏn) n. [Gk. elaiodochos, oil-containing] The preen-gland or oil-gland in birds.
- elaioplast (ĕlī'öplāst') n. [Gk. elaion, oil; plastos, moulded.] A plastid in a plant cell which forms or helps to form oil globules.

elaiosome,—elaioplast.

- elaiosphere (ĕlī'ösfēr) n. [Gk. elaion, oil; sphaira, globe.] An oil globule in a plant cell.
- elastic fibro-cartilage,—consists of cartilage cells and a matrix pervaded by a network of yellow elastic fibres which branch and anastomose in all directions.
- elastica externa,—external layer of notochordal sheath.
- elastica interna, the epitheliomorph layer of notochordal cells.
- elastin (ëlăs'tin) n. [Gk. elaunein, to draw.] The sclero-protein of which elastic fibres are composed.
- elater (ĕl'ātēr) n. [Gk. elater, driver.] One of cells with a spiral thickening which assist in dispersing spores from capsule in liverworts; one of spore appendages formed from epispore in horsetails.
- elaterophore (ĕl'ătĕröfōr) n. [Gk. elater, driver; pherein, to bear.]
 Tissue bearing the elaters, in some liverworts.
- electosome (ëlëk'tösōm) n. [Gk. eklektikos, chosen; soma, body.]
 A chondriosome regarded as a centre for elaborating and fixing chemical constituents of protoplasm.
- electric organ,—modifications of muscles or groups of muscles which discharge electric energy, found in certain fishes.
- electroendosmotic layer,—a hypothetical 'membrane' present between two neurons or between neuron and muscle cell.
- electrolemma (ëlĕk'trölĕm'ă) n. [Gk. elektron, amber; lemma, skin.] Membrane surrounding an electroplax.
- electrophoresis (ělěk'tröförē'sis) n. [Gk. elektron, amber; pherein, to

- bear.] Transport of substances, as of colloidal particles, resulting from differences in electrical potential.
- electropism,—electrotropism, q.v. electroplax (ëlěk'tröplāks) n. [Gk. elektron, amber; plax, plate.] One of the constituent plates of an
- electric organ.
 electrotaxis (ëlěk'trötāk'sīs) n. [Gk. elektron, amber; taxis, arrangement.] Orientation of movement within an electric field.
- electrotonic (ëlëk'trötŏn'ik) a. [Gk. elektron, amber; tonos, tension.]

 Pert. a state of electric tension.
- electrotonus (élěktrőt'önűs, ëlěktrötön'űs) n. [Gk. elektron, amber; tonos, tension.] The modified condition of a nerve when subjected to a constant current of electricity.
- electrotropism (ělěktrŏt'röpĭzm) n. [Gk. elektron, amber; trope, turn.] Reaction of an organism to electric stimuli; plant curvature in an electric field.
- eleidin (ĕlē'ídin) n. [Gk. elaia, olive.] Substance found as small granules or droplets in stratum granulosum of epidermis.

eleo,—see elaco-, elaio-.

- elewtherodactyl (ělū'thërödāk'tĭl) a. [Gk. eleutheros, free; daktylos, finger.] Having hind toe free.
- eleutheropetalous (člū'thëröpět'ălŭs) a. [Gk. eleutheros, free;
 petalon, petal.] Having petals or
 components of whorl free or
 separate.
- eleutherophyllous (ĕlū'thĕröfil'ūs)

 a. [Gk. eleutheros, free; phyllon, leaf.] Having components of perianth whorls free.
- eleutherosepalous (člū'thërösĕp'ălūs) a. [Gk. eleutheros, free;
 F. sépale, sepal.] Having sepals free
 or separate.
- elevator (ĕl'ĕvātör) n. [L. elevare, to lift up.] Any muscle which raises a part.
- eligulate (ēlig'ūlāt) a. [L. ex, out; ligula, little tongue.] Having no ligule; appl. certain club-mosses.

elimination bodies,-nucleic acid material expelled from each chromosome during meiosis, remaining in middle of spindle and disintegrating

during telophase.

elliptical (ělĭp'tĭkăl) a. [Gk. elleipsis, falling short.] Oval-shaped; appl. leaves of about same breadth at equal distances from base and apex, which are slightly acute.

eluvial (ēlū'vĭăl) a. [L. ex, out; luere, to wash.] Appl. leached upper layers or A horizon of soil.

elytriform (ělĭt'rĭfôrm) a. [Gk. elytron, sheath; L. forma, shape.] Shaped like an elytrum.

elytroid (ĕl'ĭtroid) a. [Gk. elytron, sheath; eidos, resemblance.] Re-

sembling an elytrum.

- elytrum (ĕl'ĭtrŭm) n. [Gk. elytron, The anterior wing of sheath.] certain insects, hard and case-like; one of scales or shield-like plates found on dorsal surface of some worms; also elytron.
- emarginate (ēmâr'jĭnāt) a. [L. ex, out; marginare, to delimit.] Having a notch at apex; having a notched margin.

embole (ěm'bölē) n. [Gk. embole, a throwing in.] Invagination; also emboly.

embolic (ĕmbŏl'ĭk) a. [Gk. embole, a throwing in.] Pushing or growing in.

- embolium (ĕmbō'lĭŭm) n. embolos, wedge.] Outer or costal part of wing, or basal part of hemelytron, in certain insects.
- embolomerous (ĕm'bölŏm'ërŭs) a. [Gk. embolos, wedge; meros, part.] Having two vertebral rings in each segment, due to union of hypocentra with neural arch, and union of two pleurocentra below notochord.

embolus (ĕm'bölŭs) n. [Gk. embolos, wedge.] Apical division of the palpus in certain spiders; a clot blocking a blood-vessel.

embryo (ĕm'briö) n. [Gk. embryon, embryo.] A young organism in early stages of development before it becomes self-supporting.

embryo cell,—one of two cells formed from first division of fertil-

in certain plants, egg developing later into embryo, the other developing into suspensor.

embryogenesis (ěm'briöjěn'ësis) n. [Gk. embryon, embryo; genesis, Origin of the embryo; descent.] embryogeny, q.v.

- embryogeny (ěmbrioj'ëni) n. [Gk. embryon, embryo; gennaein, to produce.] The processes by which the embryo is formed; origin, cellular pattern, and functions of the embryo.
- embryology (ěmbričl'öji) n. [Gk. embryon, embryo; logos, discourse.] That part of biology dealing with formation and development of the embryo.
- embryonal knot,-inner cell mass of blastodermic vesicle.
- embryonic (ĕmbrĭŏn'ĭk) a. [Gk. embryon, embryo.] Pert. embryo.
- embryonomy (ěmbriŏn'ömi) n. [Gk. embryon, embryo; nomos, law.] The laws of embryonic development; classification of embryos. a. Embryonomic.
- embryophore (ěm'briöfōr) n. [Gk. embryon, embryo; pherein, bear.] Ciliated mantle enclosing embryo in many tape-worms, and formed from superficial blastomeres of embryo.
- embryophyta (ĕm'brĭöfī'tă) n. plu. [Gk. embryo, embryo; phyton, plant.] Plants having an enclosed embryo, as those with an archegonium, or bearing seeds.

embryo-sac,—the megaspore; female gametophyte in angiosperms.

- embryotectonics (ĕm'brĭötĕktŏn'ĭks) n. [Gk. embryon, embryo; tekton, builder.] The structure or cellular pattern of the embryo.
- embryotega (ěm'briŏt'ěgă) n. [Gk. embryon, embryo; tegos, roof.] Small hardened portion of testa which marks micropyle in some seeds and separates like a little lid at period of germination.
- embryotrophy (ěm'briŏt'röfi) n. [Gk. embryon, embryo; trophe, nourishment.] Nourishment of embryo. or means adapted therefor.

emergence (ēmēr'jens) n. [L. ex, out; mergere, to dip.] An outgrowth from subepidermal tissue;

an epidermal appendage.

emersed (ēměrs't) a. [L. ex, out; mergere, to dip.] Rising above surface of water; appl. leaves.

- eminence (ĕm'ĭnëns) n. [L. eminens. eminent.] Ridge or projection on surface of bones; eminentia.
- emissary (ěm'īsărĭ) a. [L. emittere, to send out.] Coming out; name appl. veins passing through apertures in cranial wall and establishing connection between sinuses inside and veins outside.
- emmenine (ĕm'ēnĭn) n. [Gk. emmenos, monthly.] placental gonadotrophic hormone.
- empodium (ěmpo'dĭŭm) n. [Gk. en, in; pous, foot.] A small variable median structure between claws of feet in many insects.
- emulsin (ēmŭl'sĭn) n. [L. emulgere, to milk out.] A hydrolytic enzyme found in certain plants and some invertebrates.
- enamel (ĕnăm'ël) n. [O.F. esmaillier, to coat with enamel.] The hard material containing over 90 per cent. calcium and magnesium salts which forms a cap over dentine, or may form a complete coat to tooth or scale.
- enamel cells,—cells which form enamel, collectively the enamel organ; adamantoblasts, ameloblasts.
- enantiobiosis (ĕnăn'tĭöbiō'sis) [Gk. enantios, opposite; bios, life.] Antagonistic symbiosis.
- enantioblastic (ěnăn'tiöblăs'tik) a. [Gk. enantios, opposite; blastos, Formed at end of seed bud.] opposite placenta.
- enantiomorphic (ěnăn'tĭömôr'fĭk) a. [Gk. enantios, opposite; morphe, form.] Similar but contraposed, as mirror image, right and left hand; deviating from normal symmetry.
- enarthrosis (ĕn'arthrō'sĭs) n. [Gk. en, in; arthron, joint.] Ball-andsocket joint.
- enation (ena'shun) n. [L. enatus, grown from.] A non-reproductive

accessory part emerging from surface of telome; outgrowth from a previously smooth surface.

END-

encephalisation (ĕnkēf'ālīzā'shun, -sĕf'-) n. [Gk. engkephalos, brain.] Brain formation by the forwardshifting and centralising tendency of co-ordinating neurones.

encephalocoel (ĕnkĕf'ălösēl, -sĕf-) n. [Gk. engkephalos, brain; koilos, hollow.] Cavity within the brain; cerebral ventricle, the anterior dilatation of neurocoel.

- encephalomere (ěnkěf'ălömēr, -sěf-) n. [Gk. engkephalos, brain; meros, part.] A brain segment.
- encephalon (ĕnkĕf'ālŏn, -sĕf-) n. [Gk. engkephalos, brain.] The brain.
- encephalospinal (ĕnkĕf'ălöspī'năl, -sĕf-) a. [Gk. engkephalos, brain; L. spina, spine.] Pert. brain and spinal cord.
- enchondral,—endochondral, intracartilaginous, q.v.
- enchylema (ĕnkīlē'mă) n. [Gk. en, in; chylos, juice.] The more fluid portion of a cell; cell sap.
- encretion (ĕnkrē'shun) n. [Gk. en, within; krinein, L. cernere, to put apart.] Endocrine secretion; hormone.
- encyst (čnsist') v. [Gk. en, in; kystis, bladder.] Of a cell or small organism, to surround itself with an outer coat or capsule.
- encystation (ĕnsĭstā'shun), encystment (čnsist'mënt) n. [Gk. en, in; kystis, bladder.] Formation of a firm, resistant envelope or capsule.
- endarch (ĕnd'ârk) a. [Gk. endon, within; arche, beginning.] central protoxylem, or with several surrounding a central pith.
- endaspidean (ěnd'aspid'ean) a. [Gk. endon, within; aspis, shield.] With scutes extending on inner surface of tarsus.
- end-brain,—telencephalon, q.v.
- end-bulbs,-minute cylindrical or oval bodies, consisting of capsule containing a semi-fluid core in which axis cylinder terminates either in a bulbous extremity or in a coiled plexiform mass.

- end cell —a cell incapable of further differentiation.
- endemic (ěnděm'ík) a. [Gk. endemos, native.] Restricted to a certain region or part of a region.
- enderon (ĕn'dĕrŏn) n. [Gk. en, in; deros. skin.] The inner or endodermal layer.
- enderonic (ěn'dërŏn'ĭk) a. [Gk. en, in; deros, skin.] Endodermal.
- endites (ēndīts) n. plu. [Gk. endon, within.] Offshoots on mesial border of certain appendages of arthropods.
- endo-,-see also ento-.
- endobasal (ĕn'döbā'săl) a. [Gk. endon, within; basis, base.] Appl. body, the kinetic element of central intranuclear structure; cf. endosome.
- endoblast (ěn'döbläst) n. [Gk. endon, within; blastos, bud.] Hypoblast; coeloblast and myoblast.
- endocardiac (ĕn'dökâr'dĭāk) a. [Gk. endon, within; kardia, heart.] Situated within the heart; endocardial.
- endocardium (ĕn'dökâr'dĭŭm) n. [Gk. endon, within; kardia, heart.] The membrane which lines inner surface of heart.
- endocarp (ĕn'dokârp) n. [Gk. endon, within; karpos, fruit.] The innermost layer of pericarp, usually hard, in drupaceous fruits.
- endocarpoid (ěn'dökâr'poid) a. [Gk. endon, within; karpos, fruit; eidos, form.] Having the disc-like ascocarps embedded in the thallus.
- endochiton (ĕndökī'tŏn) n. [Gk. endon, within; chiton, coat.] Innermost layer of oogonial wall, as in Fucales; endochite; other layers: exochiton, mesochiton.
- endochondral (ěn'dökôn'drăl) a. [Gk. endon, within; chondros, cartilage.] Beginning or forming inside the cartilage, appl. ossification; cf. perichondral.
- endochondrostosis (ĕn'dökôndröstő'sis) n. [Gk. endon, within; chondros, cartilage; osteon, bone.]
 Ossification in cartilage from within outwards

- endochone (ěn'dökōnē) n. [Gk. endon, within; choane, funnel.]
 Spacious sub-cortical crypt in sponge tissue, from which arise incurrent canals.
- endochorion (ĕn'dökō'riŏn) n. [Gk. endon, within; chorion, chorion.] Inner lamina of chorion of insect eggs.
- endochrome plate,—a band of yellowish chromatophores found in protoplasmic portion of certain diatoms.
- endochromidia (ěn'dökrōmid'iă) n. plu. [Gk. endon, within; chroma, colour; idion, dim.] Metachromatic corpuscles, formed from colloidal solution of metachromatin
- endochylous (ěn'dökī'lūs) a. [Gk. endon, within; chylos, juice.] With water-cells within internal tissue.
- endocoelar (ěn'dosē'lăr) a. [Gk. endon, within; koilos, hollow.]

 Pert. inner wall of coelom, or splanchnopleure.
- endocoelic (ěn'dösē'lĭk) a. [Gk. endon, within; koilos, hollow.] In sea-anemones, appl. radial area on disc covering space between two mesenteries of the same pair; appl. inner cycle or cycles of tentacles, app. exocoelic.
- endocone (ĕn'dökōn) n. [Gk. endon, within; konos, cone.] A conical structure formed in certain cephalopod shells.
- endocranium (ĕn'dökrā'nĭŭm) n. [Gk. endon, within; kranion, skull.] Process on inner surface of cranium of certain insects; neurocranium, q.v.
- endocrine (ĕn'dökrin) n. [Gk. endon, within; krinein, to separate.] A ductless gland. a. Appl. or pert. organs of internal secretion. Opp. exocrine.
- endocrinology (ĕn'dökrinŏl'öjĭ) n. [Gk. endon, within; krinein, to separate; logos, discourse.] Study of endocrine glands and secretions, and of hormonal substances and their effects.

endocuticula (ĕn'dökūtīk'ūlă) n. [Gk. endon, within; L. dim. of cutis, skin.] The elastic inner layer of insect cuticle; inner layer of integument in spiders.

endocycle (ĕn'dösī'kl) n. [Gk. endon, within; kyklos, circle.] A layer of tissue separating internal

phloem from endodermis.

endocyclic (ěn'dösĭk'lik) a. [Gk. endon, within; kyklos, circle.] With the mouth remaining in axis of coil of gut, appl. crinoids; having an apical system with double circle of plates surrounding anus, appl. echinoids; pert. endocycle.

endocyst (ĕn'dösĭst) n. [Gk. endon, within; kystis, bladder.] The soft body wall in a polyzoan zooid; the membranous inner lining of a protozoan cyst; cf. epicyst.

- endoderm (ĕn'dödērm) n. [Gk. endon, within; derma. skin.] The hypoblast; the epithelium of digestive and respiratory organs, and of glands appended to digestive tract.
- endoderm disc,—posterior unpaired thickening on ventral surface of blastoderm of crayfish.
- endoderm lamella,—a thin sheet of endoderm stretching between adjacent radial canals, and between circular canal and enteric cavity in certain Coelenterata.
- endodermis (ĕn'dödĕr'mĭs) n. [Gk. endon, within; derma, skin.] Innermost layer of cortex in plants; layer surrounding pericycle.

endoenzyme (ĕn'döĕn'zīm) n. [Gk. endon, within; en, in; zyme, leaven.] Any intracellular enzyme.

- endogamy (ĕndög'āmī) n. [Gk. endon, within; gamos, marriage.] Zygote formation within the cyst by reciprocal fusion of division products of daughter nuclei; self-pollination; inbreeding.
- endogastric (ěn'dögăs'trĭk) a. [Gk. endon, within; gaster, belly.] Having curvature of body with enclosing shell towards ventral side; within the stomach.
- endogenous (ěndoj'ënus) a. [Gk.

endon, within; -genes, producing.] Originating within the organism; endogenic, opp. exogenous; autogenic, opp. allogenic; developing from a deep-seated layer; appl. metabolism concerned with tissue waste and growth

endogenous multiplication,—spore

formation, q.v.

endogeny (ěndőj'énĭ) n. [Gk. endon, within; genos, descent.] Development from a deep-seated layer.

- endognath (en'donâth') n. [Gk. endon, within; gnathos, jaw.] The inner branch of oral appendages of Crustacea.
- endognathion (ĕn'dönāth'iŏn) n. [Gk. endon, within; gnathos, jaw.] Mesial segment of human premaxilla.
- endogonidium (ën'dögönid'iŭm) n. [Gk. endon, within; dim. of gone, seed.] A gonidium formed in a gonidangium or receptacle; the colony-forming cells in such forms as Volvox.
- endolabium (ĕn'dölā'biūm) n. [Gk. endon, within; L. labium, lip.] A membranous lobe in interior of mouth on middle parts of front of labium.
- endolaryngeal (ěn'dölārīn'jēăl) a. [Gk. endon, within; laryngx, larynx.] Pert. or in the larynx.
- endolithic (ĕn'dölĭth'īk) a. [Gk. endon, within; lthos, stone.] Burrowing or existing in stony substratum, as algal filaments.
- endolymph (ĕn'dölimf) n. [Gk. endon, within; L. lympha, water.]

 The fluid in membranous labyrinth of ear.
- endolymphangial (ĕn'dölimfăn'jiăl)

 a. [Gk. endon, within; L. lympha,
 water; Gk. anggeion, vessel.] Situated in a lymphatic vessel.
- endolymphatic (ĕn'dölimfăt'îk) a. [Gk. endon, within; L. lympha, water.] Pert. lymphatics, or to ear labyrinth ducts.
- endolysin (ěndől'īsīn) n. [Gk. ėndon, within; lysis, loosing.] Intracellular substance of leucocytes which destroys engulfed bacteria.

- endomere (ĕn'dömēr) n. [Gk. endon, within; meros, part.] A hypoblast cell which gives rise to endoderm.
- endometrium (ĕn'dömē'trīŭm) n. [Gk. endon, within; metra, womb.] Mucous membrane lining the uterus. endomitosis (ĕn'dömītō'sīs) n. [Gk. endon, within; mitos, thread.] A form of mitosis occurring in endopolyploidy; multiplication of

A form of mitosis occurring in endopolyploidy; multiplication of chromonemata or chromosomes without division of nucleus.

endomixis (ěn'dömik'sis) n. [Gk. endon, within; mixis, mixing.] A stage comparable with partheno-

stage comparable with parthenogenesis in the reproductive rhythm of some protozoa; a type of nuclear reorganisation.

endomysium (ĕn'dömĭz'iŭm) n. [Gk. endon, within; mys, muscle.] The connective tissue binding muscle fibres.

- endoneurium (ĕn'dönū'rĭŭm) n. [Gk. endon, within; neuron, nerve.] The delicate connective tissue holding together and supporting nerve fibres within funiculus.
- endoparasite (ĕn'döpăr'ăsīt) n. [Gk. endon, within; parasitos, eating at another's table.] Any organism living parasitically within another.

endoperidium (ĕn'döpērĭd'iŭm) n. [Gk. endon, within; peridion, little pouch.] Inner layer of peridium.

- endophragm (ĕn'döfrăm) n. [Gk. endon, within; phragma, fence.]
 A septum formed by cephalic and thoracic apodemes in Crustacea.
- endophragmal (ĕn'döfrăg'măl) a. [Gk. endon, within; phragma, fence.] Pert. the endophragm.
- endophyllous (ĕn'döfĭl'ús) a. [Gk. endon, within; phyllon, leaf.] Sheathed by a leaf; living within a leaf, appl. parasites.

endophyte (čn'döfīt) n. [Gk. endon, within; phyton, plant.] A plant growing within another, either as parasite or otherwise.

endophytic (ĕn'döfīt'īk) a. [Gk. endon, within; phyton, plant.] Living in the tissues of plants.

endoplasm (ěn'döplázm) n. [Gk. endon, within; plasma, mould.]

The endosarc or inner portion of protoplasm in a cell.

endoplastule (čn'döplas'tūl) n. [Gk. endon, within; plastos, moulded.]
The micronucleus of certain Protista.

endopleura (ĕn'döploo'ră) n. [Gk. endon, within; pleura, side.] The inner seed-coat or tegmen.

endopleurite (en'döploo'rīt) n. [Gk. endon, within; pleura, side.] The epimeral portion of an apodeme; infolding between pleurites.

endopodite (ĕn'dŏpödīt) n. [Gk. endon, within; pous, foot.] The inner or mesial branch of a biramous crustacean limb, or the only part of biramous limb remaining.

- endopolyploidy (ĕn'döpŏl'ĭploidĭ) n.
 [Gk. endon, within; polys, many; aploos, onefold; eidos, form.]
 Polyploidy resulting from repeated doubling of chromosome number without normal mitosis.
- endoral (ĕndō'rāl) a. [Gk. endon, within; L. os, mouth.] Pert. structures situated in the vestibule of certain protozoa.
- endorhachis (ĕn'dörā'kis) n. [Gk. endon, within; rhachis, backbone.] A layer of connective tissue lining canal of vertebral column and cavity of skull.
- endosarc (ĕn'dösârk) n. [Gk. endon, within; sarx, flesh.] Endoplasm, q.v.
- endoscopic (ěn'döskŏp'ĭk) a. [Ġk. endon, within; skopein, to look.] With apex directed inwards toward base of archegonium, appl. embryo. Opp. exoscopic.
- endosiphuncle (čn'dösī'fŭngkl) n. [Gk. endon, within; L. siphunculus, little tube.] The tube leading from protoconch to siphuncle in certain Cephalopoda.

endoskeleton (ĕn'döskĕl'čtŏn) n. [Gk. endon, within; skeletos, dried up.] Internal skeleton, opp. exoskeleton.

endosmosis (ĕn'dŏsmō'sĭs) n. [Gk. endon, within; osmos, impulse.] The passage inwards through a permeable or semipermeable membrane, of a less concentrated solution. Opp. exosmosis.

- endosome (ĕn'dösōm) n. [Gk. endon, within; soma, body.] Chromatinic mass near centre of a vesicular nucleus; karyosome.
- endosperm (ěn'döspěrm) n. [Gk. endon, within; sperma, seed.] The nutritive tissue of certain seeds; nutritive residue of female prothallus surrounding an embryo.
- endospore (ĕn'döspōr), endosporium (ĕn'döspō'rĭŭm) n. [Gk. endon, within; sporos, seed.] Inner coat of sporocyst in some protozoa; an asexual spore; a sporangial spore; the inner coat of a spore wall.
- endosteal (ĕndŏs'tĕăl) a. [Gk. endon, within; osteon, bone.] Pert. endosteum.
- endosternite (ĕn'döstĕr'nīt) n. [Gk. endon, within; L. sternum, sternum.] Internal skeletal plate for muscle attachment; median sternal apodeme; a free skeleton situated in prosoma between alimentary canal and nerve cord in arachnids.
- endosteum (ĕndŏs'tëŭm) n. [Gk. endon, within; osteon, bone.] The internal periosteum lining the cavities of bones.
- endostosis (ĕn'dŏstō'sĭs) n. [Gk. endon, within; osteon, bone.] Ossification which begins in cartilage.
- endostracum (ĕndős'trākŭm) n. [Gk. endon, within; ostrakon, shell.]
 The inner layer of mollusc shell.
- endostyle (ěn'dostīl) n. [Gk. endon, within; stylos, pillar.] A band of thickened epithelium on oesophageal wall of a tornaria; two ventral longitudinal folds separated by a groove in pharynx of Tunicata; a longitudinal groove lined by ciliated epithelium on ventral wall of pharynx of Amphioxus; precursor of thyroid gland.
- endotergite (ěn'dötěr'jīt, -gīt) n. [Gk. endon, within; L. tergum, back.] An infolding from a tergite of insects, for muscle attachment; phragma.
- endotheca (ěn'döthē'kă) n. [Gk. endon, within; theke, box.] The system of dissepiments in a coral calyx; the oval surface of Cystidea.

- endothecial (ĕn'döthē'sīăl) a. [Gk. endon, within; theke, box.] Pert. endothecium; with asci in an ascocarp.
- endothecium (ĕn'döthē'sĭūm) n. [Gk. endon, within; theke, box.] The central region of an epibasal octant of oospore of liverworts and mosses; inner lining of an anther; inner dehiscing layer in ginko and angiosperms.
- endotheliocyte (ĕn'döthē'liösīt) n. [Gk. endon, within; thele, nipple; kytos, hollow.] A mononuclear phagocyte derived from endothelium; endothelial phagocyte or primitive wandering cell; a histiocyte; a macrophage.
- endothelium (ĕn döthē'lĭum) n. [Gk. endon, within; thele, nipple.] A squamous epithelium which lines serous cavities, the heart, blood and lymphatic vessels.
- endothorax (ĕn'dothō'rāks) n. [Gk. endon, within; thorax, chest.] The apodeme system in a crustacean thorax; cf. entothorax.
- endotoxin (ĕn'dotŏk'sīn) n. [Gk. endon, within; toxikon, poison.] A toxin within bacterial protoplasm; opp. exotoxin.
- endotrachea (ĕn'dötrākē'ā) n. [Gk. endon, within; L. trachea, windpipe.] The innermost, chitinous coat of tracheal tubes of insects.
- endotrophic (ĕn'dotrŏf'ik) a. [Gk. endon, within; trophe, nourishment.] Appl. space within peritrophic membrane of insects; finding nourishment from within; appl. fungi inhabiting root cortex of host. Opp. ectotrophic.
- endozoochore (ĕn'dözō'ökōr) n. [Gk. endon, within; zoon, animal; chora, place.] Any spore, seed, or organism dispersed by being carried within an animal. Opp. epizoochore.
- end-plates,—motor end-organs, the ramified expansions within the muscular fibre which form the ends of a motor nerve.
- end-sac,—the sac-like vestigial portion of coelom in excretory glands of certain Crustacea.

- endysis (ĕn'disīs) n. [Gk. endysis, putting on.] The development of a new coat; ef. ecdysis.
- energesis (ĕn'ërjë'sĭs) n. [Gk. energein, to be active.] The process by which energy is liberated through katabolic action.
- energid (ĕn'ërjĭd) n. [Gk. energos, working; idion, dim.] Any living uninucleated protoplasmic unit with or without a cell wall.
- enervose (ĕnër'vōs) a. [L. ex, without; nervus, sinew.] Having no veins, appl. certain leaves.
- engram (ĕn'grăm) n. [Gk. en, in; graphein, to write.] A character impression in the mnemic theory of heredity; a latent memory image.
- engraved (engrav'd) a. (F. en, in; A.S. grafan, to dig.] With irregular linear grooves on the surface.
- ensiform (ĕn'sifôrm) a. [L. ensis, sword; forma, shape.] Swordshaped; xiphoid.
- entad (ěn'tăd) adv. [Gk. entos, within; L. ad, towards.] Towards the interior; inwards; internally. Opp. ectad.
- ental (ĕn'tăl) a. [Gk. entos, within.] Inner; internal. Opp. ectal.
- entelechy (ĕntĕl'ĕkĭ) n. [Gk. en, in; telos, end; echein, to hold.] Vital principle or influence guiding living organisms in right direction.
- entepicondylar (ěnt'épikön'dilăr) a. [Gk. entos, within; epi, upon; kondylos, knob.] Pert. lower or condylar end of humerus; appl. ulnar foramen.
- enteral (ĕn'tĕrăl) a. [Gk. enteron, gut.] Within intestine; also appl. the parasympathetic portion of the autonomic nervous system.
- enteric (ĕntĕr'ik) a. [Gk. enteron, gut.] Pert. alimentary canal.
- enteroblast (ĕn'tëröblăst) n. [Gk. enteron, gut; blastos, bud.] The hypoblast after formation of the mesoblast.
- enterocoel (ěn'těrösěl') n. [Gk. enteron, gut; koilos, hollow.] A coelom arising as a pouch-like outgrowth of archenteric cavity, or as a series of such outgrowths.

- enterocrinin (ĕn'terökrī'nīn) n. [Gk. enteron, gut; krinein, to separate.] A hormone of small intestine, which stimulates secretion of intestinal juice.
- enteroderm (ěn'teröderm) n. [Gk. enteron, gut; derma, skin.] Enteroblast.
- enterogastrone (ěn'tërögăs'trōn) n.
 [Gk. enteron, gut; gaster, stomach.]
 A duodenal hormone which inhibits
 secretion and motility of stomach.
- enterokinase (en'tërökinās') n. [Gk. enteron, gut; kinein, to move.] Incomplete enzyme of intestinal juice which converts trypsinogen into trypsin.
- enteron (ěn'tërŏn) n. [Gk. enteron, gut.] The alimentary tract.
- enteronephric (ĕn'tërönĕf'rīk) a. [Gk. enteron, gut; nephros, kidney.] With nephridia opening into gut; opp. exonephric, appl. Oligochaeta.
- enteroproct (ĕn'teröprökt) n. [Gk. enteron, gut; proktos, anus.] The opening from endodermal gut into proctodaeum.
- enterostome (ĕn'tëröstōm) n. [Gk. enteron, gut; stoma, mouth.] The aboral opening of the actinopharynx, leading to coelenteron; the posterior opening of stomodaeum into endodermal gut.
- enterosympathetic (čn'tërösimpăthět'îk) a. [Gk. enteron, gut; syn, with; pathos, feeling.] Appl. that part of the nervous system supplying the intestine.
- enterozoon (ĕn'tërözō'ŏn) n. [Gk. enteron, gut; zoon, animal.] Any animal parasite inhabiting intestines.
- entire (ĕntīr') a. [M.E. enter, untouched.] Unimpaired; with continuous margin, appl. leaves, bacterial colony.
- ento,-see also endo-.
- entobranchiate (ĕn'töbrăng'kiāt) a.
 [Gk. entos, within; brangchia, gills.] Having internal gills.
- entobronchus (ĕn'töbrŏng'kŭs) n.
 [Gk. entos, within; brongchos,
 windpipe.] The dorsal secondary
 branch of bronchus in birds;
 entobronchium.

- entochondrite (ěn'tökôn'drīt) n. [Gk. entos, within; chondros, cartilage.]
 Plastron or endosternum of Limulus.
- entochondrostosis (ĕn'tökŏndrŏstō'sis) n. [Gk. entos, within; chondros, cartilage; osteon, bone.] Ossification from within outwards.
- entocodon (ĕn'tokō'dŏn) n. [Gk. entos, within; kodon, bell.] The lens-shaped mass of cells, in development of medusoid, which sinks below level of superficial ectoderm, and ultimately develops a cavity.
- entocoel (ĕn'tosēl) n. [Gk. entos, within; koilos, hollow.] The space enclosed by a pair of mesenteries in Anthozoa.
- entocondyle (ĕn'tokŏn'dil) n. [Gk. entos, within; kondylos, knob.]
 Condyle on mesial surface of a bone.
- entoconid (ĕn'tökō'nĭd) n. [Gk. entos, within; konos, cone.] The postero-internal cusp of a lower molar.
- entocuneiform (ĕn'tökūnē'ifôrm) n. [Gk. entos, within; kuneos, wedge; L. forma, shape.] The most internal of distal tarsal bones.
- entocyemate (ĕn'tösīē'māt) a. [Gk. entos, within; kyema, cmbryo.] With embryos having amnion and allantois.
- entoderm(al),—endoderm(al), q.v. entogastric (ĕn'tögăs'trīk) a. [Gk. entos, within; gaster, belly.] Pert. interior of stomach; appl. gastric budding in medusae.

entoglossal (ĕn'toglŏs'āl) a. [Gk. entos, within; glossa, tongue.] Lying in substance of tongue.

- entoglossum (ĕn'töglös'um) n. [Gk. entos, within; glossa, tongue.] Extension of basihyal into tongue in some fishes; also glossohyal.
- entomogenous (ĕn'tomŏj'ĕnŭs) a. [Gk. entomon, insect; genos, birth.] Growing on insects, as certain fungi.
- entomology (ĕn'tömöl'öji) n. [Gk. entomon, insect; logos, discourse.] That part of zoology which deals with insects.
- entomophagous (ĕn'tömŏf'ăgus) a.

- [Gk. entomon, insect; phagein, to eat.] Insect-eating; insectivorous.
- entomophilous (ĕn'tomŏf'ilŭs) a. [Gk. entomon, insect; philein, to love.] Pollinated by agency of insects.
- entoneural (en'tonu ral) a. [Gk. entos, within; neuron, nerve.]

 Appl. system of aboral ring and genital nerves in echinoderms.

entophyte,—endophyte, q.v.

entopic (ěntop'ik) a. [Gk. en, in; topos, place.] In normal position, opp. ectopic.

entoplasm, —endoplasm, q.v.

entoplastron (ěn'töplās'trŏn) n. [Gk. entos, within; F. plastron, breastplate.] The anterior median plate in chelonian plastra, often called episternum, probably homologous with interclavicle of other reptiles.

entopterygoid (en'topter'igoid) n. [Gk. entos, within; pteryx, wing; eidos, form.] A dorsal membrane bone behind the palatine in some fishes; cf. ectopterygoid.

entosphere (ĕn'tösfēr) n. [Gk. entos, within; sphaira, globe.] The inner portion of attraction sphere.

entosternite, -- endosternite, q.v.

entosternum (ĕntöstĕr'nŭm) n. [Gk. entos, within; L. sternum, breastbone.] Entoplastron, q.v.; an internal process of sternum of numerous arthropods.

entothorax (ĕn'töthō'rāks) n. [Gk. entos, within; thorax, chest.] An insect apophysis or sternite.

entoturbinals (en'totur'binalz) n. plu.
[Gk. entos, within; L. turbo, whorl.]
A division of ethmoturbinals.

entotympanic (ĕn'tötimpăn'ik) n.
[Gk. entos, within; tympanon,
drum.] A separate tympanic
element in some genera; also
metatympanic.

entovarial (ĕnt'ōvā'rīāl) a. [Gk. entos, within; L. ovum, egg.] Pert. canal formed in ovaries of some fishes by insinking and closure of a groove formed by covering epithelium.

entozoa (ĕn'tözō'ā) n. plu. [Gk. entos, within; zoon, animal.]
Internal animal parasites.

- entozoic (ĕn'tözō'ĭk) a. [Gk. entos, within; zoe, subsistence.] Living within the body or substance of another animal or plant; pert. entozoa.
- entrochite (ĕn'trōkīt) n. [Gk. en, in; trochos, wheel.] The joint of fossil stem of a stalked crinoid.
- enucleate (ēnū'kleāt) v. [L. e, out of; nucleus, kernel.] To deprive of a nucleus, as in microdissection of cells. a. Lacking a nucleus.

envelope (ĕn'vĕlōp) n. [F. enveloppe, covering.] An outer covering of an egg; any surrounding structure,

e.g. floral envelope.

- environment (envi/ronment) n. [F. environ, about.] The sum-total of external influences acting on an organism or on part of an organism.
- enzootic (ĕn'zōŏt'îk) a. [Gk. en, in; zoon, animal.] Afflicting animals; appl. disease restricted to a locality.
- enzyme (ěn'zīm) n. [Gk. en, in, zyme, leaven.] A catalyst produced by living organisms and acting on one or more specific substrates; a ferment; cf. apo-enzyme, coenzyme.
- Eocene (ē'ösēn) n. [Gk. eos. dawn; kainos, recent.] Early epoch of the Tertiary period, between Palaeocene and Oligocene.
- Eogaea (ē'öjē'à) n. [Gk. eos, dawn; gaia, land.] A zoogeographical division including Africa, South America, and Australasia; cf. Caenogaea.
- eosinophil (ē'ösĭn'öfĭl) a. [Gk. eos. dawn; philein, to love.] Appl. cells which readily stain red with eosin; oxyphil.

Eozoic (ē'özō'îk) a. [Gk. eos, dawn; zoe, life.] Appl. Archaean or Pre-Cambrian period.

epacme (ĕpāk'mē) n. [Gk. epi, upon; akme, prime.] The stage in phylogeny of a group just previous to its highest point of development.

epalpate (ēpāl'pāt) a. [L. ex, without; palpus, palp.] Not furnished

with palpi.

- epanthous (ĕpăn'thŭs) a. [Gk. epi, upon; anthos, flower.] Living on flowers; appl. certain fungi.
- epapillate (ēpāp'ilāt) a. [L. ex, without; papilla, nipple.] Not having papillae.
- epapophysis (ĕp'apŏf'isis) n. [Gk. epi, upon; apophysis, offshoot.] A median process arising from centre of vertebral neural arch.
- eparterial (ĕp'ârtē'rīăl) a. [Gk. epi, upon; L. arteria, artery.] Situated above an artery; appl. branch of right bronchus.
- epaulettes (ĕp'ôlĕts) n. plu. [F. épaule, shoulder.] Branched or knobbed processes projecting from outer side of oral arms of many Scyphozoa; crescentic ridges of cilia in echinopluteus.
- epaxial (ĕpăk'sĭāl) a. [Gk. epi, upon; L. axis, axle.] Above the axis; dorsal; usually appl. axis formed by vertebral column.
- epedaphic (čp'ëdăf'îk) a. [Gk. epi, upon; edaphos, soil.] Pert., or depending upon, climatic conditions.
- epencephalon (ĕp'ĕnkĕf'álön, -sĕf-)
 n. [Gk. epi, upon; engkephalos, brain.] The cerebellum.
- ependyma (ĕpĕn'dimā) n. [Gr. ependyma, outer garment.] The layer of cells lining cavities of brain and spinal cord; ependyme.
- ependymal (ĕpĕn'dimăl) a. [Gk. ependyma, outer garment.] Pert. ependyma.
- ephapse (ĕfăps') n. [Gk. ephaptein, to reach.] Region of contiguity between two axons lying side by side.
- ephaptic (ĕfāp'tīk) a. [Gk. ephaptein, to reach.] Pert. an ephapse; appl. delay, the interval between stimulation of one (pre-ephaptic) axon and response of an apposed other (post-ephaptic) axon.
- epharmonic (ĕf'armŏn'Ik) a. [Gk. epi, towards; harmos, fitting.] Pert. epharmosis; adaptive; adapted to environment; appl. convergence: morphological resemblance of different species inhabiting the same environment.

epharmosis (ĕf'armō'sīs) n. [Gk. cpi, towards; harmos, fitting.]
The process of adaptation of organisms to new environmental conditions; attainment of the state of adaptation or epharmony.

ephebic (ĕfē'bīk) a. [Gk. ephebos, adult.] Adult; pert. stage in development or phylogeny between childhood and old-age

stages.

ephemeral (ĕfĕm'ērāl) n. [Gk. ephemeros, lasting for a day.] A short-lived plant or animal species. a. Short-lived; taking place once only, appl. plant movements, as expanding of buds; completing life cycle within a brief time period.

ephippial (ĕfip'ĭăl) a. [Gk. ephippion, saddle-cloth.] Pert. ephippium; appl. winter eggs, as of

rotifers and daphnids.

- ephippium (ĕfip'iŭm) n. [Gk. ephippion, saddle-cloth.] The pituitary fossa, or fossa hypophyseos of sphenoid; a thickened and indurated part of shell separating from the rest at ecdysis; a saddleshaped modification of cuticle derived, later detached, from carapace and enclosing winter eggs, in Daphniidae.
- ephyra (ĕf'irā), ephyrula (ĕfir'ūlā)
 n. [Gk. Ephyra, nymph's name.]
 The small free-swimming jelly-fish stage of certain Scyphozoa, produced by strobilation of scyphistoma.
- epibasal (ĕp'ībā'săl) n. [Gk. epi, upon; basis, base.] Upper segment of an oospore, ultimately giving rise to the shoot. Opp. hypobasal.
- epibenthos (ĕp'Ibĕn'thŏs) n. [Gk. epi, upon; benthos, depths.] Fauna and flora found between lowwater mark and hundred fathom line.
- epibiotic (ĕp'ibiŏt'ik) a. [Gk. epibionai, to survive.] Surviving, appl. endemic species that are relics of a former flora or fauna.

- epiblast (ĕp'ĭblăst) n. [Gk. epi, upon; blastos, bud.] The outer layer of the gastrula; ectoblast; a rudimentary second cotyledon, as in grasses.
- epiblema (ĕp'iblē'mā) n. [Gk. epiblema, cover.] The outermost layer of root-tissue; piliferous layer; epiblem.
- epibole (ĕpib'ölē) n. [Gk. epibole, putting on.] Growth of one part over another in embryonic stages; also epiboly.
- epibolic (ĕp'ĭbŏl'ĭk) a. [Gk. epibole, putting on.] Growing so as to cover over; appl. type of gastrulation.
- epibranchial (ĕp'ibrāng'kiăl) a. [Gk. epi, upon; brangchia, gills.] Pert. second upper element in branchial arch; efferent branchial, appl. vessels.
- epicalyx (ĕp'îkăl'îks) n. [Gk. epi, upon; kalyx, cup.] Stipules, fused in pairs, producing an apparent outer or extra calyx; structure just below calyx produced by aggregation of bracts or bracteoles.
- epicanthus (ĕp'ĭkăn'thŭs) n. [Gk. epi, upon; kanthos, corner of eye.] A prolongation of upper eyelid over inner angle of eye; Mongolian fold.
- epicardium (ĕp'ĭkâr'dĭŭm) n. [Gk. epi, upon; kardia, heart.] The visceral part of pericardium; tubular prolongation of branchial sac in many ascidians, which takes part in budding.

epicarp (ĕp'îkârp) n. [Gk. epi, upon; karpos, fruit.] Outer layer of the

pericarp; exocarp.

epicentral (ĕp'īsĕn'trăl) a. [Gk. epi, upon; kentron, centre.] Attached to or arising from vertebral centra; appl. intermuscular bones.

epicerebral (ĕp'ĭsĕr'ĕbrăl) a. [Gk. epi, upon; L. cerebrum, brain.]

Situated above the brain.

epichilium (ĕp'îkîl'ĭum) n. [Gk. epi, upon; cheilos, lip.] Terminal lobe of lower petal of orchid; epichile.

- epichordal (ĕp'īkôr'dāl) a. [Gk. epi, upon; chorde, cord.] Upon the notochord; appl. vertebrae in which ventral cartilaginous portions are almost completely suppressed; appl. upper lobe of caudal fin in fishes.
- epicoel (ĕp'isēl) n. [Gk. epi, upon; koilos, hollow.] Cavity of mid-brain in lower vertebrates; cerebellar cavity; a perivisceral cavity formed by invagination; also epicoele, epicoelia.

epicondylar (ěp'íkŏn'dĭlār) a. [Gk. epi, upon; kondylos, knob.] Pert. epicondyle.

epicondyle (ĕp'ikŏn'dil) n. [Gk. epi, upon; kondylos, knob.] A medial and a lateral protuberance at distal end of humerus and femur.

epicone (ĕp'īkōn) n. [Gk. epi, upon; konos, cone.] The part anterior to girdle in Dinoflagellata, opp. hypocone.

- epicoracoid (ĕp'ĭkŏr'ákoid] a. [Gk. epi, upon; korax, crow; eidos, form.] Pert. an element, usually cartilaginous, at sternal end of coracoid in amphibians, reptiles, and monotremes.
- epicormic (ĕp'īkôr'mĭk) a. [Gk. epi, upon; kormos, trunk.] Growing from a dormant bud.
- epicotyl (ĕp'ikŏt'il) n. [Gk. epi, upon; kotyle, vase.] The axis of a plumule.
- epicotyledonary (ěp'îkŏtilē'dönări)
 a. [Gk. epi, upon; kotyle, cup.]
 Above the cotyledons.
- epicoxite (ĕp'îkŏk'sīt) n. [Gk. epi, upon; L. coxa, hip.] A small process at posterior end of toothed part of coxa of second to fifth pairs of appendages in Eurypterida.

epicranial (ĕp'ĭkrā'nĭāl) a. [Gk. epi, upon; kranion, skull.] Pert. cranium; appl. aponeurosis, muscles, bones, suture.

epicranium (ĕp'ĭkrā'nĭūm) n. [Gk. epi, upon; kranion, skull.] The region between and behind eyes in insect head; scalp; the structures covering the cranium.

epicranius (ĕp'ikrā'niŭs) n. [Gk. epi, upon; kranion, skull.] The scalp

muscle, consisting of occipitalis and frontalis, connected by galea aponeurotica; occipitofrontalis.

epicrine (ĕp'ikrīn) a. [Gk. epi, upon; krinein, to separate.] Appl. glands in which secretion is voided without disintegration of cells

disintegration of cells.

epicritic (ĕp'ĭkrĭt'ĭk) a. [Gk. epi, upon; krinein, to judge.] Appl. stimuli and nerve systems concerned with delicate touch and other special sensations in skin.

epictesis (ĕpĭt'ēsīs, ĕpĭtē'sīs, -ktē-) n. [Gk. epiktesis, further gain.] Capacity of a living cell to concentrate salt solutions diffusing into the cell.

epicuticula (ĕp'ĭkū'tĭk'ūlă) n. [Gk. epi, upon; L. dim. of cutis, skin.] Lamella or membrane external to exocuticle of insects.

epicyst (ĕp'isist) n. [Gk. epi, upon; kystis, bladder.] The external resistant cyst of an encysted protozoan; ef. endocyst.

epicyte (ep'isit) n. [Gk. epi, upon; kytos, hollow.] The external layer of ectoplasm in certain protozoa.

- epidemes (řp'ídēmz) n. plu. [Gk. epi, upon; demas, body.] In certain insects, small pieces closely related with articulation of wings.
- epidermis (ĕp'īděr'mĭs) n. [Gk. epi, upon; derma, skin.] The outermost protective layer of stems, roots and leaves; scarf-skin or external layer of skin, a non-vascular stratified epithelium of ectodermic origin; single layer of ectoderm in invertebrates.
- epididymis (ĕp'idid'imis) n. [Gk. epi, upon; didymos, testicle.] A mass at back of testicle composed chiefly of vasa efferentia; the coiled anterior end of Wolffian duct.

epidural (ĕp'ĭdū'răl) a. [Gk. epi, upon; L. dura, hard.] Pert. dura mater; appl. space between dura mater and wall of vertebral canal.

epigamic (ĕp'īgăm'īk) a. [Gk. epi, upon; gamos, marriage.] Tending to attract opposite sex, e.g. colour displayed in courtship. epigamous (ĕpĭg'ămŭs) a. [Gk. epi, upon; gamos, marriage.] Designating that stage in polychaetes in which immature forms become heteronereid, while sexual elements are ripening; epigamic. q.v.

epigaster (ĕp'igăs'ter) n. [Gk. epi, upon; gaster, belly.] That part of embryonic intestine which later

develops into colon.

epigastric (ĕp'īgăs'trĭk) a. [Gk. epi, upon; gaster, belly.] Pert. anterior wall of abdomen; middle region of upper zone of artificial divisions of abdomen.

epigastrium (ĕp'igăs'triŭm) n. [Gk. epi, upon; gaster, stomach.] The epigastric region; sternal portions of meso- and metathorax of insects.

epigastroid,—epipubis, q.v.

epigeal (ĕp'ijē'āl) a. [Gk. epi, upon; ge, earth.] Living near the ground, appl. insects; borne above ground, appl. cotyledons when they form first foliage leaves; also epigean, epigeic, epigeous.

epigenesis (ep'ijen'esis) n. [Gk. epi, upon; genesis, descent.] Theory of generation, that embryo is an entirely new creation, not a mere unfolding of preformed structures.

epigenetics (ep'ijenet'iks) n. [Gk. epi, upon; genesis, descent.] Study of the mechanisms causing phenotypic effects to be produced by the genes of a genotype.

epigenotype (ĕp'ijĕn'ötīp) n. [Gk. epi, upon; genos, descent; typos, image.] The concatenation of processes linking genotype and phenotype.

epigenous (ěpij'ënus) a. [Gk. epi, upon; genos, descent.] Developing

or growing on a surface.

epiglottis (ĕp'iglöt'is) n. [Gk. epi, upon; glotta, tongue.] A thin lamella of fibro-cartilage between root of tongue and entrance to larynx; epistome in Polyzoa; epipharynx in Insecta.

epignathous (ĕpignā'thūs) a. [Gk. epi, upon; gnathos, jaw.] Having upper jaw longer than lower.

epigone,-epigonium.

epigonial (ĕpĭgō'nĭăl) a. [Gk. epi, upon; gone, seed.] Appl. sterile posterior portion of genital ridge.

epigonium (epigo'nium) n. [Gk. epi, upon; gone, seed.] The young sporangial sac in liverworts.

epigynal (ĕpĭj'ĭnăl) a. [Gk. epi, upon; gyne, woman.] Pert. epigynum.

epigynous (epij'īnūs) a. [Gk. epi, upon; gyne, woman.] Having the various whorls adnate to ovary, thus apparently inserted in ovary.

epigynum (ĕpiji'num) n. [Gk. epi, upon; gyne, woman.] External female genitalia in Arachnida;

also epigyne, epigynium.

epigyny (ĕpij'ini) n. [Gk. epi, upon; gyne, woman.] Condition of having whorls apparently inserted in ovary.

epihyal (ĕp'îhî'ăl) a. [Gk. epi, upon; hyoeides, T-shaped.] Pert. upper portion of ventral part of hyoid arch. n. Upper element of ventral portion, a cartilage or bone in centre of stylohyoid ligament.

epilabrum (ĕpĭlā'brūm) n. [Gk. epi, upon; L. labrum, lip.] A process at side of labrum in Myriapoda.

- epilemmal (ěpilém'ál) a. [Ġk. epi, upon; lemma, skin.] Appl. sensory nerve endings on surface of sarco-lemma.
- epilimnion (ĕp'ilimni'ŏn, -lim'nyŏn)

 n. [Gk. epi, upon; limnaios, stagnant; limne, marshy lake.] Upper
 water layer, above thermocline, in
 lakes. Opp. hypolimnion.

epilithic (epilith'ik) a. [Gk. epi, upon; lithos, stone.] Attached on

rocks; appl. algae.

epimandibular (ĕp'imăndib'ūlăr) a. [Gk. epi, upon; L. mandibulum, jaw.] Pert. a bone in lower jaw of vertebrates.

epimeral (ĕpimē'rāl) a. [Gk. epi, upon; meros, thigh.] Pert. epimeron.

epimere (ĕp'imēr) n. [Gk. epi, upon; meros, part.] The dorsal muscle-plate of mesothelial wall.

epimerite (ĕp'Imërīt) n. [Gk. epi, upon; meros, part.] Deciduous portion of protomerite in certain Gregarinina.

- epimeron (ĕpǐmē'rŏn) n. [Gk. epi, upon; meros, thigh.] A portion of pleuron in insects which may be posterior or nearly as far forward as episternum; posterior pleurite of subcoxa; portion of arthropod segment between tergum and limb insertions.
- epimorpha (ĕp'imôr'fă) n. plu. [Gk. epi, upon; morphe, form.] Larvae hatched with all appendages developed; cf. anamorpha.

epimorphosis (ĕp'imôr'fōsīs) n. [Gk. epi, upon; morphosis, shaping.]
That type of regeneration in which proliferation of new material precedes development of new part.

epimysium (ep'imiz'iŭm) n. [Gk. epi, upon; mys, muscle.] The sheath of areolar tissue which invests the entire muscle; cf. perimysium.

- epinasty (ĕp'inăsti) n. [Gk. epi, upon; nastos, close-pressed.] The more rapid growth of upper surface of a dorso-ventral organ, e.g. a leaf, thus causing unrolling or downward curvature.
- epinephrine (ĕp'inĕf'rēn) n. [Gk. epi, upon; nephros, kidney.] Adrenaline; adrenin.
- epinephros (ĕp'inēf'rōs) n. [Gk. epi, upon; nephros, kidney.] The suprarenal or adrenal body.
- epineural (ĕp'inū'răl) a. [Gk. epi, upon; neuron, nerve.] Arising from vertebral neural arch; pert. canal external to radial nerve in certain echinoderms; appl. sinus between embryo and yolk, beginning of body cavity in insects.
- epineurium (ĕp'ĭnū'rĭūm) n. [Gk. epi, upon; neuron, nerve.] The external sheath of a nerve cord.
- epinotum (ĕp'inō'tūm) n. [Gk. epi, upon; noton, back.] Propodeon, q.v. epiopticon (ĕp'iŏp'tīkŏn) n. [Gk. epi, upon; opsis, sight.] The middle zone of optic lobes of insects.
- epiotic (ĕp'iōt'ik) a. [Gk. epi, upon; ous, the ear.] Pert. upper element of bony capsule of ear; appl. centre of ossification of mastoid process.

epiparasite (ĕp'ipăr'ăsīt) n. [Gk. epi, upon; parasitos, eating at another's table.] Ectoparasite, q.v.

epipelagic (ĕp'ĭpĕlăj'ĭk) a. [Gk. epi, upon; pelagos, sca.] Pert. deepsea water between surface and bathypelagic zone.

epiperidium,—exoperidium, q.v.

- epipetalous (ĕpi'pĕtălūs) a. [Gk. epi, upon; petalon, petal.] Having stamens inserted on petals.
- epipharyngeal (čp'ĭfărĭn'jeäl) a. [Gk. epi, upon; pharyngx, throat.]

 Pert. upper or dorsal aspect of pharynx.
- epipharynx (ĕp'ifăr'ingks) n. [Gk. epi, upon; pharyngx, throat.] A projection on roof of mouth cavity of certain insects; membranous lining of labrum and clypeus drawn out with labrum to form a piercing organ, as in Diptera; lingua.

epiphloem (ĕpĭflō'ĕm) n. [Gk. epi, upon; phloios, bark.] Outer bark.

- epiphragm (ĕp'ĭfrăm) n. [Gk. epiphragma, covering.] A layer of hardened mucous matter, or a calcareous plate, closing the opening of certain gastropod shells; membrane which closes the capsule in certain mosses; a closing membrane in sporophores of certain fungi.
- epiphyll (ep'ifil) n. [Gk. epi, upon; phyllon, leaf.] A plant which grows on leaves, e.g. various lichens.
- epiphyllous (ĕp'ĭfĭl'ūs) a. [Gk. epi, upon; phyllon, leaf.] Growing on leaves; united to perianth, appl. stamens.
- epiphysial (ĕp'ifiz'iāl) a. [Gk. epi, upon; phyein, to grow.] Pert. or similar to the epiphysis; epiphyseal.
- epiphysis (ĕpif'isis) n. [Gk. epi, upon; phyein, to grow.] Any part or process of a bone which is formed from a separate centre of ossification and later fuses with the bone; pineal body; pineal and parapineal organs; stout bar firmly fused to alveolus of each jaw and articulating with rotulae in sea-urchins; certain processes on tibia of insects; caruncle near hilum of seed.

- epiphyte (ĕp'ĭfit) n. [Gk. epi, upon; pluyton, plant.] Plant which lives on surface of other plants.
- epiphytic (ĕp'ifit'ik) a. [Gk. epi, upon; phyton, plant.] Living on, or attached to, surface of a plant, opp. endophytic; pert. or similar to an epiphyte.

epiphytotic (ĕpi'fĭtŏt'ĭk) a. [Gk. epi, upon; phyton, plant.] Pert. disease

epidemic in plants.

- epiplankton (ĕp'iplăng'ktŏn) n. [Gk. epi, upon; plangktos, wandering.] That portion of plankton from surface to one hundred fathoms.
- epiplasm (ĕp'iplăzm) n. [Gk. epi, upon; plasma, mould.] Cytoplasm of a brood mother-cell remaining unused in brood formation; cytoplasm of ascus remaining after spore formation.
- epiplastron (ĕp'ĭplăs'trŏn) n. [Gk. epi, upon; F. plastron, breastplate.] One of anterior pair of bony plates in plastron of Chelonia.
- epipleura (ĕp'īploo'ră) n. [Gk. epi, upon; pleura, rib.] Epithecal part of cingulum in diatoms; one of rib-like structures in telcosts which are not preformed in cartilage; an uncinate process in birds; the turned down outer margin of elytra of certain beetles.
- epiploic (čpřplo'ík) a. [Gk. epiploon, caul of entrails.] Pert. omentum.
- epiploic foramen,—opening between bursa omentalis and large sac of peritoneum; foramen of Winslow.
- epiploon (ĕpĭp'lōŏn) n. [Gk. epiploon, caul of entrails.] Great omentum; insect adipose tissue.
- epipodial (ĕp'ĭpō'dĭāl) a. [Gk. epi, upon; pous, foot.] Pert. epipodium.
- epipodite (ĕp'īpödīt) n. [Gk. epi, upon; pous, foot.] A process arising from basal joint of crustacean limb and usually extending into gill chamber.
- epipodium (ĕp'ĭpō'dĭŭm) n. [Gk. epi, upon; pous, foot.] The leafblade or lamina; embryonic leaflamina; ridge, fold, or lobe along edge of foot of Gastropoda; raised

- ring on an ambulacral plate in Echinoidea.
- epiprecoracoid (ĕp'iprēkŏr'ākoid) n. [Gk. epi, upon; L. prae, before; Gk. korax, crow; eidos, form.] A small cartilage at ventral end of precoracoid in pectoral girdle in some Chelonia.
- epiproct (ĕp'iprökt) n. [Gk. epi, upon; proktos, anus.] A supra-anal plate representing tergum of tenth or eleventh segment in some insects.
- epipteric (ĕp'iptĕr'ik) a. [Gk. epi, upon; pteron, wing.] Pert. or shaped like, or placed above wing; appl. a small skull bone between parietal and sphenoidal ala.

epipterygoid (ĕp'ĭtĕr'igoid) n. [Gk. epi, upon; pteryx, wing.] A small bone extending nearly vertically downwards from prootic to pterygoid; also columella cranii.

- epipubic (ĕp'îpū'bik) a. [Gk. epi, upon; L. pubes, adult.] Pert. or borne upon pubis; appl. certain cartilages or bones principally in marsupials; appl. anterior median process of ischiopubic plate.
- epipubis (ĕp'ĭpū'bis) n. [Gk. epi, upon; L. pubes, adult.] Unpaired cartilage or bone borne anteriorly on pubis; also epigastroid.
- epirhizous (ĕp'ĭrī'zŭs) a. [Gk. epi, upon; rhiza, root.] Growing upon a root.
- epirrhysa (čpĭrī'să) n. plu. [Gk. epirrhein, to flow into.] Inhalant canals in sponges, opp. aporrhysa.
- episematic (ĕp'isēmăt'ik) a. [Gk. epi, upon; sema, sign.] Aiding in recognition; appl. coloration, markings.
- episeme (ĕp'isēm) n. [Gk. epi, upon; sema, sign.] A marking or colour aiding in recognition.
- episepalous (ĕp'isĕp'ālūs) a. [Gk. epi, upon; F. sépale, sepal.] Adnate to sepals.
- episkeletal (ĕp'iskel'ëtăl) a. [Gk. epi, upon; skeletos, hard.] Outside the endoskeleton.
- episperm (ĕp'ispĕrm) n. [Gk. epi, upon; sperma, seed.] The outer coat of seed; spermoderm; testa.

- episporangium (ēp'ispörăn'jiŭm) n. [Gk. epi, upon; sporos, seed; anggeion, vessel.] An indusium.
- epispore (ěp'ispōr) n. [Gk. *epi*, The outer upon; *sporos*, seed.] layer of a spore wall; perinium, q.v. epistasis (ĕpĭs'tăsĭs) n. [Gk. epi, upon; stasis, standing.]

inance of a gene over another, non-allelomorphic gene; epistasy, q.v.

epistasy (čpis'tăsi) n. [Gk. epi, upon; *stasis*, standing.] Greater degree of modification manifested by one of two related types in phylogenesis; masking of one hereditary character by another; epistasis, q.v.

epistatic (ĕp'ĭstăt'ĭk) a. [Gk. epistates, master.] Appl. the predominating of two characters whose genes are not allelomorphs; exhibiting or pert. the condition of

epistasis; cf. hypostatic.

episternalia (ěp'istěrnā'liă) n. plu. [Gk. epi, upon; sternon, breastbone.] Two small elements preformed in cartilage frequently intervening in development between clavicles and sternum, and ultimately fusing with sternum.

episternite (ĕp'istĕr'nīt) n. [Gk. epi, upon; sternon, breast-bone.] One of portions of an ovipositor formed from side portions of a somite.

- episternum (ěp'ĭstěr'nŭm) n. epi, upon; L. sternum, breast-bone.] The interclavicle; also applied to an anterior cartilaginous element of sternum; a lateral division of an arthropod somite, above sternum and in front of epimeron; anterior pleurite of subcoxa.
- epistome (ĕp'ĭstōm) n. [Gk. *epi*, upon; stoma, mouth.] A small lobe overhanging mouth in Polyzoa and containing a part of body cavity; the region between antenna and mouth in Crustacea; subcheliceral plate in certain ticks; that portion of insect head immediately behind labrum; portion of rostrum of certain Diptera; also epistoma.

[Gk. epiepistrophe (ĕpĭs'tröfĭ) n. strophe, moving about.] The position assumed by chloroplasts along outer and inner cell-walls when exposed to diffuse light.

epistropheus (ěp'istro'fëŭs) n. [Gk. epistrophe, turning.] The second

cervical or axis vertebra.

epithalamus (ĕp'ithăl'ămŭs) n. [Gk. epi, upon; thalamos, chamber. Part of thalamencephalon, comprising trigonum habenulae, pineal body, and posterior commissure.

epithalline (ěp'ĭthăl'īn) a. [Gk. epi, upon; thallos, branch.] Growing

upon the thallus.

- epitheca (ĕp'ĭthē'kă) n. [Gk. epi, upon; theke, cup.] An external layer surrounding lower part of theca in many corals; theca covering epicone in Dinoflagellata; older half of frustule in diatoms.
- epithecium (ěp'ithē'siŭm) n. [Gk. epi, upon; theke, cup.] The surface of spore-cases in lichens and fungi.
- epithelial (ĕp'ĭthē'lĭăl) a. [Gk. epi, upon; thele, nipple.] Pert. epithelium; epitheliomorph.
- epithelial bodies,—the parathyroids. epitheliofibrillae (ěpithē'liöfībril'ē) n. plu. [Gk. epi, upon; thele, nipple; L. fibrilla, small fibre.] Parallel or reticular fibrillae of columnar epithelium analogous to myofibrillae.
- epitheliomorph (ěpithē'liömôrf) a. [Gk. epi, upon; thele, nipple; morphe, form.] Resembling epithelium; epithelioid; appl. layer of cells, or elastica interna, which secretes notochordal sheath.
- epithelium (ĕp'īthē'lĭŭm) n. epi, upon; thele, nipple.] Any cellular tissue covering a free surface or lining a tube or cavity.
- [Gk. epi, epithem (ĕp'ĭthĕm) n. upon; tithenai, to put.] A plant tissue of specialised cells and intercellular spaces forming a hydathode; the secretory layer in nectaries: an excrescence on the beak of birds; also epithema and epitheme.

- epitokous (ĕpĭt'ökŭs) a. [Gk. epi, upon; tokos, birth.] Designating the heteronereid stage of certain polychaetes.
- epitrematic (ěp'îtrēmăt'îk) a. [Gk epi, upon; trema, pore.] Appl. upper lateral bar of branchial basket of lamprey.

epitrichial (ĕp'ĭtrīk'ĭāl) a. [Gk. epī, upon; thrix, hair.] Pert. or re-

sembling the epitrichium.

- epitrichium (ĕp'ĭtrĭk'iŭm) n. [Gk. epi, upon; thrix, hair.] An outer layer of foetal epidermis of many mammals, usually shed before birth.
- epitrochlea (ĕp'ĭtrŏk'lëă) n. [Gk.epi, upon; L. trochlea, Gk. trochilea, pulley.] Inner condyle at distal end of humerus.
- epitympanic (ĕp'itimpăn'ik) a. [Gk. epi, upon; L. tympanum, kettledrum.] Situated above tympanum.
- epivalve (ĕp'ivālv) n. [Gk. cpi, upon; L. valva, fold.] The apical part of envelope in certain Dinoflagellata; epicone.
- epixylous (čpĭzī'lūs, čpĭk'silūs) a. [Gk. epi, upon; xylon, wood.] Growing upon wood.
- epizoic (ĕp'ĭzō'ĭk) a. [Gk. epi, upon; zoon, animal.] Living on or attached to the body of an animal.
- epizoochore (ĕp'īzō'ökōr) n. [Gk. epi, upon; zoon, animal; chora, place.] Any spore, seed, or organism dispersed by being carried upon the body of an animal. Opp endozoochore.
- epizoon (ĕp'izō'ŏn) n. [Gk. epi, upon; zoon, animal.] An animal living on another; an external parasite; ectozoon.
- epizootic (ép'izōŏt'ik) a. [Gk. epi, upon; zoon, animal.] Common among animals. n. Disease affecting a large number of animals simultaneously, corresponding to epidemic in man.
- epizygal (ĕpĭz'igăl) n. [Gk. epi, upon; zygon, yoke.] The upper ossicle in a syzygial pair of brachials or columnars in crinoids.

- eponychium (ĕp'ŏnīk'iŭm) n. [Gk. epi, upon; onyx, nail.] The thin cuticular fold which overlaps lunula of nail; dorsal portion of a neonychium.
- epoophoron (ĕp'ōŏf'orōn) n. [Gk. epi, upon; oon, egg; pherein, to bear.] A rudimentary organ (homologous with epididymis), remains of Wolffian body of embryo, lying in mesosalpinx between ovary and uterine tube; organ of Rosenmüller.
- equal (ē'kwāl) a. [L. aequalis, equal.] Having the portions of the lamina equally developed on the two sides of midrib; appl. leaves.
- **equation division**, homeotypic or second division in meiosis.
- equatorial furrow,—division round equator of segmenting egg.
- equatorial plate,—group of chromosomes lying at equator of spindle during mitosis; locus of new cellwall after cell-division.
- equibiradiate (ē'kwībīrā'dĭāt) a. [1.. aeguus, equal; bis, twice; radius, ray.] With two equal rays.
- equicellular (ē'kwisĕl'ūlăr) a. [L. acquus, equal; cellula, cell.] Composed of equal cells.
- equifacial (ē'kwĭfā'shăl) a. |L. aequus, equal; facies, face.| Having equivalent surfaces or sides, as vertical leaves.
- equilateral (ē'kwilăt'erăl) a. [L. aequus, equal; latus, side.] Having the sides equal; appl. shells symmetrical about a transverse line drawn through umbo.
- equilenin (ěkwilē'nin) n. [L. equus, horse.] An oestrogenic hormone present in urine of the pregnant mare; C₁₈H₁₈O₂.
- equiline (ĕk'wilēn) n. [L. equus, horse.] An oestrogenic hormone, more physiologically active than equilenin, occurring in urine of the pregnant mare; C₁₈H₂₀O₂.
- equipotent (ēkwip'ötënt) a. [L. aequus, equal; potens, powerful.] Totipotent, q.v.
- equitant (ěk'wĭtănt) a. [L. equitare, to ride.] Overlapping saddlewise, as leaves in leaf-bud.

- equivalve (ē'kwĭvālv') a. [L. aequus, equal; valva, valve.] Having two halves of a shell alike in form and size.
- erect (ěrěkt') a. [L. erigere, to raise up.] Directed towards summit of ovary, appl. ovule; not decumbent. erectile (ěrěk'třl) a. [L. erigere, to

raise up.] Capable of being erected. erectile tissue,—a tissue capable of being made rigid by distention of

blood-vessels within it.

erection (ĕrĕk'shūn) n. [I. erigere, to raise up.] The state of a part which has become swollen and distended through accumulation of blood in erectile tissue.

erector (ĕrĕk'tŏr) n. [L. erigere, to raise up.] A muscle which raises

up an organ or part.

ereidesm (ĕrei'dezm) n. [Gk.ereidein, to support; desma, bond.] An epithelial intracellular fibre.

- eremobic (ĕrēmo'bik) a. [Gk. eremos, solitude; bios, life.] Growing or living in isolation; having a solitary existence.
- eremochaetous (ĕr'ēmökē'tūs) a. [Gk. eremos, lonely; chaeta, bristle.] Having no regularly arranged system of bristles; appl. flies.

eremophyte (ĕr'ēmöfīt') n. [Gk. eremos, solitude; phyton, plant.] A desert plant.

erepsin (erep'sin) n. [L. eripere, to set free.] A proteolytic enzyme of intestinal juice and body tissues.

- ergaloid (ĕr'găloid) a. [Gk. ergon, work; eidos, form.] Having the adults sexually capable though wingless.
- ergastic (ĕrgăs'tīk) a. [Gk. ergastikos, fit for working.] Pert. metaplasm; appl. lifeless cell-inclusions, as fat, starch, etc.

ergastoplasm (ĕrgās'töplāzm) n. [Gk. ergazesthai, to work; plasma, mould.] Archoplasm; kinoplasm.

- ergastoplasmic (ĕrgăs'töplăz'mik) a. [Gk. ergazesthai, to work; plasma, mould.] Appl. fibrillae of gland cells which may induce production of secretory granules.
- ergatandromorph (ergatan'drömôrf)

n. [Gk. ergates, worker; aner, male; morphe, form.] An ant or other social insect in which worker and male characters are blended.

ergatandrous (ĕrgātăn'drŭs) a. [Gk. ergates, worker; aner, man.] Hav-

ing worker-like males.

ergataner (ĕrgātā'nër) n. [Gk. ergates, worker; aner, male.] A male ant resembling a worker; an ergatoid or ergatomorphic male.

ergate (ĕr'gāt) n. [Gk. ergates, worker.] A worker-ant; ergates.

- ergatogyne (ĕrgā'töjīn'ē) n. [Gk. ergates, worker; gyne, female.] A female ant resembling a worker; an ergatoid or ergatomorphic female.
- ergatogynous (ĕrgātoj'inūs) a. [Gk. ergates, worker; gyne, woman.] Having worker-like females.
- ergatoid (ĕrgā'toid) a. [Gk. ergates, worker; eidos, form.] Resembling a worker, appl. ants; ergatomorphic.
- ergones (ĕr'gōnz) n. plu. [Gk. ergon, work.] Organic substances of which small amounts suffice for activation or regulation of a physiological process, as enzymes, hormones, and vitamins; also ergines.
- ergonomy (ĕr'gŏn'ömi) n. [Gk. ergon, work; nomos, law.] The differentiation of functions; physiological differentiation associated with morphological specialisation.

ergosterol,—a sterol occurring in plants and animals, with photochemical reaction products leading to formation of its isomer vitamin D₂; C₂₈H₄₄O.

- ergot (ĕrgot) n. [O.F. argot, spur.]
 A small bare patch found on limbs of horse-tribe, representing last remnant of naked palm of hand and sole of foot; condition of ovary of grasses produced by a fungus; rye smut; sclerotium of Claviceps, yielding several alkaloids, e.g. ergotoxine and ergometrine, which stimulate uterine muscle.
- erichthoidina (ĕrīk'thoid'īnă) n. [Gk. erechthein, to break; eidos, form.]
 Larval stage of Stomatopoda comparable with zoaea.

- erichthus (ĕrīk'thŭs) n. [Gk. erechthein, to break.] Larval stage of Stomatopoda comparable with pseudozoaea.
- eriocomous (ĕriŏk'ömŭs) a. [Gk. erion, wool; kome, hair.] Having woolly hair; fleece-haired.
- eriophyllous (ĕr'iöfil'ūs) a. [Gk. erion, wool; phyllon, leaf.] Having leaves with a cottony appearance.
- erose (ĕrōs') a. [L. erodere, to wear away.] Having margin irregularly notched; appl. leaf, bacterial colony.
- erosion (ĕrō'zhun) n. [L. erodere, to wear away.] Decay which usually starts at apex of many gastropod shells.
- erostrate (črŏs'trāt) a. [L. ex, without; rostrum, beak.] Having no beak; appl. antlers.
- ersaeome (ĕr'sēōm) n. [Gk. erse, young.] The free monogastric generation of Siphonophora.
- eruciform (ĕroo'sifôrm) a. [L. eruca, caterpillar; forma, shape.] Having the shape of, or resembling a caterpillar; appl. insect larvae; appl. spores of certain lichens.
- erumpent (ērūm'pent) a. [L. erumpere, to break out.] Breaking through suddenly; appl. fungal hyphae.
- erythrin (ĕrĭth'rĭn) n. [Gk. erythros, rcd.] A red colouring matter found in certain algae and lichens; C₂₀H₂₂O₁₀.
- erythrism (erith'rizm) n. [Gk. erythros, red.] Abnormal presence, or excessive amount, of red colouring matter, as in petals, feathers, hair, eggs; cf. rufinism.
- erythroblasts (ĕrīth'röblāsts) n. plu. [Gk. erythros, red; blastos, bud.] Nucleated cells, derived from mesoderm, which later contain haemoglobin and develop into red blood corpuscles.
- erythrocruorin (ĕrĭth'rökroo'örin) n. [Gk. erythros, red; L. cruor, blood.] Red iron-containing respiratory pigment in some invertebrates.
- erythrocyte (ĕrĭth'rösīt) n. [Gk.

- erythros, red; kytos, hollow.] A red blood corpuscle.
- formed by action of pyloric gland secretion (intrinsic factor) on extrinsic factor in food, and stored in liver, and necessary for maturation of red blood cells in bonemarrow; symbol: EMF.
- erythrocytolysis (ĕrĭth'rösītöl'isĭs) n. [Gk. erythros, red; kytos, cell; lysis, loosing.] Destruction of red blood corpuscles; haemolysis.
- erythron (erith'ron) n. [Gk. erythros, red; on, being.] The red cells in bone marrow and circulating blood, collectively.
- erythrophilous (ĕr'ĭthrŏf'ilŭs) a. [Gk. erythros, red; philein, to love.] Having special affinity for red stains; appl. structures in a cell or to a type of cells.
- erythrophore (ĕrith'röför) n. [Gk. erythros, red; pherein, to bear.] A reddish purple pigment bearing
- erythrophyll (ĕrĭth'röfil) n. [Gk. erythros, red; phyllon, leaf.] A red colouring matter of some leaves and of red algae.
- erythropoiesis (ĕrĭth'ropoi'ēsīs) n. [Gk. erythros, red; poiesis, making.] The production of red blood corpuscles.
- erythropsin (ĕrĭthrŏp'sĭn) n. [Gk. erythros, red; opsis, sight.] Red colouring matter in insect eyes; rhodopsin. q.v.
- erythrotin (ĕrith'rötin) n. [Gk. erythros, red.] Vitamin B₁₂, extracted from liver, anti-pernicious anaemia factor, and growth factor for certain micro-organisms.
- erythrozyme (ĕrĭth'rözīm) n. [Gk. erythros, red; zyme, leaven.] An enzyme capable of decomposing ruberythric acid, and acting upon glucosides.
- escape (eskāp') n. [M.E. escapen.]
 A plant originally cultivated, now found wild.
- escutcheon (ěskůch'ůn) n. [O.F. escuchon, shield.] Area on rump of many quadrupeds which is either

- variously coloured or has the hair specially arranged; mesoscutellum of certain insects; ligamental area of certain bivalves.
- eseptate (ēsĕp'tāt) a. [L. ex, without; septum, enclosure.] Not supplied with septa.
- esophagus,—oesophagus, q.v.
- esoteric (esoter'ik) a. [Gk. esoterikos, arising within.] Arising within the organism.
- espathate (ēspăth'āt) a. [L. ex, without; spatha, broad blade.] Having no spathe.
- esquamate (ēskwā'māt) a. [L. ex, without; squama, scale.] Having no scale.
- essential oils,—volatile oils, composed of various constituents and contained in plant organs, with characteristic odour.
- esth-,-see aesth-.
- estipulate (ēstip'ūlāt) a. [L. ex, without; stipula, stem.] Having no stipules.
- estr-,-see oestr-.
- estuarine (ĕs'tūārīn) a. [L. aestuarium, estuary.] Pert. or found in an estuary; appl. organisms.
- etaerio (ĕtē'riō) n. [Gk. etairia, association.] A fruit composed of achenes or drupels carried on an enlarged thalamus, or of follicles; eterio.
- etheogenesis (ē'thëöjēn'esis) n. [Gk. etheos, youth; genesis, descent.]
 Parthenogenesis producing males; development of a male gamete without fertilisation.
- Ethiopian (ē'thĭō'piān) a. [Gk. aithiops, burned-face.] Appl. or pert. a zoogeographical region including Africa south of the Sahara and southern Arabia, and divisible into African and Malagasy subregions.
- ethmohyostylic (ĕth'möhi'ostil'îk) a. [Gk. ethmos, sieve; T; stylos, pillar.] With mandibular suspension from ethmoid region and hyoid bar.
- ethmoid (eth'moid) a. [Gk. ethmos, sieve; eidos, shape.] Pert. bones which form a considerable part of walls of nasal cavity.

- ethmoidal (ĕthmoi'dăl) a. [Gk. ethmos, sieve; eidos, shape.] Pert. ethmoid bones or region.
- ethmoidal notch,—a quadrilateral space separating the two orbital parts of the frontal bone; incisura ethmoidalis.
- ethmolysian (ĕth'mölĭs'ĭăn) a. [Gk. ethmos, sieve; lyein, to loosen.] Pert. an apical system in which the madreporite extends backwards till it separates the two posterolateral genitals.
- ethmopalatine (ĕth'möpăl'ătin) a. [Gk. ethmos, sieve; L. palatus, palate.] Pert. ethmoid and palatine bones, or their region.
- ethmophract (ĕth'möfrākt) a. [Gk. ethmos, sieve; phrassein, to fence in.] Pert. a simple, compact, apical system with pores occurring only in right anterior corner.
- ethmoturbinals (ĕth'mötŭr'bĭnălz)

 n. plu. [Gk. ethmos, sieve; L.
 turbo, whorl.] Cartilages or bones
 in nasal cavity which are folded so
 as to increase olfactory area.
- ethmovomerine (ĕth'mövō'mërin) a. [Gk. ethmos, sieve; L. vomer, ploughshare.] Pert. ethmoid and vomer regions; appl. the cartilage which forms nasal septum in early embryo.
- ethnography (ethnog'rafi) n. [Gk. ethnos, nation; graphein, to write.]
 The description of the races of mankind.
- ethnology (ĕthnŏl'öjī) n. [Gk. ethnos, nation; logos, discourse.] Science dealing with the different races of mankind, their distribution, relationship, and activities.
- ethology (ēthŏl'öji) n. [Gk. ethos, custom; logos, discourse.] Bionomics; study of habits in relation to habitat; study of behaviour.
- ethomerous (ēthōm'ērūs) a. [Gk. ethos, custom; meros, part.] Having the normal number of parts or segments; with normal number of chromosomes.
- etiolation (ē'tiölā'shūn) n. [F. étioler, to blanch.] Blanched condition produced in plants reared in darkness, or by disease.

etiolin (ē'tīölǐn) n. [F. étioler, to blanch.] A yellowish pigment found in chloroplasts of plants grown in darkness; protochlorophyll.

etiology,—aetiology, q.v.

euapogamy (ū'apōg'āmī) n. [Gk. eu. well; apo, away; gamos, marriage.] Diploid apogamy, haploid apogamy being meiotic euapogamy.

euaster (ūăs'tër) n. [Gk. eu, good; aster, star.] An aster in which the rays meet at a common centre.

- eucarpic (ū'kârpĭk) a. [Gk. eu, well; karpos, fruit.] Appl. Phycomycetes having rhizoids, or haustoria. Opp. holocarpic.
- eucentric (ūsčn'trik) a. [Gk. eu, well; kentron, centre.] Pericentric, q.v.
- eucephalous (ūkěf'ălūs, -sěf-) a. [Gk. eu, good; kephale, head] With well-developed head; appl. certain insect larvae.
- euchromatic (ū'krōmăt'īk) a. [Gk. eu, well; chroma, colour.] Pert. euchromatin; appl. chromosome regions which never become heteropycnotic. Opp. heterochromatic.
- euchromatin (ūkrō'mătīn) n. [Gk. eu, well; chroma, colour.] Chromatin making up bulk of chromosome and including active genes.
- euchromosome (ūkrö'mösōm) n.
 [Gk. eu, well; chroma, colour; soma, body.] A typical chromosome, or autosome.
- eucoen (ūsēn) n. [Gk. eu, well; koinos, common.] Those members of a biocoenosis which are unable to live in a different environment. Opp. tychocoen.
- eucone (ūkōn) a. [Gk. eu, good; konos, cone.] Having crystalline cones fully developed in single elements of compound eye.

eudipleural (ū'dīploo'rāl) a. [Gk. eu, good; dis, double; pleuron, side.] Symmetrical about a median plane; bilaterally symmetrical.

eudoxome (ū'dōksōm) n. [Gk. eudoxos, glorious.] Monogastric freeswimming stage of a siphonophore without nectocalyx.

eugamic (ūgăm'īk) a. [Gk. eu, well; gamos, marriage.] Appl. mature period, opp. agamic or youthful, and aged or senescent.

eugenic (ūjěn'īk) a. [Gk. eu, well; genos, birth.] Pert. or tending towards racial improvement.

eugenics (ūjēn'īks) n. [Gk. eu. well; genos, birth.] The science dealing with the factors which tend to improve or impair stock.

euglenoid (ūglē'noid) a. [Gk. eu, well; glene, eyeball, puppet; eidos, form.] Pert. or like Euglena; appl. characteristic movement of Euglena.

euhaline (ūhăl'īn) a. [Gk. eu, well; halinos, saline.] Living only in saline inland waters; cf. euryhaline.

eumelanin (ū'mĕl'ānĭn) n. [Gk. eu, well; melas, black.] Black melanin; cf. phaeomelanin.

eumerism (ū'merizm) n. [Gk. eu, well; meros, part.] An aggregation of like parts.

- eumeristem (ū'mĕr'istĕm) n. [Gk. cu, well; meristos, divided.] Meristem composed of isodiametric thinwalled cells.
- eumerogenesis (ū'mer'öjen'ësīs) n. [Gk. eu, well; nucros, part; genesis, descent.] Segmentation in which the units are similar at least for a certain time.
- eumitosis (ūmštō'sis) n. [Gk. eu, well; mitos, thread.] Typical mitosis.
- eumitotic (ūmltot'īk) a. [Gk. eu, well; mitos, thread.] Anaschistic, q.v.; pert. cumitosis.
- euphotic (ūfō'tik) a. [Gk. eu, well; phos, light.] Well illuminated, appl. zone, surface waters to depth of about 80 metres; upper layer of photic zone; cf. dysphotic.

euphotometric (ū'fōtömĕt'rīk) a. [Gk. eu, well; phos, light; metron, measure.] Appl. leaves oriented to receive maximum diffuse light; cf. panphotometric.

euplastic (ūplăs'tik) a. [Gk. eu, well; plastos, moulded.] Readily organised, easily forming a tissue.

- euploid (ū'ploid) a. [Gk. eu, well; haploos, onefold; eidos, form.] Polyploid when total chromosome number is an exact multiple of the haploid number. Opp. aneuploid.
- eupotamic (ūpŏt'āmĭk) a. [Gk. eu, well; potamos, river.] Thriving both in streams and in their backwaters; appl. potamoplankton.

eupyrene (ū'pīrēn') a. [Gk. eu, well; pyren, fruit-stone.] Appl. sperms of normal type; cf. apyrene, oligopyrene.

eurybaric (ū'rībăr'īk) a. [Gk. eurys, wide; baros, weight.] Appl. animals adaptable to great differences

in altitude, opp. stenobaric. eurybathic (ū'rībāth'īk) a. [Gk.

eurybathic (ū'rībāth'īk) a. [Gk. eurys, wide; bathys, deep.] Having a large vertical range of distribution, opp. stenobathic.

eurychoric (ū'rīkō'rīk) a. [Gk. eurys, wide; choros, place.] Widely distributed, opp. stenochoric.

- eurycyst (ū'risist) n. [Gk. eurys, wide; kystis, bladder.] Large cell of middle vein in mosses; deuter cell; pointer cell.
- euryhaline (ū'rīhāl'īn) a. [Gk. eurys, wide; halinos, saline.] Appl. marine organisms adaptable to a wide range of salinity, opp. stenohaline.
- euryhygric (ū'rĭhī'grĭk) a. [Gk. eurys, wide; hygros, wet.] Appl. organisms adaptable to a wide range of atmospheric humidity.
- euryphagous (ūrif'āgūs) a. [Gk. eurys, wide; phagein, to eat.] Subsisting on a large variety of foods, opp. stenophagous; cf. omnivorous.
- eurypylous (ū'rīpī'lūs) a. [Gk. eurys, broad; pyle, gate.] Wide at the opening; appl. canal system of sponges in which the chambers open directly into excurrent canals by wide apopyles, and receive water from incurrent canals through prosopyles.
- eurysome (ū'risōm) a. [Gk. eurys, broad; soma, body.] Short and stout, opp. leptosome.
- eurythermic (ū'rĭthěr'mĭk) a. [Gk.

eurys, wide; therme, heat.] Appl. organisms adaptable to a wide range of temperature; eurythermous, opp. stenothermic.

eurytopic (ū'rĭtŏp'īk) a. [Gk. eurys, wide; topos, place.] Having a wide range of geographical distribution,

opp. stenotopic.

eusporangiate (ū'spörăn'jīāt) a. [Gk. eu, well; sporos, seed; anggeion, vessel.] Having sporogenous tissue derived from inner cell that follows periclinal division of superficial initial; cf. leptosporangiate.

Eustachian (ūstā'kiān) a. [B. Eustachio, Italian physician]. Pert. tube or canal connecting tympanic cavity with pharynx; pert. valve guarding orifice of inferior vena

cava in atrium of heart.

eustele (ūstē'lē) n. [Gk. eu, well; stele, pillar.] The arrangement of vascular tissue into collateral or bicollateral bundles with conjunctive tissue between, as in gymnosperms and dicotyledons.

eusternum (ūstěr'nům) n. [Gk. eu, well; sternon, breastplate.] A sternal sclerite of insects; antester-

nite, basisternum.

eustomatous (ūstŏm'ătŭs) a. [Gk. eu, well; stoma, mouth.] Having a distinct mouth-like opening.

- eutelegenesis (ū'tělĕjĕn'ësĭs) n. [Gk. eu, well; tele, afar; genesis, descent.] Improved breeding by artificial insemination.
- euthenics (ūthěn'īks) n. [Gk. euthenein, to thrive.] The science of betterment of human race on the side of intellect and morals; the study of environmental agencies contributing to racial improvement.

eutherian (ūthē'rĭăn) a. [Gk. eu, well; therion, small animal.] Appl. placental mammals with development uterine till full-time.

euthycomous (ūthĭk'ömūs) a. [Gk. euthys, straight; kome, hair.]

Straight-haired.

euthyneurous (ū'thinū'rŭs) a. [Gk. euthys, straight; neuron, nerve.] Having visceral loop of nervous system untwisted.

- eutrophic (ūtrŏf'îk) a. [Gk. eu, well; trophe, nourishment.] Providing, or pert., adequate nutrition. Opp. dystrophic.
- evagination (ē'vajinā'shun) n. [L. e, out; vagina, sheath.] The process of unsheathing, or product of this process; an outgrowth.

evanescent (ev'anes'ent) a. [L. evanescere, to vanish.] Disappearing early; appl. flowers which fade

quickly.

evection (ēvěk'shun) n. [L. e, out; vehere, to convey.] Displacement of parent cell at septum of a filament, causing dichotomous appearance, as in certain algae.

eviscerate (ẽvis'erāt) v. [L. ex, out; viscera, entrails.] To disembowel; to eject the viscera, as do holo-

thurians on capture.

- evocator (ĕv'ökātŏr) n. [L. evocator, caller forth.] The chemical stimulus furnished by an organiser, q.v.
- evolute (ĕv'ŏlūt) a. [L. evolvere, to unroll.] Turned back; unfolded.
- evolutility (čv'ŏlūtil'iti) n. [L. evolvere, to unroll.] Capability to evolve or change in structure; capacity to change in growth and form as a result of nutritional or other environmental factors.
- evolution (čv'olū'shūn) n. [L. evolvere, to unroll.] The gradual development of organisms from preexisting organisms since the dawn of life.

exalate (ěksā'lāt) a. [L. ex, without; ala, wing.] Not having wing-like

appendages; apterous.

exalbuminous (ek'salbu'minus) a. [L. ex, without; albumen, white of egg.] Without albumen; appl. seeds without endosperm or perisperm; exendospermous.

exannulate (ěksăn'ūlāt) a. [L. ex, without; annulus, ring.] Having a sporangium not furnished with an annulus; appl. certain ferns.

exarate (ěks arāt) a. [L. exaratus, ploughed up.] Appl. a pupa with free wings and legs.

exarch (ěks'ârk) n. [L. ex, without;

Gk. *arche*, beginning.] With protoxylem strands outside metaxylem, or in touch with pericycle.

exasperate (ĕgzăs'përāt) a. [L. exasperare, to roughen.] Furnished

with hard, stiff points.

excentric (ĕksĕn'trĭk) a. [L. ex, out of; centrum, centre.] One-sided; having the two portions of lamina unequally developed.

exciple (ĕk'sīpl) n. [L. excipula, receptacles.] The marginal wall, or outer covering, of apothecium in certain lichens; excipulum.

excitability,—capacity of a living cell, or tissue, to respond to an environmental change or stimulus.

- excitation (ëk'sĭtā'shŭn) n. [L. excitare, to rouse.] Act of producing or increasing stimulation; immediate response of protoplasm to a stimulus.
- excitatory cells,—motor cells in sympathetic nervous system.
- excitonutrient (ëksī'tönū'triënt) a. [L. excitare, to rouse; nutriens, feeding.] Causing or increasing nutrient activities.
- exconjugant (ěkskön'joogănt) n. [L. ex, out; conjugare, to yoke.] An organism which is leading an independent life after conjugation with another.
- excreta (ěkskrě'tă) n. plu. [L. excretum, separated.] Waste material eliminated from body or any tissue thereof; deleterious substances formed within a plant.
- excrete (ěkskrēt') v. [L. ex, out; cernere, to sift.] To eliminate waste material from body; to withdraw useless materials from the place of most active metabolism in plant.
- excretion (ĕkskrē'shūn) n. [L. ex, out; cernere, to sift.] Act of eliminating waste material, or the product of the elimination.
- excurrent (ěkskůr'ënt) a. [L. ex, out; currere, to run.] Pert. ducts, channels, or canals in which there is an outgoing flow; with undivided main stem; having midrib projecting beyond apex.

- excurved (ĕkskŭrvd') a. [L. ex, out; curvare, to curve.] Curved outwards from centre.
- excystation (ĕks'-sīstā'shun) n. [L.ex, out of; Gk. kystis, bladder.] Emergence from encysted condition.
- exendospermous (ěks'ěndöspěr'můs)

 a. [L. ex, without; Gk. endon, within; sperma, seed.] Exalbuminous, a.v.
- exflagellation (ĕksflăj'ēlā'shūn) n.
 [L. ex, out of; flagellum, whip.]
 Process of microgamete formation by microgametocyte in Haemosporidia.
- exfoliation (ĕksfō'liā'shūn) n. [L. ex, out; folium, leaf.] The shedding of leaves or scales from a bud.
- exhalant (ĕks'hā'lānt) a. [L. ex, out; halare, to breathe.] Capable of carrying from the interior outwards.
- exindusiate (ĕk'sĭndū'zĭāt) a. [L. ex, out; indusium, cover.] Having the sporangia uncovered or naked.

exine,—extine, q.v.

- exinguinal (ĕk'sing'gwinăl) a. [L. ex, out; inguen, groin.] Occurring outside the groin; pert. second joint of arachnid leg.
- exites (ëk'sīts) n. plu. [Gk. exo, without.] Offshoots on outer lateral border of axis of certain arthropod limbs.
- exocardiac (ĕk'sökâr'dĭāk) a. [Gk. exo, without; kardia, heart.] Situated outside the heart.
- exocarp (ĕk'sökârp) n. [Gk. exo, without; karpos, fruit.] Outer layer of the pericarp; epicarp.
- exoccipital (ĕk'sŏksip'ftăl) a. [L. ex, without; occiput, back of head.]

 Pert. a skull bone on each side of the foramen magnum.
- exochiton (ĕk'söki'tŏn) n. [Gk. exo, without; chiton, coat.] Outermost layer of oogonial wall, as in Fucales; exochite; cf. endochiton, mesochiton.
- exochorion (ĕk'sökō'riŏn) n. [Gk. exo, without; chorion, chorion.]

 Outer layer of membrane secreted by follicular cells surrounding the egg in ovary of insects.
- exocoel (ěk'sösēl) n. [Gk. exo, without; koilos, hollow.] The space

- between mesenteries of adjacent couples in certain Zoantharia; exocoelom, q.v.
- exocoelar (ĕk'sösē'lăr) a. [Gk. exo, without; koilos, hollow.] Pert. parietal wall of coelom.
- exocoelic (ĕk'sösē'lĭk) a. [Gk. exo, without; koilos, hollow.] In Zoantharia, pert. space between adjacent couples of mesenteries; appl. radial areas on disc; appl. outermost cycle of tentacles.
- exocoelom (ěk'sösē'lŏm) n. [Gk. exo, without; koilos, hollow.] Extraembryonic body cavity of embryo.
- exocone (ĕk'sökōn) a. [Gk. exo, without; konos, cone.] Appl. insect compound eye with cones of cuticular origin.
- exocrine (ĕk'sökrı̈n) a. [Gk. exo, without; krinein, to separate.] Appl. glands whose secretion is drained by ducts; cf. endocrine, apocrine.
- exocuticula (čk'sökūtīk'ūlā) n. [Gk. exo, without; L. dim. of cutis, skin.] Middle layer of insect cuticle, between endocuticula and epicuticula; outer layer of integument in spiders.
- exoderm (ĕk'södĕrm) n. [Gk. exo, without; derma, skin.] The dermal layer of sponges.
- exodermis (ëk'södër'mis) n. [Gk. exo, without; L. dermis, skin.] A specialised layer below the piliferous layer; ectoderm, q.v.
- exoenzyme (ěk'söĕn'zīm) n. [Gk. exo, outside; zyme, leaven.] Any extracellular enzyme.
- exogamete (ĕk'sögămēt') n. [Gk. exo, without; gametes, mate.] A reproductive cell which fuses with one derived from another source.
- exogamy (ĕksög'āmi) n. [Gk. exo, without; gamos, marriage.] Conjugation or fusion of isogametes with others of a different brood; outbreeding.
- exogastric (ĕk'sögăs'trik) a. [Gk. exo, without; gaster, stomach.] Having the shell coiled towards dorsal surface of body.

- exogastrula (ĕk'sögăs'troolā) n. [Gk exo, without; gaster, stomach.]
 An hour-glass shaped sea-urchin larva induced experimentally.
- exogenous (ĕksŏj'ēnūs) a. [Gk. exo, outside; -genes, producing.] Originating outside the organism; developed from superficial tissue, the superficial meristem; growing from parts which were previously ossified; appl. metabolism concerned with effector activities and temperature.
- exognathion (ěk'sögnåth'iŏn) n. [Gk. exo, without; gnathos, jaw.] The maxillary portion of upper jaw; the maxilla with exception of endognathion and mesognathion.
- exo-intine (ĕk'sŏĭn'tĭn) n. [Gk. exo, without; L. intus, within.] Middle layer of a spore-covering, between extine and intine.
- exonephric (ĕk'sönĕf'rĭk) a. [Gk. exo, without; nephros, kidney.] With nephridia opening to exterior; opp. enteronephric, appl. Oligochaeta.
- exoperidium (ĕk'söpērĭd'ĭŭm) n. [Gk. exo, without; peridion, a small wallet.] The outer layer of spore case in certain fungi; epiperidium.
- exophylaxis (ék'sofilāk'sis) n. [Gk. exo, without; phylax, guard.] Protection afforded against pathogenic organisms by skin secretions.
- exoplasm (čk'söplăzm) n. [Gk. exo, without; plasma, mould.] Ectoplasm, q.v.
- exopodite (ěks'öpŏdīt) n. [Gk. exo, without; pous, foot.] The outer branch of a typical biramous crustacean limb.
- exoscopic (ĕk'sösköp'ĭk) a. [Gk. exo, without; skopein, to look.] With apex emerging through archegonium, appl. embryo. Opp. endoscopic.
- exoskeleton (ěk'söskěl'ëtön) n. [Gk. exo, without; skeletos, hard.] A hard supporting structure secreted by ectoderm or by skin.
- exosmosis (ěk'sŏsmō'sĭs) n. [Gk. exo, without; osmos, impulse.] The passing out through a membrane of a gas or fluid. Opp. endosmosis. exospore (ěk'söspōr), exosporium

- (ĕk'söspō'rīūm) n. [Gk. exo, without; sporos, seed.] Outer coating of sporangial wall; a conidium.
- exostome (ěk'söstōm) n. [Gk. exo, without; stoma, mouth.] Opening or foramen in outer wall of ovule.
- exostosis (ĕk'sŏstō'sis) n. [Gk. exo, without; osteon, bone.] Formation of knots on surface of wood; formation of knob-like outgrowths of bone at a damaged portion, or of dental tissue in a similar way.
- exotheca (ěk'sothē'kă) n. [Gk. exo, without; theke, box.] The extracapsular tissue of a coral.
- exothecal (ěk'söthē'kăl) a. [Gk. exo, without; theke, box.] Pert. tissue outside the theca of a coral.
- exothecate (čk'sothē'kāt) a. [Gk. exo, without; theke, box.] Having an exotheca.
- exothecium (ěk'söthē'sĭum) n. [Gk. exo, without; theke, case.] The outer specialised dehiscing cell layer of the gymnosperm sporangum; cf. endothecium.
- exotic (ĕgzŏt'ik) a. [Gk. exotikos, foreign.] Introduced or nonendemic. n. A foreign plant or animal not acclimatised.
- exotospore (čksō'tospōr) n. [Gk. exotos, outward; sporos, seed.]
 A sporozoite.
- exotoxin (ĕk'sotŏk'sīn) n. [Gk. exo, without; toxikon, poison.] A soluble toxin excreted by bacteria; opp. endotoxin.
- exotropism (čksŏťröpĭzm) n. [Gk. exo, without; trope, turn.] Curvature away from axis, exhibited by a laterally geotropic organ.
- expantin, presumable neurohumor inducing expansion of chromatophores in crustaceans. Opp. contractin.
- expiration (čk'spĭrā'shŭn) n. [L. ex, out; spirare, to breathe.] The act of emitting air from lungs; emission of carbon dioxide by plants and animals.
- explantation (ĕk'splāntā'shūn) n. [L. ex, out of; plantare, to plant.] Tissue culture away from organism of its origin.

- explosive,—appl. flowers in which pollen is suddenly discharged on decompression of stamens by alighting insect, as of Cytisus and Ulex; appl. fruits with sudden dehiscence, seeds being discharged to some distance; appl. evolution, rapid formation of numerous types; tachytypogenesis; appl. speciation, rapid formation of species from a single species in one locality.
- **expressivity,**—the degree to which a gene produces an effect.
- exsculptate (ěks'skŭlp'tāt) a. [L. ex, out; sculpere, to carve.] Having the surface marked with more or less regularly arranged raised lines with grooves between.
- exscutellate (ĕkskū'tēlāt) a. [L. ex, without; scutellum, small shield.] Having no scutellum; appl. insects.
- exserted (ĕksĕr'tëd) a. [L. exserere, to stretch out.] Protruding beyond some including organ or part; appl. stamens which project beyond corolla.
- exsertile (ěksěr'til) a. [L. exserere, to stretch out.] Capable of extrusion.
- exstipulate (ĕkstip'ūlāt) a. [L. ex, without; stipula, stem.] Without stipules.
- exstrophy (eks'ströfi) n. [Gk. exo, outwards; strophe, turning.] Eversion, as normal or anomalous projection of luteal tissue to exterior of ovary.
- exsuffiation (ěks'sūflā'shūn) n. [L. ex, out; suffiare, to blow.] Forced expiration from lungs.
- extend (ĕkstěnd') v. [L. ex, out; tendere, to stretch.] To straighten out, opp. to flex or bend any organ.
- extensor (ěkstěn'sŏr) n. [L. ex, out; tendere, to stretch.] Any muscle which extends a limb or part. Opp. flexor.
- exterior (ekste'rior) a. [L. externus, on outside.] Situated on side away from axis or definitive plane.
- external (ěkstěr'năl) a. [L. externus, outside.] Outside or near the outside; away from the mesial plane.

externum (ĕkstĕr'nŭm) n. [L. externus, outward.] Outer region or cortex of a mitochondrium or of Golgi apparatus.

exteroceptor (ĕk'stërösĕp'tŏr) n. [L. exter, outside; capere, to take.]
A receptor which receives stimuli from outside the body; a contact receptor, or distance receptor.

extine (ĕk'stǐn) n. [L. exter, outside.]
Outer coat of spore or pollen grain;
exosporium. Opp. intine.

extra-axillary (ĕk'strâ-āks'îlărī) a. [L. extra, beyond; axilla, armpit.] Arising above axil of leaf, said of branches which develop from upper bud when there are more than one in connection with axil.

extrabranchial (ěk'străbrăng'kĭăl) a. [L. extra, beyond; Gk. brangchia, gills.] Arising outside the branchial arches.

- extracapsular (ĕk'străkāp'sūlăr) a. [L. extra, outside; capsula, small box.] Arising or situated outside a capsule; appl. ligaments, etc., in connection with a joint; appl. protoplasm lying outside the central capsule in some protozoa; appl. dendrites.
- extracellular (ĕk'străsĕl'ūlăr) a. [L. extra, outside; cellula, little cell.]
 Occurring outside the cell; diffused out of the cell.
- extracolumella (ěk'străkŏl'ūměl'ă) n. [L. extra, beyond; columella, small column.] Distal element of auditory skeletal structure; also hyostapes.
- extraembryonic (ĕk'strâĕm'brĭŏn'îk)
 a. [L. extra, outside; Gk. embryon, foetus.] Situated outside the embryo proper, as portion of blastoderm.
- extraenteric (ĕk'strāĕntĕr'īk) a. [L. extra, outside; Gk. enteron, gut.]
 Outside the alimentary tract.
- extrafloral (ĕk'strāflō'rāl) a. [L. extra, outside; flos, flower.] Situated outside the flower; appl. nectaries.
- extrafoveal (ěk'străfô'věäl) a. [L. extra, beyond; fovea, depression.]

 Pert. macula lutea surrounding fovea centralis; appl. rod vision.

 Opp. foveal.

- extranuclear (čk'strănū'kleăr) a. [I.. extra, outside; nucleus, kernel] Pert. structures or forces acting outside the nucleus; situated outside the nucleus.
- extraocular (ěk'străŏk'ūlăr) a. [L. extra, outside; oculus, eye.] Exterior to the eye; appl. antennae of insects.

extraperitoneal,—subperitoneal.

- extraspicular (ěk'stráspik'ūlăr) a. [L. extra, outside; spicula, small spike.] With spicules having one end embedded in spongin and the other end free.
- extrastapedial (ĕk'străstăpē'dĭăl) a. [L. extra, beyond; stapes, stirrup.] Extending beyond the stapedio-columellar junction.
- extrastelar (ěk'străstē'lăr)a. [L. extra, outside; Gk. stele, column.] Pert. ground tissue outside vascular tissue.
- extravaginal (čk'stravaj'inal) a. [L. extra, outside; vagina, sheath.] Forcing a way through the sheath, as shoots of many plants.
- extravasate (ĕkstrāv'āsāt) v. [L. extra, outside; vas, vessel.] To force its way from the proper channel into the surrounding tissue, said of blood, etc.
- extraventricular(ěk'străvěntrĭk'ūlăr)
 a. [L. extra, beyond; ventriculus, belly.] Situated or arising beyond the ventricle.
- extraxylary (ëk'străzī'lărĭ) a. [L. extra, outside; Gk. xylon, wood.]
 On the outside of the xylem; appl. fibres.
- extremity (ĕkstrĕm'îtĭ) n. [L. extremitas, limit.] The limb, or distal portion of a limb; distal end of any limb-like structure.
- extrinsic (ekstrin'sik) a. [L. extrinsecus, on outside.] Pert. muscles not wholly within the part; appl. brightness due to objective light intensity. Opp. intrinsic.
- extrorse (ěkstrôrs') a. [L. extrorsus, outwardly.] Turned away from axis; appl. dehiscence of anthers.
- exudation (ěk'sūdā'shūn) n. [L. exudare, to sweat.] Any discharge

- through an incision or pore, e.g. gums, resins, moisture, etc.
- exumbral (čksům'brāl) a. [L. ex, out; umbra, shade.] Pert. rounded upper surface of a jelly-fish.
- exumbrella (ěks'úmbrěl'ă) n. [L. ex, out; umbra, shade.] Upper, convex surface of jelly-fish.
- exuviae (čksū'vĭē) n. plu. [L. exuere, to strip off.] Cast-off skins, shells, etc., of animals.
- exuvial (ĕksū'vĭāl) a. [L. exuere, to strip off.] Ecdysial; appl. insect glands whose secretion facilitates ecdysis.
- eye (i) n. [A.S. éage.] The organ of sight or vision; a pigment spot in various animals and in lower plants; the bud of a tuber.
- eye-spots,—certain pigment spots in many lower plants and animals, and also in some vertebrates, which are supposed to have a visual function; ocelli.
- eye-teeth, -upper canine teeth.
- F₁,—denotes first filial generation, or hybrids arising from a first cross, successive generations arising from this one being denoted by F₂, F₃, etc. P₁ denotes parents of F₁ generation, P₂ the grandparents, etc.
- fabella (făběl'ă) n. [L. fabella, small bean.] A small fibro-cartilage ossified in the gastrocnemius.
- fabiform (fāb'ífôrm) a. [L. faba, bean; forma, shape.] Bean-shaped.
- Fabrician [J. C. Fabricius, Danish entomologist]. Appl. a classification of the Arthropoda based on the anatomy of the mouth-
- facet (făs'ĕt, făsĕt') n. [L. facies, face.] A smooth, flat, or rounded surface for articulation; an ocellus; corneal portion of insect eye.

facial (fā'shăl) a. [L. facies, face.]
Pert. face; appl. artery, nerve, bone, vein, etc.

faciation (fāsīā'shun) n. [L. facies, face.] Formation or character of facies; a grouping of dominant species within an association; geographical differences in abundance or proportion of dominant species in a community; cf. lociation.

facies (fā'shīez) n. [L. facies, face.] The face; a surface, in anatomy; the general aspect of a plant; aspect, as superior and inferior; a particular modification of a biotope; a grouping of dominant plants in the course of a successional series; one of different types of deposit in a geological series or system; the palaeontological and lithological character of a deposit.

facilitation (făsilită'shun) n. [L. facilitas, easiness.] Diminution of resistance to a stimulus subsequent to previous stimulusion, as of

nerves; Ger. Bahnung.

faciolingual (fā'sīölīng'gwăl) a. [L. facies, face; lingua, tongue.]

Pert. or affecting face and tongue.

factor (făk'tör) n. [L. facere, to make.] Any agent (biotic, climatic, nutritional, etc.) contributing to a result; a Mendelian factor or gene; a determinant.

factorial (făktō'riăl) a. [L. facere, to make.] Pert. genetic factors or genes.

facultative (făk'ūltā'tīv) a. [L. facultas, faculty.] Having the power of living under different conditions; conditional; appl. organisms which may be normally self-dependent, but which are adaptable to a parasitic or semiparasitic mode of life; appl. aerobes, anaerobes; appl. parthenogenesis, symbionts, saprophytes, gametes, etc. Opp. obligate.

faeces (fē'sēz) n. plu. [L. faeces, dregs.]

Excrement from alimentary canal.

falcate (făl'kāt) a. [L. falx, sickle.] Sickle-shaped; hooked.

falces (făl'sēz) n. plu. [L. falces, sickles.] Chelicerae, of arachnids.

falciform (făl'sĭfôrm) a. [L. falx, sickle; forma, shape.] Sickle-shaped or scythe-shaped; appl.

ligament, a dorso-ventral fold of peritoneum, attached to under surface of diaphragm and anterior and upper surfaces of liver; appl. process, processus falciformis, a fold of choroid penetrating retina near optic disc and ending at back of lens, functioning in accommodation in teleosts; appl. body, a sporozoite.

falcula (făl'kūlă) n. [L. falcula, little hook.] A curved scythe-like claw.

falculate (făl'kūlāt) a. [L. falcula, little hook.] Curved, and sharp at the point.

Fallopian tube [G. Fallopio, Italian anatomist]. Uterine tube, upper portion of oviduct in mammals; anterior portion of the Müllerian duct.

false or asternal ribs,—those ribs whose cartilaginous ventral ends do not join the sternum directly.

false vocal cords,—ventricular folds of larynx, two folds of mucous membrane, each covering a ligament, anterior to true vocal cords.

falx (fälks) n. [L. falx, sickle.] A sickle-shaped fold of the dura mater; inguinal aponeurosis of transverse and internal oblique muscles of abdomen.

family (făm'îli) n. [L. familia, household.] Term used in classification, signifying a group of related genera, families being grouped into orders.

fan (făn) n. [A.S. fann, fan.] A bird's tail feathers; a flabellum, q.v.; a rhipidium, q.v.; vannus, q.v.

fang (fáng) n. [A.S. fang, grip.] A long-pointed tooth, especially the poison tooth of snakes; the root of a tooth.

faradisation (făr'ădĭzā'shūn) n. [M. Faraday, English physicist]. Method of stimulation inducing partial or complete tetanus.

farctate (fârk'tāt) a. [L. farctus, stuffed.] Filled, not hollow.

farina (fărē'nă, fărī'nă) n. [L. farina, flour.] The pollen of plants; the fine mealy-like powder found on some insects.

- farinaceous (făr'inā'shus) a. [L. farina, flour.] Containing flour; starchy; farinose.
- farinose (făr'inōs) a. [L. farina, flour.] Producing, or covered with, fine powder or dust.
- fascia (făs'ĭă, făsh'ĭă) n. [L. fascia, band.] An ensheathing band of connective tissue.
- fascial (făs'ĭăl, făsh'ĭăl) a. [L. fascia, bundle.] *Pert*. a fascia, ensheathing and binding.
- fasciated (fas'īa'ted, fash'īa'ted) a. [L. fascia, bundle.] Banded; arranged in fascicles; appl. stems or branches malformed and flattened.
- fasciation (făshĭā'shŭn) n. [L. fascia, bundle.] The formation of fascicles; coalescent development of branches of a shoot-system, as in cauliflower.
- fascicle (făs'īkl) n. [L. fasciculus, small bundle.] A small bundle or tuft, as of fibres, or of leaves.
- fascicular (făsīk'ūlār) a. [L. fasciculus, small bundle.] Pert. a fascicle; arranged in bundles or tufts; appl. cambium, tissue.
- fasciculus (făsĭk'ūlūs) n. [L. fasciculus, small bundle.] A fascicle; a group, bundle, or tract of nerve fibres, as of medulla spinalis.
- fasciola (făsī'ölă) n. [L. fasciola, small bandage.] A narrow colour band; a delicate lamina continuous with supracallosal gyrus.
- fasciole (făs'iōl) n. [L. fasciola, small bandage.] Ciliated band on certain echinoids for sweeping water over surrounding parts.
- fastigiate (făstij'îāt) a. [L. fastigare, to slope up.] With branches close to stem and erect, opp. patent; in pyramidal or conical form.
- fastigium (făstij'iŭm) n. [L. fastigium, gable.] Angular top of roof of fourth ventricle, formed by contact of anterior and posterior medullary vela of cerebellum.
- fat (făt) n. [A.S. faet, fat.] Adipose tissue; any part of animal tissue which has its cells filled with a greasy or oily reserve material.

- fat-body,—one of the vascularised tissue structures filled with fat globules and associated with gonads in Amphibia; one of the subcutaneous organs along ventral sides and enlarged during breeding season in Lacertilia; tissue of indeterminate form distributed throughout body of insects and functioning as nutritive reserve; corpus adiposum; epiploon.
- fat soluble,—appl. vitamins A, D, E, and K.
- fatigue (fătēg') n. [L. fatigare, to weary.] Effect produced by long stimulation on cells of an organ.
- fauces (fôs'ēz) n. plu. [L. fauces, throat.] Upper or anterior part of throat between palate and pharynx; mouth of a spirally coiled shell; throat of a corolla.
- fauna (fôn'ă) n. [L. faunus, god of woods.] All the animals peculiar to a country, area, or period.
- faunal region,—an area characterised by a special group or groups of animals.
- faunula (fôn'ūlā) n. [Dim. of fauna.]
 Animal population of a small unit area, as of intestine, bark, etc.
- favella (făvěl'à) n. [L. favus, honeycomb.] A conceptacle of certain red algae.
- faveolate (făvē'ölāt) a. [L. faveolus, dim. of favus, honey-comb.] Honeycombed or alveolate.
- faveolus (făvē'olŭs) n. [L. faveolus, small honey-comb.] A small depression or pit; alveola.
- favoid (fav'oid) a. [L. favus, honeycomb; Gk. eidos, form.] Resembling a honey-comb.
- favose (făvos') a. [L. favus, honeycomb.] Honeycombed; alveolate.
- feather-veined,—appl. leaf in which veins run out from mid-rib in regular series at an acute angle; pinnately veined.
- feces,—facces.
- Fechner's Law [G. T. Fechner, German psychophysicist]. The tendency of intensity of sensation to vary as the logarithm of the stimulus.

fecundity (fekŭn'dĭtĭ) n. [L. fecundus, fertile.] Power of a species to multiply rapidly; capacity to form reproductive elements.

female (fē'māl) n. [L. femina, women.] A pistillate flower; an egg-producing or young-producing animal—symbol ♀.

female pronucleus,—the nucleus left in the ovum after maturation.

femoral (fem'öral) a. [L. femur, thigh.] Pert. thigh; appl. artery, vein, nerve, etc.; crural.

femur (fē'mūr) n. [L. femur, thigh.]
The thigh-bone, proximal bone of hind limb in vertebrates; third joint in insect and spider leg counting from proximal end.

fenchone (fĕn'chōn) n. [Ger. Fenchel, fennel.] A ketone, the essential oil in oil of fennel; C₁₀H₁₆O.

fenestra (fënës'tră) n. [L. fenestra, window.] An opening in a bone, or between two bones, or in a plant membrane; a pit on head of cockroach; a transparent spot on wings of insects.

fenestrate (fëněs'trāt) a. [L. fenestra, window.] Having small perforations or transparent spots, appl. insect wings; having numerous perforations, appl. leaves, dissepiments.

fenestrated membrane,—a close network of yellow elastic fibres resembling a membrane with perforations, as in inner tunic of arteries.

fenestrule (fěněs'trool) n. [Dim. of L. fenestra, window.] Small opening between branches of a polyzoan colony.

feral (fē'rāl) a. [L. fera, wild animal.]
Wild, or escaped from cultivation
or domestication and reverted to
wild state.

ferment (fer'ment) n. [L. fermentum, ferment.] An organised substance, capable of producing fermentation; an enzyme.

fermentation (fer'menta'shun) n.
[L. fermentum, ferment.] A transformation occurring in organic substance, usually of a carbo-

hydrate, caused by action of a ferment; zymosis.

ferruginous (fĕroo'jinŭs) a. [L. ferruginus, rusty.] Having the appearance of iron rust.

fertile (fĕr'tīl) a. [L. fertilis, fertile.]
Capable of producing living offspring; of eggs or seeds, capable
of developing.

fertilisation (fer'tillza'shun) n. [L. fertilis, fertile.] The union of male and female pronuclei; pollination

fertilisation - tube,—process of an antheridium, penetrating oogonial wall, for passage of male gamete in certain fungi.

fertility vitamin,—α tocopherol or vitamin E; anti-sterility vitamin.

fertilizin (fer'tili'zin) n. [L. fertilis, fertile.] A soluble colloidal substance produced by certain eggs and causing sperm agglutination, also inducing cleavage.

festoon (festoon') n. [F. feston, garland.] The margin, with rectangular divisions, of integument in ticks.

fetlock (fěťlòk) n. [Icel. fet, step; lokka, tuft of hair.] The tuft of hair behind a horse's pastern joint; the pastern joint itself.

fetus,-see foetus.

fibre (fi'ber) n. [L. fibra, band.] A strand of nerve, muscle, connective, or bast tissue; elongated plant-cell for mechanical strength; fiber.

fibre tracheids,—fibres of a nature intermediate between that of libriform fibres and of tracheids.

fibril (fi'bril) n. [L. fibrilla, small fibre.] A small thread-like structure or fibre; a component part of a fibre; a root-hair; a slender filliform outgrowth on some lichens.

fibrillae (fibril'ē) n. plu. [L. fibrilla, small fibre.] Thread-like branches of roots; minute elastic fibres secreted within spongin cells; minute muscle-like threads found in various infusorians; fibrils.

fibrillate (fi'brilat) a. [L. fibrilla, small fibre.] Having fibrillae or

hair-like structures.

- fibrillose (fī'brīlös, fībrīl'ös) a. [I.. fibrilla, small fibre.] Furnished with fibrils; appl. mycelia of certain fungi.
- fibrin (fi'brin) n. [L. fibra, band.]
 An insoluble protein found in blood
 after coagulation, readily digested
 in gastric juice.
- fibrinogen (fibrin'öjen) n. [L. fibra, band; Gk. -genes, producing.] A soluble protein of blood, which, by activity of thrombin, yields fibrin and produces coagulation.
- fibroblast (fi'bröbläst) n. [L. fibra, band; Gk. blastos, bud.] A connective tissue cell; fibrocyte, desmocyte.
- fibro-cartilage (fi'brö-kâr'tilëj) n. [L. fibra, band; cartilago, gristle.] A kind of cartilage whose matrix is mainly composed of fibres similar to connective tissue fibres, found at articulations, cavity margins, and osseous grooves.
- fibrous (fī'brŭs) a. [L. fibra, band.] Composed of fibres; appl. tissue, roots, mycelium, etc.
- fibula (fib'ūlă) n. [L. fibula, buckle.]
 Outer and smaller shin bone.
- fibulare (fib'ūlā'rē) n. [L. fibula, buckle.] The outer element of proximal row of tarsus.
- field,—a dynamic system in which all the parts are interrelated and in equilibrium, so that a change in any part affects the whole.
- filament (fil'ămënt) n. [L. filum, thread.] A thread-like structure; the stalk of anther; a hypha, q.v.; stalk of a down-feather; a cryptoptile; slender apical end of egg-tube of insect ovary.
- filamentous (filamen'tus) a. [L. filum, thread.] Thread-like; having filaments; appl. form or margin of certain bacterial colonies.
- filial generation,—F₁, etc., q.v. filial regression,—tendency of offspring of outstanding parentage to revert to average for species.
- filicauline (fil'Ikôl'In) a. [L. filum, thread; caulis, stalk.] With a thread-like stem.
- filiciform (fil'isifôrm), filicoid (fil'-

- Ikoid) a. [L. filix, fern; forma, shape; Gk. eidos, form.] Shaped like the frond of a fern; fern-like.
- filiform (fil'ifôrm) a. [L. filum, thread; forma, shape.] Thread-like.
- filiform papillae,—papillae on the tongue, ending in numerous minute slender processes.
- filigerous (filij'ërŭs) a. [L. filum, thread; gerere, to carry.] With thread-like outgrowths or flagella.
- Filippi's glands,—paired glands with ducts conveying viscid secretion into silk ducts.
- fillet (fil'et) n. [L. filum, thread.]
 Band of white matter in midbrain and medulla oblongata;
 lemniscus.
- filoplume (fil'oploom) n. [L. filum, thread; pluma, feather.] A delicate hair-like feather with long axis and a few free barbs at apex.
- filopodia (fil'öpō'dĭā) n. plu. [L. filum, thread; Gk. pous, foot.]
 Protozoan thread-like pseudopodia.
- filose (fi'los) a. [L. filum, thread.] Slender; thread-like; appl. pseudopodia of protozoa.
- filter-passers,—organisms capable of passing through a filter which arrests bacteria; microhenads; viruses.
- filtration (filtra'shun) n. [F. filtrer, to strain.] Appl. iridial angle of cornea; straining, as of lymph through capillary walls.
- filum terminale, the terminal thread, a slender grey filament, of the spinal cord.
- fimbria (fim'bria) n. [L. fimbria, fringe.] Any fringe-like structure; a posterior prolongation of fornix to hippocampus; one of delicate processes fringing the mouth of tube or duct, as of oviduct, or of siphon of molluscs.
- fimbriated (fim'briā'tēd) a. [L. fimbria, fringe.] Fringed at margin, as petals, tubes, ducts, antennae.
- fin (fin) n. [A.S. finn, fin.] A fold of skin with fin-rays and skeletal supports, in most fishes.

- finials (fin'iălz) n. plu. [L. finis, end.]
 The ossicles of the distal rami of crinoids, which do not branch again.
- fin-rays,-horny supports of fins.
- fissilingual (fis'illing'gwăl) a. [L. fissus, cleft; lingua, tongue.] With bifid tongue.
- fission (fish'ŭn) n. [L. fissus, cleft.]
 Cleavage of cells; division of a unicellular organism into two or more parts.
- fissiparous (fisip'arus) a. [L. fissus, cleft; parere, to beget.] Reproducing by fission.
- fissiped (fis'iped) n. [L. fissus, cleft; pes, foot.] With cleft feet, that is, with digits of feet separated.
- fissirostral (fis'īrŏs'trăl) a. [L. fissus, cleft; rostrum, beak.] With deeply-cleft beak.
- fissure (fish-ūr) n. [L. fissus, cleft.]

 A cleft, deep groove, or furrow dividing an organ into lobes, or subdividing and separating certain areas of the lobes; sulcus.
- fistula (fis'tūlă) a. [L. fistula, pipe.] Pathological or artificial pipe-like opening; trachea, water-conducting vessel.
- fistular (fis'tūlăr) a. [L. fistula, pipe.] Like a fistula; pipe-like; hollow, as stems of Umbelliferae.
- fix (fiks) v. [L. fixus, fixed.] To kill, and preserve; to establish; to retain.
- fixation muscles, muscles which prevent disturbance of body equilibrium generally, and fix limbs in case of limb-movements.
- flabellate (flăběl'āt) a. [L. flabellare, to fan.] Fan-shaped; appl. pectinate antennae with long processes.
- flabelliform (flabel'ifôrm) a. [L. flabellum, fan; forma, shape.]
 Fan-shaped.
- flabellinerved (flabel'inervd) a. [L. flabellum, fan; nervus, sinew.]
 Appl. leaves with many radiating nerves.
- flabellum (flăběl'ům) n. [L. flabellum, fan.] Any fan-shaped organ or structure; distal exite of branchiopodan limb; epipodite of

- certain crustacean limbs; terminal lobe of glossa in certain insects.
- flagellate (flaj'élat) a. [L. flagellum, whip.] Furnished with flagella; like a flagellum.
- flagelliform (flajěl'ífôrm) a. [L. flagellum, whip; forma, shape.]
 Lash-like; like a flagellum.
- flagellula (flăjěl'ūlă) n. [L. flagellula, dim. of flagellum, whip.] A flagellate zoospore or flagellispore.
- flagellum (flajel'ŭm) n. [L. flagellum, whip.] The lash-like process of many Protista and of cells, as in choanocytes and certain male gametes; external structure on basal joint of chelicera of Pseudoscorpiones; distal part of antenna in some arthropods, as in Diptera; a long slender runner or creeping stem.
- flame cells,—the terminal cells of branches of excretory system in many worms, with cavity continuous with lumen of duct, and containing a cilium or bunch of cilia, the motions of which give a flickering appearance similar to that of a flame; a pronephridiostome.
- flavedo (flave'dö) n. [L. flavus, yellow.] Exocarp of hesperidium; cf. albedo.
- flavescent (flaves'ent) a. [L. flavescere, to turn yellow.] Growing yellow.
- flavin (flā'vin) n. [L. flavus, yellow.] A water-soluble yellow pigment of cells; lyochrome.
- flavoxanthin (flā'vözăn'thin) n. [L. flavus, yellow; Gk. xanthos, yellow.] A yellow colouring matter in petals, as of Ranunculaceae; C₄₀H₅₆O₃.
- flex (fleks) v. [L. flectere, to bend.]
 To bend; appl. movement of limbs
- flexor (flek'sor) n. [L. flexus, bent.]
 A muscle which bends a limb, or part, by its contraction.
- flexor plate,—a median plate supporting praetarsus of insects, for attachment of tendon of claw flexor.

flexuous (flěk'sūŭs) a. [L. flexus, Curving ın a zig-zag manner; flexuose.

flexure (flěk'sūr) n. [L. flexus, bent.] A curve or bend; appl. curve in embryonic brain, curve of intestine.

float (flot) n. [A.S. fleotan, to float.] The pneumatophore of siphonophores; one of four tracheal sacs in aquatic larva of Culicidae; large spongy mass serving as a float in some pteridophytes.

floating ribs,—ribs not uniting at their ventral end with the ster-

num.

floccose (flŏk'ōs) a. [L. floccus, a lock of wool.] Covered with wool-like tufts; appl. bacterial growth.

floccular (flok'ūlar) a. [L. floccus, lock of wool.] Pert. the flocculus. flocculence (flok'ūlëns) n. [L. floccus,

lock of wool.] Adhesion in small flakes, as of a precipitate.

flocculent (flok'ūlënt) a. [L. floccus, lock of wool.] Covered with a soft waxy substance giving appearance of wool; covered with small woolly tufts.

flocculus (flok'ūlūs) n. [L.L. dim. of L. floccus, lock of wool.] A small accessory lobe on each lateral lobe of the cerebellum; a posterior hairy tuft in some Hymenoptera.

floccus (flok'ŭs) n. [L. floccus, lock of wool.] The tuft of hair terminating a tail; downy plumage of young birds; mass of hyphal filaments in algae and fungi.

flora (flo'ră) n. [L. flos, flower.] The plants peculiar to a country,

area, or period.

floral (flo'ral) n. [L. flos, flower.] Pert. the flora of a country or area; pert. flowers.

florescence (flōrĕs'ens) n. florescere, to begin to flower.] Bursting into bloom.

floret (flo'rët) n. [L. flos, flower.] One of the small individual flowers of a composite flower; flower with lemma and palea, of grasses.

floricome (flo'rikom) n. [L. flos, flower; coma, hair.] A form of branched hexaster spicule.

florigen (flo'rijen) n. [L. flos, flower; gignere, to produce.] A plant substance which stimulates change in buds to flowering condition; flowering hormone.

florigenic (flo'rijen'ik) a. [L. flos, flower; gignere, to produce.] Appl. principle originating in leaves which

stimulates flowering.

florula (flo'rūlă) n. [Dim. of flora.] Plant population of a small unit area, as of compost heap, etc.

floscelle (flösěl') n. [L. flosculus, little flower.] Flower-like structure round the mouth, composed of five bourrelets and five phyllodes, in some echinoids.

flosculus (flos'kūlus) n. [L. flosculus, little flower.] A small flower; a

floret, q.v.

floss (flos) n. [O.F. flosche, down.] A downy or silky substance; the loose pieces of silk in cocoon.

flower (flow'ër) n. [L. flos, flower.] The blossom of a plant, comprising generally sepals, petals, stamens, and pistil; a leafy shoot adapted for reproductive purposes.

flowering glume,—lemma, q.v.

fluviatile (floo'vĭătil) a. [L. fluviatilis, pert. river.] Growing in or near streams; inhabiting and developing in streams, appl. certain insect larvae; caused by rivers, appl. deposits.

fluviomarine (floo'viomaren') a. [L. fluvius, stream; mare, sea.] Pert. or inhabiting rivers and sea.

fluvioterrestrial (floo'viöteres'trial) [L. fluvius, stream; terra, land.] Found in streams and in the land beside them.

flux (fluks) n. [L. fluere, to flow.] Term appl. species that are not yet stable.

foetal (fē'tăl) a. [L. foetus. offspring.] Embryonic; pert. a foetus.

foetid glands,—small sac-like glands which secrete an ill-smelling fluid, in Orthoptera.

foetus (fē'tus) n. [L. foetus, offspring.] An embryo in egg or in uterus.

- foliaceous (fö'liā'shǔs) a. [L. folium, leaf.] Having the form or texture of a foliage leaf; thin and leaf-like.
- Folian process [C. Folli or Folius, Italian anatomist]. Anterior process of malleus; processus gracilis.

foliar (fö'liăr) a. [L. folium, leaf.] Pert. or consisting of leaves.

foliation (föliā'shūn) n. [L. folium, leaf.] The production of leaves;

leafing.

- folic (fō'līk) a. [I.. folium, leaf.] Appl. acid obtained from spinach, and liver extract, C₁₅H₁₅O₈N₅, and existing in various forms, with hacmatopoietic and other effects; e.g. vitamins Be. B₁₂, M, factors R and S, rhizopterine, etc.
- foliicolous (foliïk'ölŭs) a. [L. folium, leaf; colere, to dwell.] Growing on leaves; appl. certain fungi and lichens.
- foliobranchiate (fö'lĭöbrăng'kĭāt) a. [L. folium, leaf; branchiae, gills.] Possessing leaf-like gills.
- foliolae (fö liölē) n. plu. [L. folium, dim., leaf.] Leaf-like appendages of telum.
- foliolate (fō'liŏlāt) a. [L. folium, dim., leaf.] Pert., having, or like, leaflets.
- foliole (fō'liōl) n. [L. folium, dim., leaf.] Small leaf-like organ or appendage; a leaflet, as of a compound leaf.

foliose (fō'liōs) a. [L. folium, leaf.] With many leaves; leafy.

folium (fō'lĭum) n. [L. folium, leaf.]
A flattened structure in the cerebellum, expanding laterally into superior semilunar lobules.

follicle (föl'īkl) n. [L. folliculus, small sac.] A capsular fruit which opens on one side only; cavity or sheath; an ovarian follicle; a hair follicle.

- follicles of Langerhans [P. Langerhans, German anatomist]. Groups of cells in submucosa at junction of fore-gut and mid-gut of larval cyclostomes, secreting an insulinlike substance and being homologous to islets of Langerhans.
- follicle-stimulating hormone, a

gonadotrophic hormone, prolan A, which stimulates ovarian follicles and testis; symbol FSH.

follicular (fölĭk'ūlăr), folliculate (fölĭk'ūlāt) a. [L. folliculus, small sac.] Pert., like, or consisting of follicles; appl. an ovarian hormone.

folliculose (fŏlĭk'ūlōs) a. [L. folliculus, small sac.] Having follicles.

fontanelle (fŏn'tăněl) n. [F. fontanelle, little fountain.] A gap or space between bones in the cranium, closed only by membrane; depression on head of termites.

fonticulus (fŏntĭk'ūlūs) n. [L. fonticulus, dim. of fons, fountain.] A fontanelle; depression at anterior end of sternum, the jugular notch.

food-chain,—sequence of organisms in which each is food of a later member of the sequence.

food vacuole,—a small vacuole containing fluid and food-particles, in endosarc of many Protista.

- foot (foot) n. [A.S. fot, foot.] An embryonic structure in vascular cryptogams through which nourishment is obtained from prothallus; basal portion of sporophyte in mosses; an organ of locomotion, differing widely in different animals, from tube-foot of echinoderms, muscular foot of gastropods and other molluscs, tarsus of insects, to foot of vertebrates.
- foramen (förā'men) n. [L. foramen, opening.] The opening through coats of ovule; any small perforation; aperture through a bone or membranous structure.
- foramen (occipitale) magnum, the opening in occipital region of skull through which passes the spinal cord.
- foramen of Monro [A. Monro (primus), Scottish anatomist]. Interventricular foramen, passage between third and lateral ventricles; porta or foramen interventriculare.

foramina,-plu. of foramen.

foraminate (fŏrăm'înāt) a. [L. foramen, opening.] Pitted; having foramina or perforations. foraminiferous (főräm'ĭnĭf'črŭs) a. [L. foramen, opening; ferre, to carry.] Having foramina; containing shells of Foraminifera.

forceps (fôr'sĕps) n. [L. forceps, tongs.] The clasper-shaped anal cercus of some insects; large fighting or seizing claw of crabs and lobsters; fibres of corpus callosum curving into frontal and occipital lobes.

forcipate (fôr'sIpāt) a. [L. forceps, tongs.] Resembling forceps, or

forked like forceps.

forcipulate (fôrsip'ūlāt) a. [Dim. of L. forceps, tongs.] Shaped like a small forceps; appl. asteroid pedicellariae.

fore-brain,—prosencephalon, q.v.

forespore,—early stage in endospore formation, in bacteria.

forfex (fôr'fēks) n. [L. forfex, shears.] A pair of anal organs which open and shut transversely, occurring in certain insects.

forficate (fôr'fikāt) a. [L. forfex,

shears.] Deeply notched.

forma (fôr'mă) n. [L. forma, shape.]
Form; taxonomic unit consisting
of individuals that differ from those
of a larger unit by a single character; smallest category in botanical classification.

formation (fôrmā'shūn) n. [L. forma, shape.] Structure arising from an accumulation of deposits; the vegetation proper to a definite type of habitat; production.

formative (fôr'mătīv) a. [L. forma, shape.] Plastic; appl. matter which is living and developable.

formicarian (för'mikä'riän) a. [L. formica, ant.] Pert. ants; appl. plants which attract ants by means of sweet secretions.

formicarium (fôr'mĭkā'riŭm) n. [L. formica, ant.] Ants' nest, particularly an artificial arrangement for purposes of study; formicary.

fornicated (fôr'nīkā'tēd) a. [L. fornicatus, vaulted.] Concave within, convex without; arched.

fornix (fôr'niks) n. [L. fornix, vault.] An arched recess, as be-

tween eyelid and eye-ball, or between vagina and cervix uteri; an arched sheet of white longitudinal fibres beneath corpus callosum; scutum of Cheilostomata; one of arched scales in the orifice of some flowers.

fossa (fős'ă) n. [L. fossa, ditch.] A pit or trench-like depression.

fosse (fos) n. [L. fossa, ditch.] A fossa; a circular groove formed by upper part of parapet in sea-anemones.

fossette (fösět') n. [F. fossette, small pit, from L. fossa, ditch.] A small pit; a socket containing base of antennule in arthropods.

fossil (fos'il) n. [L. fossilis, dug up.]
Petrified animal or plant, or portion

thereof, as found in rocks.

fossiliferous (fős'îlĭf'ĕrŭs) a. [L. fossilis, dug up; ferre, to carry.]
Containing fossils.

fossorial (fösö'riăl) n. [L. fossor, digger.] Adapted for digging; appl. animals, claws, feet.

fossula (fős'ūlă) n. [Dim. of L. fossa, ditch.] A small fossa; small pit with reduced septa on one side of a corallite cup in Rugosa.

fossulate (fŏs'ūlāt) a. [Dim. of L. fossa, ditch.] With slight

hollows or grooves.

fossulet (fős'űlët) n. [Dim. of L. fossa, ditch.] A long narrow depression.

fourchette (foorshet') n. [F. fourchette, fork.] Frenulum of labia minora.

fovea (fō'vëā) n. [L. fovea, depression.] A small pit, fossa, or depression; a small hollow at leaf base in Isoëtes, containing a sporangium; pollinium base in orchids.

fovea centralis,—central and thinnest part of macula lutea, without rods and with long and slender cones.

fovea dentis,—facet on atlas, for articulation with dens of axis.

foveal (fô'veăl) a. [L. fovea, depression.] Pert. fovea; pert. fovea centralis; appl. cone vision. Opp. extrafoveal.

- foveate (fō'vëāt) a. [L. fovea, depression.] Pitted.
- foveola (föve'ölä) n. [L. foveola, small depression.] A small pit; a shallow cavity in bone; a small depression just above fovea in leaf of Isoëtes.
- foveolae opticae,—two pigmented areas in depressions of neural plate of amphibian embryo, the primordia of eyes.
- foveolate (fövē'olāt) a. [L. foveola, small depression.] Having regular small depressions.
- fragmentation (frag'menta'shun) n.
 [L. frangere, to break.] Division into small portions; nuclear division by simple splitting; amitosis.
- fraternal,- dizygotic, appl. twins.
- free (frē) a. [A.S. freo, acting at pleasure.] Motile; unattached; distinct; separate.
- free central placentation,—axile placentation, fixation of ovules to central axis of ovary.
- free-martin,—a sterile female twinborn with a male.
- frenate (frē'nāt) a. [L. frenare, to bridle.] Having a frenum or frenulum.
- frenulum (frěn'ūlům) n. [L. frenulum, dim. of frenum, bridle.] A fold of membrane, as of tongue, clitoris, etc.; a process on hindwing of Lepidoptera for attachment to fore-wing; a thickening of sub-umbrella of certain Scyphomedusae.
- frenum (frē'num) n. [L. frenum, bridle.] A frenulum; a fold of integument at junction of mantle and body of Cirripedia, ovigerous in Pedunculata; also fraenum.
- frigofuge (frig'öfüj) n. [L. frigor, cold; fugere, to flee.] An organism which does not tolerate cold.
- frond (frônd) n. [L. frons, leafy branch.] A leaf, especially of fern or palm; thallus of certain sea-weeds.
- frondescence (frondes'ens) n. [L. frondescere, to put forth leaves.]
 Development of leaves.

- frons (fronz) n. [L. frons, fore-head.] Forehead; or comparable structure.
- frontal (frun'tăl) a. [L. frons, forehead.] In region of forehead; appl. artery, vein, lobe, convolution; appl. head-organ of nemertines; a prostomial ridge of polychaetes; palps of certain nereids; specialised feeding surface in certain ciliates; ganglion, gland, and pore in insects.
- frontalis (fröntā'lĭs) n. [L. frons, forehead.] Frontal part of the scalp muscle or epicranius.
- frontocerebellar fibres, fibres passing from frontal region to cerebellum.
- frontoclypeus (frūn'töklĭp'ëŭs) n.
 [L. frons, forehead; clypeus, shield.] Frons and clypeus fused, in insects.
- fronto-ethmoidal,—pert. frontal and ethmoidal bones; appl. suture.
- frontonasal (frun'tönā'zāl) a. [L. frons, forehead; nasus, nose.] Pert. forehead or frontal region and nose; appl. ducts and process.
- frontosphenoidal (frun'tösfenoid'ăl)

 a. [L. frons, forehead; Gk. sphen,
 wedge; eidos, form.] Pert. frontal
 and sphenoid bones; appl. a process of zygomatic bone articulating
 with frontal.
- fructification (frūk'tĭfikā'shūn) n. [L. fructus, fruit; facere, to make.] Fruit formation; any spore-producing structure in cryptogams. fructose (frūk'tōs) n. [L. fructus.]
- fructose (frūk'tōs) n. [L. fructus, fruit.] Fruit-sugar; laevulose; $C_6H_{12}O_8$.
- frugivorous (froojiv'örŭs) a. [L. frux, fruit; vorare, to devour.] Fruit-eating; appl. certain animals.
- fruit (froot) n. [F. fruit, from L. fructus, fruit.] The fertilised and developed ovary of a plant.
- fruit-body,—the spore-bearing structure, as a sporangiocarp, basidiocarp, conidiocarp.
- fruit-spot,-sorus, as of ferns.
- frustule (frus'tūl) n. [L. frustulum, small fragment.] The siliceous twovalved shell and protoplasm of a diatom.

frutescent (frootĕs'ënt) a. [I..
frutex, shrub.] Shrub-like; fruticose.

frutex (froo'těks) n. [L. frutex, shrub.] Shrub.

fruticose (froo'tikōs) a. [L. frutex, shrub.] Like a shrub; appl. certain lichens.

fruticulose (frootik'ūlōs) a. [L. frutex, shrub.] Like a small shrub.

fucivorous (fūsĭv'örŭs) a. [L. fucus, seaweed; vorare, to devour.] Appl. seaweed-eating animals.

fucoid (fū'koid) a. [I.. fucus, seaweed; Gk. eidos, form.] Pert. or resembling seaweed.

fucosan (fū'kösăn) n. [L. fucus, seaweed.] Product of carbonassimilation in brown seaweeds.

fucoxanthin (fū'közăn'thĭn) n. [L. fucus, seaweed; Gk. xanthos, yellow.] The main carotenoid pigment of brown algae; C₄₀H₅₆O₆.

fugacious (fūgā'shūs) a. [L. fugax, swift.] Evanescent; falling off early; caducous; appl. petals, etc.

fulcral (fül'krăl) a. [L. fulcrum, support.] Pert. or acting as a fulcrum; appl. triangular plates aiding in movement of stylets in Hymenoptera.

fulcrate (fŭl'krāt) a. [L. fulcrum, support.] Having a fulcrum.

fulcrum (fūl'krūm) a. [L. fulcrum, support.] A supporting organ such as a tendril or stipule; sporophore in lichens; plate supporting rami of incus in mastax of rotifers; the lower surface of a ligula; a chitinous structure in base of insect rostrum; hinge-line of brachiopods; spine-like scale on anterior fin-rays of many ganoids.

function (fungk'shun) n. [L. functus, performed.] The action proper to

any organ or part.

functional (fungk'shönăl) a. [L. functus, performed.] Acting normally; acting or working part of an organ as distinct from remainder.

fundament,—primordium, q.v.

fundamentum,—hypocotyl, q.v. fundatrix (fūn'dātrīks) n. [L. fundare, to found.] Stem mother, a female founding a new colony by

a female founding a new colony by oviposition; appl. Aphides.

fundiform (fūn'dĭfôrm) a. [L. funda, sling; forma, shape.]
Looped; appl. a ligament of

penis.

fundus (fǔn'dǔs) n. [L. fundus, bottom.] The base of an organ, as of stomach, urinary bladder, etc.; boundary between underground and above-ground portions of plant axis.

fungicolous (funjik'ölus) a. [L. fungus, mushroom; colere, to inhabit.] Living in or on fungi.

fungiform (fūn')ifôrm) a. [L. fungus, mushroom; forma, shape.] Fungoid or shaped like a fungus; appl. tongue papillae.

fungine (fun'jin) n. [L. fungus, mushroom.] Chitinous substance forming cell-wall of fungi.

fungistatic (fŭn'jīstăt'īk) a. [L. fungus, mushroom; Gk. statikos, causing to stand.] Inhibiting the development of fungi.

fungivorous (fŭnjiv'örŭs) a. [L. fungus, mushroom; vorare, to devour.] Appl. fungus-eating animals and plants.

fungous (fung'gus) a. [L. fungus, mushroom.] With character or consistency of fungus; fungoid.

funicle (fū'nīkl) n. [L. funiculus, small cord.] An ovule stalk; a slender strand attaching peridiolum to peridium; a small cord or band, as of nerve fibres; a large double strand of cells passing from aboral end of coelom to aboral wall of zooecium of Molluscoidea; also funiculus.

funicular (fūnik'ūlăr) a. [L. funiculus, small cord.] Consisting of a small cord or band; pert. a funiculus or funicle.

funiculus (fūnĭk'ūlŭs) n. [L. funiculus, small cord.] A funicle, q.v.; one of the ventral, lateral, and dorsal columns of white matter of the spinal cord.

funnel (fŭn'ël) n. [L. fundere, to pour.] Siphon of cephalopods.

funnelform (fun'elform) a. [L. fundere, to pour; forma, shape.] Widening gradually from a narrow base; infundibuliform.

furca (fŭr'kă) n. [L. furca, fork.] The apophysis or entothorax of insect metathorax; forked intercoxal plate, as in Copepoda.

furcal (fŭr'kăl) a. [L. furca, fork.] Forked; appl. a branching nerve

of lumbar plexus.

furcasternum (fŭr'kăstër'nŭm) n. [L. furca, fork; sternum, breast-bone.]
Forked poststernite or sternellum in many insects.

furcate (fŭr'kāt) a. [L. furca, fork.] Branching like prongs of a fork.

furciferous (fŭršif'erūs) a. [L. furca, fork; ferre, to carry.] Bearing a forked appendage, as some insects.

furcula (fūr'kūlā) n. [L. furcula, dim. of furca, fork.] A forked process or structure; the merrythought bone; a transverse ridge in embryonic pharynx; partially fused abdominal appendages forming springing organ in Collembola.

furred (furd) a. [M.E. furren, sheath.] Having short decumbent hairs thickly covering the surface.

fuscin (fūs'sin) n. [L. fuscus, tawny.] A brown pigment in retinal epithelium.

fuseau (fūzō) n. [F. fuseau, from L. fusus, spindle.] A spindle-shaped structure; a spindle-shaped, thick-walled spore divided by septa, in certain fungi; a fusiform macroconidium.

fusi (fū'zī) n. plu. [L. fusus, spindle.] In spiders, organs composed of two retractile processes which issue from mammulae and form threads.

fusiform (fū'zifôrm) a. [L. fusus, spindle; forma, shape.] Spindle-shaped; tapering gradually at both ends; appl. innermost layer of cerebral cortex; appl. a gyrus of temporal lobe.

fusion-nucleus,—central nucleus of embryo-sac formed by fusion of odd nuclei from each end. fusulae (fū'zūlē) n. plu. [Dim. of L. fusus, spindle.] Spools, minute tubes of spinneret.

galactase (gălăk'tās) n. [Gk. gala, milk.] An enzyme, trypsin-like in action, found in milk.

galactin (gălăk'tīn) n. [Gk. gala, milk.] The prepituitary lactogenic hormone, prolactin; a polysaccharide occurring in certain plants, e.g. in lupin.

galactoblast (gălāk'töblāst) n. [Gk. gala, milk; blastos, bud.] A fat-containing globule or colostrum corpuscle in mammary acini.

galactophorous (găl'áktôf'örŭs) a. [Gk. gala, milk; pherein, to carry.] Lactiferous; appl. ducts of mammary glands.

galactosis (găl'ăktō'sĭs) n. [Gk. gala, milk.] Milk secretion.

galactotropic (gălăk'tötröp'ĭk) a. [Gk. gala, milk; trope, turn.] Stimulating milk secretion; appl. hormone: prolactin, q.v.

galbulus (găl'būlŭs) n. [L. galbulus, cypress nut.] A modified cone with fleshy scales, as in cypress.

galea (gāl'eā) n. [L. galea, helmet.]
A helmet-shaped petal, or other similarly-shaped structure; epicranial aponeurosis, the galea aponeurotica, of the scalp muscle or occipitofrontalis; galea capitis, thin sheath covering head of spermatozoon; outer division of stipes or endopodite of first maxilla of insects, itself divided into basigalea and distigalea; a prominence of movable digit of chelicerae in Pseudoscorpiones.

galeate (gāl'ēāt) a. [L. galea, helmet.] Helmet-shaped; hooded. Galen, veins of [Galen, Greek physician]. Internal cerebral veins and great cerebral vein formed by their union.

gall (gôl) n [A.S. gealla, gall.] Bile, secretion of liver. [L. galla, gallnut.] An excrescence on plants, caused by fungi, mites, and insects,

- especially by Cynipidae and Cecido-
- gall bladder, --- pear shaped of spherical sac which stores bile.
- galloxanthin (gălözăn'thin) n. [L. gallus, cock; Gk. xanthos, yellow.] Carotenoid pigment associated with retinal cones in domestic fowl.
- galvanotaxis (găl'vănötăk'sīs), galvanotropism (găl'vănŏt'röpīzm) n. [L. Galvani, Italian physiologist]. Response or reaction to electrical stimulus.
- gametal (gămē'tăl) a. [Gk. gametes, spouse.] Pert. a gamete; reproductive.
- gametangium (găm'ētăn'jiŭm) n. [Gk. gametes, spouse; anggeion, vessel.] A structure producing sexual cells.
- gametes (gămēts') n. plu. [Gk. gametes, spouse.] Cells derived from gametocytes which conjugate and form zygotes; sexual cells.
- gametic (gămět'ík) a. [Gk. gametes, spouse.] Pert. gamete; appl. a mutation occurring before maturation of gamete; appl. linkage.
- gametids (gămē'tidz) n. plu. [Gk. gametes, spouse.] Primary sporoblasts destined to become gametes.
- gametoblast (gămē'töblāst) n. [Gk. gametes, spouse; blastos, bud.]
 Plasson; formulative substance.
- gametocyst (gămē'tösīst) n. [Gk. gametes, spouse; kystis, bladder.] Cyst surrounding two associated free forms in sexual reproduction of gregarines.

gametocyte (gămē'tösīt) n. [Gk. gametes, spouse; kytos, hollow.]
The mother-cell of a gamete.

- gametogenesis (găme'töjen'esis) n. [Gk. gametes, spouse; genesis, origin.] Gamete formation; gametogeny.
- gametogenic (gămē'töjěn'ik) a. [Gk. gametes, spouse; genos, descent.] Arising from spontaneous changes in chromosomes of gametes; appl. variation.

- gametogonium (gămē'tögō'niŭm) n. [Gk. gametes, spouse; gonos, offspring.] A cell producing a gamete, a gametocyte.
- gametoid (gamet'oid) n. [Gk. gametes, spouse; eidos, form.] A structure behaving like a gamete, as apocytes uniting to form a zygotoid.
- gametokinetic (gămē'tökinēt'īk) a. [Gk. gametes, spouse; kinein, to move.] Stimulating gamete formation; appl. hormones, as folliclestimulating hormone or prolan A.
- gametophore (gāmē'töfōr) n. [Gk. gametes, spouse; pherein, to bear.] A special part of a gametophyte on which gametangia are borne; a hyphal outgrowth which fuses with a similar neighbouring outgrowth to form a zygospore.

gametophyll (gămē'töfil) n. [Gk. gametes, spouse; phyllon, leaf.] A modified leaf bearing sexual organs; a micro- or macro-sporophyll.

- gametophyte (găme tofit) n. [Gk. gametes, spouse; phyton, plant.]
 The gamete-forming phase in alternation of plant generations; haplophase; sexual generation of plants; pollen grain and embryo-sac; cf. sporophyte.
- gamic (găm'ik) a. [Gk. gamos, marriage.] Fertilised.
- gammation (gămā'shūn) n. [Gk. gammation, dim. of gamma.] An angular bar beside the branchial arches of Palaeospondylus.
- gamobium (gămō'bium) n. [Gk. gamos, marriage; bios, life.] The sexual generation in alternation of generations; opp. agamobium.
- gamocyst (găm'ösist) n. [Gk. gamos, marriage; kystis, bladder.] Oocyst, or sporocyst.
- gamodeme (găm'ödēm) n. [Gk. gamos, marriage; demos, people.] A deme forming a relatively isolated intrabreeding community.
- gamodesmic (găm'ödĕs'mik) a. [Gk. gamos, marriage; desma, bond.] Having the vascular bundles fused together instead of separated by connective tissue.

- gamogastrous (găm'ogăs'trūs) a. [Gk. gamos, marriage; gaster, belly.] Appl. a pistil formed by union of ovaries, but with styles and stigmata free.
- gamogenesis (găm'öjĕn'ësĭs) n. [Gk. gamos, marriage; genesis, descent.]
 Sexual reproduction.
- gamogenetic (găm'öjënět'ĭk) a. [Gk. gamos, marriage; genesis, descent.] Sexual; reproduced from union of sex elements.
- gamogony (gămŏg'önĭ) n. [Gk. gamos, marriage; gone, descent.]
 Sporogony in protozoa.

gamont (gămont') n. [Gk. gamos, marriage; on, being.] A sporont.

- gamopetalous (găm'opĕt'ălūs) a. [Gk. gamos, marriage; petalon, petal.] With coherent petals; sympetalous.
- gamophase (găm'ofāz) n. [Gk. gamos, marriage; phasis, aspect.] The haploid phase of a life-cycle; haplophase; cf. zygophase.
- gamophyllous (găm'öfil'ŭs) a. [Gk. gamos, marriage; phyllon, leaf.] With united perianth leaves; monophyllous.
- gamosepalous (găm'ösĕp'álūs) a. [Gk. gamos, marriage; F. sépale, sepal.] With coherent sepals; monosepalous.
- gamostele (găm'östē'lē) n. [Gk. gamos, marriage; stele, pillar.] Stele formed from fusion of several steles.
- gamostelic (găm'östē'lĭk) a. [Gk. gamos, marriage; stele, pillar.]
 Appl. condition in which steles of a polystelic stem are fused together.
- gamostely (găm'östē'lĭ) n. [Gk. gamos, marriage; stele, pillar.]
 The arrangement of polystelic stems when the separate steles are fused together surrounded by pericycle and endodermis.
- gamotropism (gămŏt'röpĭzm) n. [Gk. gamos, union; trope, turn.] Tendency to mutual attraction, exhibited by movements of gametes.

ganglia,—plu. of ganglion.

gangliar (găng'gliăr) a. [Gk. gangglion, little tumour.] Pert. a ganglion or ganglia. gangliate (găng'gliāt) a. [Gk. gangglion, little tumour.] Having ganglia.

- gangliform (găng'glifôrm) a. [Gk. gangglion, little tumour; L. forma, shape.] In the form of a ganglion.
- ganglioblast (găng'glĭoblăst) n. [Gk. gangglion, little tumour; blastos, bud.] Mother-cell of gangliocyte.
- gangliocyte (găng'gliosīt) n. [Gk. gangglion, little tumour; kytos, hollow.] A ganglion cell outside the central nervous system.

ganglioid (găng'glĭoid) a. [Gk. gangglion, little tumour; eidos, form.] Like a ganglion.

ganglion (găng'glion) n. [Gk. gangglion, little tumour.] A mass of nerve cell bodies and giving origin to nerve fibres; a nerve centre.

ganglionated (găng'glĭŏnā'tëd) a. [Gk. gangglion, little tumour.] Supplied with ganglia; gangliate.

- ganglioneural (găng'glĭonū'răl) a. [Gk. gangglion, little tumour; neuron, nerve.] Appl. a system of nerves, consisting of a series of ganglia connected by nerve strands.
- ganglioneuron (găng'gliönū'rŏn) n. [Gk. gangglion, little tumour; neuron, nerve.] A nerve cell of a ganglion.
- ganglionic (găng'gliŏn'îk) a. [Gk. gangglion, little tumour.] Pert., consisting of, or in neighbourhood of a ganglion; appl. layer of retina, arteries, arterial system of brain.
- ganglioplexus (găng'gliöplek'sŭs) n. [Gk. gangglion, little tumour; L. plexus, braided.] A diffuse ganglion.
- ganoblast (găn'öblăst) n. [Gk. ganos, sheen; blastos, bud.] An ameloblast.
- ganoid (găn'oid) a. [Gk. ganos, sheen; eidos, form.] Appl. scales of ganoid fishes, rhomboidal, joined like parquetry and consisting of a layer of bone with superficial enamel.
- ganoine (găn'öin) n. [Gk. ganos, sheen.] The outer layer of a ganoid scale, formed by the corium; enamel-like substance in formation of ameloblasts; ganoin.

- gape (gāp) n. [A.S. geapan, to open wide.] The distance between the open jaws of birds, fishes, etc.
- garland cells,—a chain of nephrocytes, in Diptera.
- gas gland,—glandular portion of airbladder of fishes.
- Gasserian ganglion [A. P. Gasser, German anatomist]. The semilunar ganglion on sensory root of fifth cranial nerve.

gastero-,-also gastro-.

- gasteromycetous (găs'tërömīsē'tŭs)

 a. [Gk. gaster, stomach; mykes,
 mushroom.] Having the spores
 developed in a gleba within a
 peridium.
- gastraea (găstrē'ā) n. [Gk. gaster, stomach.] A hypothetical gastrula-like animal; the ancestral metazoan, according to Haeckel.

gastraeum (găstrē'ŭm) n. [Gk. gaster, stomach.] Ventral side of body.

gastral (găs'trăl) a. [Gk. gaster, stomach.] Pert. stomach, as gastral cavity, cortex, layer, etc.

- gastralia (găstră'liă) n. plu. [Gk. gaster, stomach.] Microscleres in the gastral membranes of Hexactinellida; abdominal ribs, as in some reptiles.
- gastric (găs'trik) a. [Gk. gaster, stomach.] Pert. or in region of stomach; appl. arteries, glands, nerves, veins.
- gastrin (găs'trin) n. [Gk. gaster, stomach.] A hormone secreted by pyloric mucosa and which stimulates gastric secretion.

gastro-,--also gastero-.

- gastrocentrous (găs'trösen'trus) a. [Gk. gaster, stomach; kentron, centre.] Appl. vertebrae with centra formed by pairs of interventralia, while the basiventralia are reduced.
- gastrocnemius (găs'tröknē'mĭŭs) n. [Gk. gaster, stomach; kneme, tibia.] Large muscle of calf of leg.

gastrocoel (găs'trösēl) n. [Gk. gaster, stomach; koilos, hollow.] The archenteron of a gastrula.

- gastrocolic (găs'trökŏl'îk) a. [Gk. gaster, stomach; kolon, gut.] Pert. stomach and colon; appl. ligament, the greater omentum.
- gastrocutaneous (găs'trokūtā'nëŭs)
 a. [Gk. gaster, stomach; L. cutis,
 skin.] Appl. pores leading from
 intestine to surface in Hemichorda.
- gastrocystis (găs'trösis'tis) n. [Gk. gaster, stomach; kystis, bladder.]
 Blastocyst.
- gastrodermis (găs'trödēr'mīs) n. [Gk. gaster, stomach; derma, skin.] Enteroblast.
- gastroduodenal (găs'trödūodē'nāl)
 a. [Gk. gaster, stomach; L. duodeni, twelve each.] Pert. stomach and duodenum; appl. an artery.
- gastroepiploic (găs troepiplo'ik) a. [Gk. gaster, stomach; epiploon, omentum.] Pert. stomach and great omentum; appl. arteries, veins.
- gastrointestinal (găs'tröintěs'tinăl)
 a. [Gk. gaster, stomach; L. intus,
 within.] Pert. stomach and intestines.
- gastrolienal (găs'trölīē'năl) a. [Gk. gaster, stomach; L. lien, spleen.] Pert. stomach and spleen; appl. ligament; gastrosplenic.
- gastrolith (găs'trolith) n. [Gk. gaster, stomach; lithos, stone.] A mass of calcareous matter found on each side of gizzard of crustaceans before a moult.
- gastroparietal (gäs'tröpărī'etāl) a. [Gk. gaster, stomach; L. paries, wall.] Pert. stomach and body wall.
- gastrophrenic (găs'trofren'ik) a. [Gk. gaster, stomach; phren, midriff.] Pert. stomach and diaphragm; appl. ligament.
- gastropod (găs'tropŏd) n. [Gk. gaster, stomach; pous, foot.] A mollusc with ventral muscular disc adapted for creeping; gasteropod.
- gastropores (găs'tröpōrz) n. plu. [Gk. gaster, stomach; poros, channel.] The larger pores, for nutrient persons, of hydroid corals.
- gastropulmonary(găs'tröpul'mönări)
 a. [Gk. gaster, stomach; L. pulmo, lung.] Pert. stomach and lungs.

- gastrosplenic (găs'trösplēn'ĭk) a. [Gk. gaster, stomach; splen, spleen.] Pert. stomach and spleen; gastrolienal.
- gastrostege (găs'tröstēj) n. [Gk. gaster, stomach; stege, roof.] A ventral scale of snakes.
- gastrovascular (găs'trövăs'kūlăr) a. [Gk. gaster, stomach; L. vasculum, small vessel.] Serving both digestive and circulatory purposes, as canals of some Coelentera.
- gastrozooid (găs'trözô'oid) n. [Gk. gaster, stomach; zoon, animal; eidos, form.] In coelenterate colonies, the nutrient person with mouth and tentacles; trophozooid in some tunicates.
- gastrula (găs'troolă) n. [Gk. gaster, stomach.] The cup- or basinshaped structure formed by invagination of a blastula.
- gastrulation (găs'troolā'shun) n. [Gk. gaster, stomach.] Formation of gastrula from blastula by invagination.
- geitonogamy (gī'tŏnŏg'amĭ) n. [Gk. geiton, neighbour; gamos, marriage.] Fertilisation of a flower by another from the same plant.
- gelatigenous (jěl'átij'ënůs) a. [L. gelare, to congeal; Gk. -genes, producing.] Gelatine-producing.
- gelatine (jĕl'átĭn) n. [L. gelare, to congeal.] A jelly-like substance obtained from animal tissue.
- gelatinous (jěläťínůs) a. [L. gelare, to congeal.] Jelly-like in consistency. geminate (jěm'ināt) a. [L. gemini,
- twins.] Growing in pairs; binate; paired; appl. species or subspecies: corresponding forms in corresponding but separate regions, as reindeer and caribou.
- gemini (jem'ini) n. plu. [L. gemini, twins.] Bivalent chromosomes; pairs of paternal and maternal chromosomes at parasyndesis.
- geminiflorous (jĕm'iniflō'rūs) a. [L. gemini, twins; flos, flower.] Appl. a plant whose flowers are arranged in pairs.
- gemma (jěm'ă) n. [L. gemma, bud.]
 A bud or outgrowth of a plant

or animal which develops into a new organism; a leaf-bud, opp. flower-bud; a chlamydospore, q.v.; a hypothetical unit, q.v.

gemmaceous (jemā'shus) a. [L. gemma, bud.] Pert. gemmae or buds.

gemma-cup,—cyathus, q.v.

gemmate (jem'at) a. [L. gemmare, to bud.] Having buds.

- gemmation (jemā'shun) n. [L. gemma, bud.] Budding; budformation by means of which new independent individuals are developed in plants and animals; arrangement of buds.
- gemmiferous (jěmif'ërus) a. [L. gemma, bud; ferre, to bear.] Budbearing; gemmate.
- gemmiform (jĕm'ïfôrm) a. [L. gemma, bud; forma, shape.] Shaped like a bud; appl. pedicellariae of echinoderms.
- **gemmiparous** (jĕmĭp'ărŭs) a. [L. gemma, bud; parere, to produce.] Reproducing by bud-formation.
- **gemmulation** (jĕm'ūlā'shŭn) n. [L. gemmula, little bud.] Gemmule-formation.
- gemmule (jem'ūl) n. [L. gemmula, little bud.] A pangen; a moss bud; one of the internal buds of Porifera arising asexually and coming into activity on death of parent organism; one of the minute protoplasmic processes on branch of a dendrite, contact point in synapse.
- gena (jë'na) n. [L. gena, cheek.]
 The cheek or side part of head;
 antero-lateral part of prosoma of
 trilobites, and of insect head.
- genal (jě'năl) a. [L. gena, cheek.]

 Pert. the cheek; appl. facial suture
 of trilobites and angle of cheek.
- gene (jēn) n. [Gk. genos, descent.] A unit hereditary factor in the chromosome; also gen; regarded as multiple, composed of genomeres.
- gene flow,—the spreading of genes resulting from outcrossing and from subsequent crossing within a group; genorheithrum, q.v.
- gene mutation,—a heritable variation caused by changes at a particular locus; point-mutation.

- genecology (jěn'ēköl'öji) n. [Gk. genos, descent; oikos, household; logos, discourse.] Ecology in relation to genetics.
- Gené's organ [C. G. Gené, Italian zoologist]. Subscutal or cephalic gland secreting a viscid substance used in transferring eggs to dorsal surface, in ticks.

geneogenous (jěněoj'čnůs) a. [Gk. genea, birth; gennaein, to produce.]

Congenital.

geneology (jĕn'ĕŏl'ŏjĭ) n. [Gk. genos, descent; logos, discourse.] The study of development of individual and race; embryology and palaeontology combined.

genera,-Plu. of genus.

generalised (jen'eraliz'd) a. [L. generalis, of one kind.] Combining characteristics of two or more groups, as in many fossils.

generation (jĕn'ērā'shūn) n. [L. generatio, reproduction.] Production; formation; the individuals of a species equally remote from a common ancestor.

generative (jěn'ërātĭv)a. [L. generare, to beget.] Concerned in reproduction; appl. smaller of two cells into which a pollen grain primarily divides.

generative ferment,—a specific substance, present in small quantities in blood, necessary for formation of gonadial internal secretions.

generic (jënër'ĭk) a. (L. genus, race.]
Common to all species of a genus;
pert. a genus.

generitype (jënër'itip) n. [L. genus, race; typus, image.] The typical species of a genus.

genesiology (jěn'esĭŏl'ojĭ) n. [Gk. genesis, descent; logos, discourse.] Science dealing with reproduction.

genesis (jen'esis) n. [Gk. genesis, descent.] Formation, production, or development of a cell, organ, individual, or species.

genetic (jěně'tik) a. [Gk. genesis, descent.] Pert. genesis; pert.

genetics.

genetic factor,—gene, q.v.

- genetic spiral,—in spiral phyllotaxis, imaginary spiral line following points of insertion of successive leaves.
- genetics (jěnět'íks) n. [Gk. genesis, descent.] That part of biology dealing with heredity and variation.

genetype, --genotype.

- genial (jĕn'iăl) a. [Gk. geneion, chin.] Pert. the chin; appl. chuplates of reptiles; appl. tubercles on inside of mandible, for insertion of genioglossal and geniohyoid muscles.
- genic (jēn'ik) a. [Gk. genos, descent.]

 Pert. genes.

genic balance,—harmonious interaction of genes.

genicular (jënĭk'ūlār) a. [L. geniculum, little knee.] Pert. region of the knee; appl. arteries, etc., pert. geniculum.

geniculate (jënīk'ūlāt) n. [L. geniculum, little knee.] Bent like a kneejoint; appl. antenna; pert. geniculum, appl. a ganglion of the facial nerve; appl. bodies, lateral and medial corpora geniculata, constituting the metathalamus; having upper part of filament forming an angle more or less obtuse with lower.

geniculum (jenĭk'ūlŭm) a. [L. geniculum, little knee.] Sharp bend in a nerve; part of the facial nerve in temporal bone where it turns abruptly towards stylo-mastoid foramen.

genioglossal (jėnī'öglŏs'āl) a. [Gk. geneion, chin; glossa, tongue.] Connecting chin and tongue; appl. muscle; geniohyoglossal.

geniohyoid (jënī'öhī'oid) a. [Gk. geneion, chin; hyoeides, Y-shaped.]
Pert. chin and hyoid; appl. muscles.

genital (jĕn'ĭtăl) a. [L. gignere, to beget.] Pert. the region of reproductive organs; appl. corpuscles, glands, ridge, tubercle, etc.

genitalia (jen'ītālīā) n. plu. [L. gignere, to beget.] Genitals, the organs of reproduction, especially the external organs.

- genito-anal (jěn'ítöā'năl) a. [L. gignere, to beget; anus, vent.] In the region of genitalia and anus.
- genitocrural (jen'itökroo'răl) a. [L. gignere, to beget; crus, leg.] In the region of genitalia and thigh; appl. a nerve originating from first and second lumbar nerves.

genito-enteric (jën'îtöëntër'îk) a. [L. gignere, to beget; Gk. enteron, gut.] Pert. genitalia and intestine. genitofemoral (jën'îtöfëm'oral) a.

[L. gignere, to beget; femur, thigh-

bone.] Genitocrural.

Gennari's band [F. Gennari, Italian anatomist]. A layer of white fibres in middle cell-lamina of cerebral cortex, especially of occipital lobe; line of Gennari.

genoblast (jĕn'öblăst) n. [Gk. genos, offspring; blastos, bud.] A mature germ-cell exclusively male or female.

- genoholotype (jen'öhöl'ötip) n. [Gk. genos, race; holos, whole; typos, image.] A species defined as typical of its genus.
- genome (jen'om) n. [Gk. genos, offspring.] Minimum group or set of chromosomes derived from a zygote or gamete; genom.

genomere (jěn'ömēr) n. [Gk. genos, offspring; meros, part.] A unit of a gene, regarded as a multiple.

genonema (jěn'onē'mā) n. [Gk. genos, descent; nema, thread.] Axial thread on which genes are located in chromosome; axoneme; chromonema; a chromatid in its genetical aspect.

genonomy (jěnŏn'ömĭ) n. [Gk. genos, descent; nomos, law.] The study of laws of relationships with reference to classification of

organisms.

genophenes (jen'öfenz) n. plu. [Gk. genos, offspring; phainein, to appear.] Reaction types of the same genotype.

genorheithrum (jěn'örē'thrum) n. [Gk. genos, descent; rheithron, stream.] The passage or descent of genes in phylogenesis.

genosome (jen'osom) n. [Gk. genos, descent; soma, body.] The part

of the chromosome bearing the locus of a gene.

genospecies (jen'öspe'shez) n. [Gk. genos, race; L. species, particular kind.] A species consisting of individuals having the same genotype.

genosyntype (jěnösin'tīp) n. [Gk. genos, race; syn, with; typos, image.] A series of species together defined as typical of their genus.

- genotype (jen'ötīp) n. [Gk. genos, race; typos, image.] Genetic or factorial constitution of an individual; group of individuals possessing the same genetic constitution; biotype; genoplast; type species of a genus, generitype.
- genotypic (jen'ötip'ik) a. [Gk. genos, race; typos, image.] Pert. genotype; appl. characters arising from hereditary endowment.
- genovariation,—point mutation, q.v. genu (jěn'ū) n. [L. genu, knee.] A knee-like bend in an organ or part; anterior end of corpus callosum.
- genus (jē'nŭs) n., genera (jēn'ērā) plu. [L. genus, race.] A group of closely related species, in classification of plants or animals.
- genys (jěn'is) n. [Gk. genys, jaw.] Lower jaw.
- geobionts (jē'öbīönts) n. plu. [Gk. ge, earth; bion, living.] Organisms permanently inhabiting the soil.
- geobios (jē'öbī'ŏs) n. [Ğk. ge, earth; bios, life.] Terrestrial life; edaphon, a.v.
- geoblast (jē'öbläst) n. [Gk. ge, earth; blastos, bud.] A germinating plumule of which the cotyledons remain underground.

geocarpic (jē'ökâr'pĭk) a. [Gk. ge, earth; karpos, fruit.] Having the fruits maturing underground.

- geocryptophyte (jë'ökrip'töfit) n.
 [Gk. ge, earth; kryptos, hidden; phyton, plant.] A plant with dormant parts underground; geophyte.
- geology (jēŏl'öjĭ) n. [Gk. ge, earth; logos, discourse.] The science dealing with structure, activities, and history of the earth.

- geomalism (jēom'ālizm) n. [Gk. ge, earth; omalos, level.] Response to the influence of gravitation; horizontal habitus.
- geonastic (jēönās'tĭk) a. [Gk. ge, earth; nastos, pressed.] Curving towards the ground.
- geonemy (jēŏn'ēmi) n. [Gk. ge, earth; nemein, to inhabit.] The geographical distribution of organisms; biogeography; chorology.
- geophilous (jēŏf'īlŭs) a. [Gk. ge, earth; philein, to love.] Living in or on the earth.
- geophyte (jē'öfīt) n. [Gk. ge, earth; phyton, plant.] A land plant; a plant with dormant parts (tubers, bulbs, rhizomes) underground.
- geosere (jē'ösēr) n. [Gk. ge, earth; L. serere, to put in a row.] A sere originating on a clay substratum.
- geotaxis (jē'ötāk'sīs) n. [Gk. ge, earth; taxis, arrangement.] Locomotor response to gravity.
- geotonus (jēŏt'önŭs) n. [Gk. ge, earth; tonos, tension.] Normal position in relation to gravity.
- geotropism (jēōt'ropizm) n. [Gk. ge, earth; trope, turn.] Tendency to respond to stimulus of gravity, usually positive, by turning downwards, as in growth of a root.
- gephyrocercal (jef'iröser'käl, jefi'röser'käl) a. [Gk. gephyra, bridge; kerkos, tail.] Appl. secondary diphycercal caudal fin brought about by reduction of extreme tip of heterocercal or homocercal fin.
- geratology (jer'atol'oji) n. [Gk. geras, old age; logos, discourse.] Study of the factors of decadence and old age of populations; cf. gerontology.
- germ (jerm) n. [L. germen, bud.]
 A unicellular micro-organism; a seed; a bud; a developing egg.
- germ band,—primitive streak, of early embryo.
- germ-cell,—a reproductive cell, opp. somatic cell; a primitive male or female element.
- germ-centre,—an area of lymphcorpuscle division in nodules of lymph gland tissue.

- germ-disc,—a small green cellular
 plate of the germ tube of liverworts;
 cf. germinal disc.
- germ-layer,—an early differentiated layer of cells.
- germ nucleus,—an egg or sperm nucleus.
- germ plasm,—idioplasm, the physical basis of inheritance.
- germ pore,—the exit pore of a germ tube in the spore integument.
- germ stock,-stolon of tunicates.
- germ theory,—biogenesis; the theory that hving organisms can be produced or developed only from living organisms.
- germ track,—lineage of zygote in developing organism; continuity of germ cells.
- germ tube,—short filamentous tube put forth by a germinating spore.
- germ vitellarium, an organ, of platyhelminths, producing both ova and vitelline material.
- germ yolk gland,—in some Rhabdocoelida, an embryonic structure consisting of fertile portion of egg and a sterile portion which functions as a yolk gland feeding the fertile portion.
- germarium (jermā'riŭm) n. [L. germen, bud.] An ovary; distal portion of an ovariole.
- germen (jěr'měn) n. [L. germen, bud.] A mass of undifferentiated cells, the primary form of germ cells.
- germiduct (jër'midŭkt) n. [L. germen, bud; ducere, to lead.] Oviduct, of trematodes.
- germigen (jĕr'mĭjĕn) n. [L. germen, bud; generare, to beget.] Ovary, of trematodes.
- germinal (jĕr'mĭnăl) a. [L. germen, bud.] Pert. a seed, a germ-cell, or reproduction.
- germinal bands,—two sets of rows of cells in early development of annulates.
- germinal cells,—the cells concerned in reproduction, set apart early in embryonic life.
- germinal disc,—the disc-like area of an egg yolk on which segmentation first appears; blastodisc.

- germinal epithelium,—the layer of columnar epithelial cells covering the stroma of an ovary.
- **germinal layers,**—primary layers of cells in a developing ovum: epiblast, hypoblast, and later, mesoblast; histogens, q.v.

germinal lid,—operculum of a pollen-grain.

germinal spot,—the nucleolus of an ovum.

- germinal vesicle,—the nucleus of an ovum before formation of polar bodies.
- germination (jer'mina'shun) n. [L. germen, bud.] Beginning of growth; budding; sprouting; development.
- germination hormone, substance formed in endosperm of Gramineae and which stimulates growth of the coleoptile and inhibits that of the root; blastanin.

germiparity (jěr'mĭpăr'ĭtĭ) n. [L. germen, bud; parere, to beget.] Reproduction by germ-formation.

- germogen (jĕr'möjĕn) n. [L. germen, bud; Gk. genos, offspring.] The central cell of gastrula-like phase, or infusorigen, in development of Rhombozoa; the residual nucleus, or unused portion, after formation of rhombogen by division of primary germogen or primitive central cell.
- gerontal (jëron'tal) a. [Gk. geron, old man.] Senile.

gerontic (jeron'tik) a. [Gk. geron, old man.] Pert. old age; gerontal; appl. stage in phylogeny.

gerontology (jer'ontol'oji) n. [Gk. geron, old man; logos, discourse.] The study of senescence and senility; geratology, q.v.

gestalt (gestalt') n. [Ger. Gestalt, form.] Organised or unified response to an arrangement of stimuli; co-ordinated movements or configuration of motor reactions; a mental process considered as an organised pattern, involving explanation of parts in terms of the whole; a pattern considered in relation to background or environment; appl. morphology irrespective of taxonomic relationships.

gestation (jëstā'shun) n. [L. gestare, to bear.] The intra-uterine period in development of an embryo.

giant cells,—large nerve-cells in annelids; myeloplaxes; osteoclasts, large multinuclear protoplasmic masses found in marrow, spleen; megakaryocytes, q.v.; Langhans' cells, q.v.; Betz cells, q.v.; gigantocytes.

giant chromosomes,—polytene or large chromosomes, as in salivary gland cells of larval Diptera.

giant fibres,—greatly enlarged and modified nerve-fibres running longitudinally through ventral nerve cord of some invertebrates.

gibbous (gĭb'ūs) a. [L. gibbus, hump.] Inflated; saccate or pouched, as the lateral sepals of Cruciferae; gibbose.

gigantocyte (jīgăn'tösīt) n. [Gk. gigas, giant; kytos, hollow.] Giant

cell, q.v.

gill (gil) n. [M.E. gille, gill.] A plate-like or filamentous outgrowth; respiratory organ of aquatic animals; radial lamella on under side of pileus of agarics.

gill arch,—part of visceral skeleton in region of functional gills; branchial arch.

gill basket,—the branchial skeleton of lampreys, composed of continuous cartilage.

gill book,—the respiratory organ of certain Palaeostraca, consisting of a large number of leaf-like structures between which water circulates.

gill cleft,—a branchial cleft formed on side of pharynx.

gill cover,-an operculum.

gill helix,—a spirally coiled gill-like organ in certain Clupeidae.

gill plume,—the gill or ctenidium of the majority of Gasteropoda.

- gill pouch,—an oval pouch containing gills and communicating directly or indirectly with exterior, as in Myxine and Petromyzon.
- gill rakers,—small spine-like structures attached in a single or double row to branchial arches, preventing escape of food.

- gill remnants,—epithelial, postbranchial, or suprapericardial bodies arising in pharynx of higher vertebrates.
- gill rods,—gelatinous rods supporting the pharynx in Cephalochorda; branchial rays in certain fishes.
- gill slits,—a series of perforations leading from pharynx to exterior, persistent in lower vertebrates, embryonic in higher.

gingival (jǐnjī'văl) a. [L. gingivae,

gums.] Pert. the gums.

- ginglymoid (ging'glimoid) a. [Gk. gingglymos, hinge-joint; eidos, form.] Constructed like a hinge-joint.
- ginglymus (ging'glimus) n. [Gk. gingglymos, hinge-joint.] An articulation constructed to allow of motion in one plane only.

Giraldès' organ [J. A. C. C. Giraldès, Portuguese surgeon]. The para-

didymis.

- girdle (gër'dl) n. [A.S. gyrdan, to gird.] In appendicular skeleton, the supporting structure at shoulder and hip, each consisting typically of one dorsal and two ventral elements; transverse groove in Dinoflagellata, containing transverse flagellum and separating epicone and hypocone; the cingulum of diatoms.
- girdle bundles,—leaf-trace bundles which girdle the stem and converge at the leaf insertion, as in Cycadales.
- girdle scar,—a series of scale scars on axis of bud.

gito-,-geito-.

gizzard (gĭz'ard) n. [M.E. gizer, gizzard.] Muscular grinding chamber of alimentary canal of various animals; proventriculus of insects.

- glabella (glăběl'ă) n. [L. glaber, bald.] The space on forehead between superciliary ridges; the elevated median region of cephalic shield of Trilobita.
- glabrate (glā'brāt) a. [L. glaber, smooth.] Becoming smooth; glabrescent; with a nearly smooth surface.
- glabrous (glā'brus) a. [L. glaber,

smooth.] With a smooth, even surface.

- glacial (glā'siăl) a. [L. glacies, ice.] Pert. or appl. the Pleistocene epoch of the Quaternary period, characterised by periodic glaciation.
- gladiate (glăd'iāt) a. [L. gladius, sword.] Shaped like a sword; ensiform.
- gladiolus (glădi'ölŭs, glădiô'lŭs) n. [L. gladiolus, small sword.] The mesosternum or corpus sterni.
- gladius (glăd'ĭŭs) n. [L. gladius, sword.] The pen or chitinous shell in Chondrophora; cf. phragmocone, sepion.
- glairine (glā'rēn) n. [F. glaire, white of egg.] Glairy film found on thermal springs and formed by pectic zoogloea.
- gland (gland) n. [L. glans, acorn.]
 Single cell or mass of cells specialised for elaboration of secretions
 either for use in the body or for
 excretion; glans.
- gland cell,—an isolated secreting cell; a cell of glandular epithelium.
- glandula (glăn'dūlă) n. [L. glandula, small acorn.] A gland; a glutinous gland subserving cohesion of pollinia; arachnoid granulation on outer surface of dura mater.
- glandula vesiculosa, seminal vesicle.
- **glandulae Pacchionii,** arachnoideal granulations or Pacchionian bodies, q.v.
- glandular (glăn'dūlăr) a. [L. glandula, small acorn.] Pert. a gland; with secreting function.
- glandular epithelium,—the tissue of glands, composed of polyhedral, columnar, or cubical cells whose protoplasm contains or elaborates the material to be secreted.
- glandular tissue,—tissue of single or massed cells, parenchymatous and filled with granular protoplasm, adapted for secretion of aromatic substances in plants.
- glandulose serrate (glăn'dūlōssĕr'āt) a. [L. glandula, small acorn; serratus, sawn.] Having the serrations tipped with glands.

- glans (glans) n. [L. glans, acorn.] A nut; a hard, dry, indehiscent onecelled fruit, as an acorn; a gland; the glans penis; the glans clitoridis.
- Glaserian fissure [J. H. Glaser, Swiss anatomist]. Petrotympanic fissure.

glaucescent (glôsĕs'ënt) a. [L. glaucus, sea-green.] Somewhat glaucous.

glaucous (glôk'ŭs) a. [L. glaucus, sea-green.] Bluish green; covered with a pale green bloom.

gleba (glē'bà) n. [L. gleba, clod.] The central part of the sporophore in certain fungi; the spore-forming apparatus in certain plants.

glebula (glē'būlă) n. [L. glebula, small clod.] A small prominence

on a lichen thallus.

glenohumeral (glē'nöhū'meral) a. [Gk. glene, socket; L. humerus, humerus.] *Pert.* glenoid cavity and humerus; appl. ligaments.

glenoid (gle'noid) a. [Gk. glene, socket; eidos, form.] Like a appl. cavity into which socket; head of humerus fits, the mandibular fossa, and various ligaments.

glenoidal labrum,-a fibro-cartilaginous rim attached round the margin of glenoid cavity and of acetabulum.

glia (glē'ă, glī'ă) n. [Gk. glia, glue.] Gliacyte; neuroglia cell, a supporting cell of nervous tissue.

gliadin (glī'ādĭn) n. [Gk. glia, glue.] A substance interacting with glutenin to form gluten in cereals; the prolamine of wheat and rye seeds; formerly, any prolamine.

gliding growth,—see sliding growth. gliosomes (glī'ösōmz) n. plu. [Gk. glia, glue; soma, body.] Granules in protoplasm of neuroglia, possibly in relation with mitochondria.

Glisson, Glisson's capsule English physician]. fibrous Α capsule within liver, enclosing hepatic artery, portal vein, lymphatic vessels, and bile duct.

globate (glō'bāt) a. [L. globus, globe.] Globe-shaped; globular.

globigerina ooze,—sea-bottom mud which is largely composed of shells of Foraminifera.

globin (glō'bin) n. [L. globus, globe.] The basic protein constituent of haemoglobin.

globoid (glō'boid) n. [L. globus, globe; Gk. *eidos*, form.] spherical body in aleurone grains, a double phosphate of calcium and magnesium.

globose (globos') a. [L. globus, globe.] Spherical or globe-shaped;

globular.

globule (glob'ūl) n. [L. globulus, small globe.] Any minute spherical structure ; the antheridium of Characeae: globulus.

globulin (glob'ūlin) n. [L. globus, globe.] A protein, insoluble in water, such as fibrinogen, vitellin, crystallin, legumin.

globulose (glob'ūlos) a. [L. globus, globe.] Spherical; consisting of,

or containing globules.

globulus (glob'ūlus) n. [L. globulus, small globe.] A globule; spherical or club-shaped sensory organ at bifurcation of antenna in Pauropoda.

globus major and minor,—head and

tail of epididymis.

globus pallidus,—part of lentiform nucleus of corpus striatum.

[Gk. glochidiate (glökid'iāt) a. glochis, arrow-point.] Furnished with barbed hairs.

glochidium (glökid'iŭm) n. glochis, arrow-point; idion, dim.] Hairs bearing barbed processes seen on massulae of certain rhizocarps; the larva of fresh-water mussels such as Unio and Anodon.

gloea (glē'ā) n. [Gk. gloia, glue.] An adhesive secretion of some protozoa.

gloeocystidium (glē'ösistid'iŭm) n [Gk. gloios, sticky; kystis, bag; idion, dim.] A cystidium containing a slimy or oily substance.

glomera carotica,—carotid bodies,

q.v.

glomerular (glomer'ūlar) a. glomus, ball.] Pert. or like a glomerulus.

glomerulate (glomer'ūlāt) a. ſL. glomus, ball.] Arranged clusters.

glomerule (glom'ërūl) n. [L. glomus, ball.] A condensed cyme of almost sessile flowers; a compact cluster.

glomeruliferous (glŏmĕr'ūlĭf'ērŭs) a.
[L. glomus, ball; ferre, to carry.]
Having the flowers arranged in

glomerules.

glomerulus (glŏmĕr'ūlūs) n. [L. glomus, ball.] Network of capillary blood-vessels; inturned portion of a Bowman's capsule; oval body terminating olfactory fibres in rhinencephalon; a mass of interlacing intracapsular dendrites, in sympathetic ganglia; a glomerule.

glomus (glō'mūs) n. [L. glomus, ball.] A number of glomeruli run together; coccygeal and carotid bodies, consisting largely of chrom-

affin cells.

glossa (glŏs'ā) n. [Gk. glossa, tongue.] A tongue-like projection in middle of labium of insects.

glossal (glŏs'ăl) a. [Gk. glossa, tongue.] Pert. the tongue.

glossarium (glösā'rĭum) n. [Gk. glossa, tongue.] The slender-pointed glossa of certain Diptera.

glossate (glŏs'āt) a. [Gk. glossa, tongue.] Having a tongue or

tongue-like structure.

glosso-epiglottic (glòs'öĕpiglòt'ik) a. [Gk. glossa, tongue; epi, upon; glotta, tongue.] Pert. tongue and epiglottis; appl. folds of mucous membrane.

glossohyal (glŏs'ohī'āl) n. [Gk. glossa, tongue; hyoeides, T-shaped.] Median basihyal of fishes; entoglossum.

glosso-kinaesthetic area,—a brain area in Broca's convolution immediately connected with speech.

glossopalatine (glòs'öpāl'ātīn) a. [Gk. glossa, tongue; L. palatus, palate.] Connecting tongue and soft palate; appl. arch, muscle.

glossophagine (glosof'ajin) a. [Gk. glossa, tongue; phagein, to eat.]
Securing food by means of the

tongue.

glossopharyngeal (glos'öfarin'jeäl)
a. [Gk. glossa, tongue; pharynx,

gullet.] Pert. pharynx and gullet; appl. ninth cranial nerve.

glossopodium (glŏs'öpō'dĭŭm) n.
[Gk. glossa, tongue; pous, foot.]
The sheathing leaf-base of Isoëtes.

glossotheca (glos'öthē'kă) n. [Gk. glossa, tongue; theke, box.] The proboscis-covering part of pupal integument of insects.

glottis (glŏt'ĭs) n. [Gk. glotta, tongue.] The opening into the windpipe.

glucase (glook'ās) n. [Gk. glykys, sweet.] A plant enzyme which produces grape sugar from maltose.

glucokinin (glook'okī'nĭn) n. [Gk. glykys, sweet; kinein, to move.] A plant substance capable of reducing blood-sugar.

glucoproteins,—see glycoproteins. glucose (glook'ös) n. [Gk. glykys, sweet.] The grape sugar of plants and animals; dextrose, C₆H₁₂O₆.

gluma (gloom'ă) n. [L. gluma, husk.] A bract at base of a grass inflorescence or spikelet; a chaffy or membranous bract; glume.

glumaceous (gloomā'shūs) a. [L. gluma, husk.] Dry and scaly like glumes; formed of glumes.

glume,—gluma, q.v.; lemma, q.v. glumiferous (gloomif'ërüs) a. [L. gluma, husk; ferre, to bear.]

Bearing or producing glumes. glumiflorous (gloom'iflo'rŭs) a. [L.

gluma, husk; flos, flower.] Having flowers with glumes or bracts at their bases.

glutaeal (gloote'ăl) a. [Gk. gloutos, buttock.] Pert. or in region of buttocks; appl. arteries, muscles, nerves, tuberosity, veins.

glutaeus (glootē'ŭs) n. [Gk. gloutos, buttock.] A muscle of the buttock.

glutathione (gloot'āthī'ön) n. [L. gluten, glue; Gk. theion, sulphur.] A sulphur-containing tri-peptide found in different tissues and capable of being alternately reduced and oxidised; C₁₀H₁₇O₆N₃S.

gluten (gloot'en) n. [L. gluten, glue.]
A nitrogenous substance obtainable from some cereals, a product of

gliadin and glutenin.

- glutenin (gloot'ënin) n. [L. gluten, glue.] A substance of cereals interacting with gliadin to form gluten.
- glutinous (gloot'inus) a. [L. gluten, glue.] Having a sticky or adhesive surface.
- glycerin (glis'erin), glycerol (glis'ërôl) n. [Gk. glykys, sweet.] The sweet principle of natural fats and oils; $C_8H_5(OH)_3$.

glycogen (glī'köjěn) n. [Gk. glykys, A carbohydrate storage sweet.] product of plants and animals, $(C_6H_{10}O_5)x$; animal starch.

glycogenase (glī'köjenās') n. glykys, sweet; -genes, producing.] An enzyme which causes synthesis of storage glycogen in liver.

glycolysis (glīkŏl'isis) n. [Gk. glykys, sweet; lyein, to loosen.] Decomposition of glucose or of glycogen,

by hydrolysis.

- **glycophyte** (glī'köfīt) n. [Gk. glykys, sweet; phyton, plant.] A plant unable to thrive on substratum containing more than 0.5 per cent. sodium chloride in solution; opp. halophyte.
- glycoproteins (glī'köprō'tëĭnz) n. plu. [Gk. glykys, sweet; protos, first.] Compounds of protein with a carbohydrate, including mucins and mucoids; mucoproteins.

glycosecretory (glī'kösēkrē'törī) a. [Gk. glykys, sweet; L. secretus, set apart.] Connected with the secre-

tion of glycogen.

glycotropic (glī'kötrŏp'ik) a. ſGk. glykys, sweet; trope, turn.] Appl. factor secreted by prepituitary and which inhibits peripheral action of insulin; glycotrophic.

gnathic (năth'ik) a. [Gk. gnathos, jaw.] Pert. the jaw; gnathal.

- gnathion (nath'ion) n. [Gk. gnathos, jaw.] Lowest point of the median line of the lower jaw.
- gnathism (năth'izm) n. [Gk. gna-Formation of jaw thos, jaw.] with reference to degree of projection.
- gnathites (nath'its) n. plu. gnathos, jaw.] The buccal appendages of arthropods.

gnathobase (nāth'öbās) n. gnathos, jaw; basis, base.] inwardly turned masticatory process on protopodite of appendages near mouth of Crustacea.

gnathochilarium (nāth'ökīlā'rĭŭm) n. [Gk. gnathos, jaw; cheilos, lip.] First maxillae and sternal plate in Pauropoda, united in Diplopoda.

gnathopod (nåth'öpod) n. gnathos, jaw; pous, foot.] crustacean limb in oral region modified to assist with food.

gnathopodite (năth'ŏpödīt) n. [Gk. gnathos, jaw; pous, foot.] A

maxilliped of an arthropod.

gnathos (na'thos) n. [Gk. gnathos, jaw.] A median sclerite on ventral side of ninth tergum in Lepidoptera.

- gnathosoma (nåth'ösö'mä) n. Gk. gnathos, jaw; soma, body.] mouth region, including oral appendages, of Acarina.
- gnathostegites (năthos'tějīts) n. plu. [Gk. gnathos, jaw; stege, roof.] Pair of covering plates for mouth parts of some crustaceans.

gnathostomatous (nåth'östöm'ätus) a. [Gk. gnathos, jaw; stoma, mouth.] With jaws at the mouth.

- gnathotheca (nåth'öthē'kā) n. [Gk. gnathos, jaw; theke, case.] The horny outer covering of a bird's lower jaw.
- gnathothorax (nåth'öthö'räks) [Gk. gnathos, jaw; thorax, chest.] The part of the cephalothorax posterior to protocephalon, in Malacostraca.
- gnesiogamy (nē'siŏg'āmi) n. [Gk. gnesios, lawful; gamos, marriage.] Fertilisation by an individual of the same species; intraspecific zygosis. goblet cells,-mucus-secreting cells

of columnar epithelia; chalice cells.

Golgi apparatus or complex [C]. Golgi, Italian histologist]. constituents, localised or diffuse, often consisting of separate elements, the Golgi bodies, batonettes, dictyosomes or pseudochromosomes, containing lipoprotein, and concerned with cellular synthesis and secretion; originally apparato reticolare, canalicular system, internal reticular apparatus, the reticulum being possibly an artefact.

Golgi, organs of,-Golgi-Mazzoni

corpuscles.

Golgi-Mazzoni corpuscles [C. Golgi and V. Mazzoni, Italian histologists]. Cylindrical end-organs or small Pacinian corpuscles at junction of tendon and muscle.

golgiokinesis (gŏl'jĭökĭn \tilde{e} 'sĭs) n. [C. Golgi; Gk. kinesis, movement.] Division of the Golgi apparatus

during mitosis.

golgiosomes (gŏl'jĕösōmz) n. plu. [C. Golgi; Gk. soma, body.] Golgi bodies or material produced by division of the Golgi apparatus during mitosis.

gomphosis (gomfo'sis) n. [Gk. gomphos, bolt.] Articulation by insertion of a conical process into a socket, as of roots of teeth into alveoli.

gonad (gŏn'ad) n. [Gk. gone, birth.] A sexual gland, either ovary, or

testes, or ovotestis.

gonadectomy (gŏn'ăděk'tömĭ) n. [Gk. gone, birth; ek, out; tome, cutting.] Excision of gonad, castration in the male, spaying in female.

gonadin (gŏnā'dĭn) n. [Gk. gone, Active principle of sex birth.] glands controlling secondary sexual

characteristics.

gonadotrophins (gŏnădŏt'rŏfĭnz) n. plu. [Gk. gone, birth; trephein, to nourish.] Two prepituitary hormones: 1. Follicle-stimulating hormone or prolan A; gametogenetic or gametokinetic hormones; thylakentrin. 2. Luteinising or interstitial-cell-stimulating hormone; prolan B; metakentrin. Chorionic gonadotrophin secreted by chorionic cells of placenta and excreted in pregnancy urine, resembling but not identical with luteinising hor-Serum gonadotrophins: mone. hormone in follicle-stimulating blood of pregnant mares, luteinising hormone in that of women.

gonadotropic (gŏn'adötrŏp'ik) a. [Gk. gone, birth; trope, turn.]

Affecting the gonad; appl. prepituitary hormones and certain hormones obtained from urine and other body fluids and tissues, particularly during pregnancy; appl. effects; gonadotrophic.

gonadotropin (gŏn'ādŏt'röpin) n. [Gk. gone, birth; trope, turn.] Any gonadotropic hormone or substance.

gonaduct (gŏn'ădŭkt) n. [Gk. gone, birth; L. ductus, led.] Gonoduct. gonal (gon'al) n. [Gk. gone, birth.] Appl. middle portion of genital ridge which alone forms functional

gonad; gonidial, q.v.

gonangium (gönän'jiŭm) n. gone, birth; anggeion, vessel.] A gonotheca; a dilated cup of perisarc protecting the blastostyle of Calyptoblastea.

gonapophyses (gŏn'ăpŏf'ĭsēz) n. plu. [Gk. gone, birth; apo, from; phyein, to grow.] Chitinous outgrowths or valves subserving copulation in insects; the component parts of a sting.

gone (gŏn'ē) n. [Gk. gone, generation. | One of four daughter cells of an auxocyte; the generative portion of a gonad; an organism possessing a gone. v. To produce a gone.

gongylidia (gŏng'gĭlĭd'ĭă) n. plu. [Gk. gongylos, round; idion, dim.] Hyphal swellings or modifications in fungi cultivated by certain ants.

gonia (go'niă) n. plu. [Gk. gone, seed.] Primitive sex cells, spermatogonia or oogonia.

goniale (gōniâ'lë) n. [Gk. gonia, In some vertebrates a bone of lower jaw beside articular.

gonic (gŏn'ik) a. [Gk. gone, generation.] Pert. gones; pert. semen.

gonidangium (gon'idan'jium) [Gk. dim. of gone, seed; anggeion, vessel.] A structure producing or containing gonidia.

gonidia (gŏnĭd'iă) n. plu. dim. of gone, seed.] Minute reproductive bodies of many bacteria; asexual non-motile reproductive cells produced upon gametophytes; algal constituents of lichens. Sing. gonidium.

- gonidial (gŏnĭd'ĭăl) a. [Gk. dim. of gone, seed.] Pert. gonidia.
- gonidiferous (gonidif'ërus) a. [Gk. dim. of gone, seed; L. ferre, to carry.] Bearing or producing gonidia.

gonidimium (gŏn'idim'iŭm) n. [L.L. dim. of Gk. gone, seed.] A gonidial structure smaller than a gonidium and larger than a gonimium.

gonidiogenous (gŏnid'iŏj'enŭs) a. [Gk. dim. of gone, seed; -genes, producing. Bearing or producing gonidia; gonidiferous.

gonidioid (gŏnĭd'ioid) a. [Gk. dim. of gone, seed; eidos, form.] Like a gonidium; appl. certain algae.

gonidiophore (gonid'iöfor) n. [Gk. dim. of gone, seed; pherein, to bear.] An aerial hypha supporting a gonidangium.

gonidiophyll (gŏnĭd'iöfĭl) n. [Gk. dim. of gone, seed; phyllon, leaf.] A gametophyte leaf bearing gonidia.

gonidium,—sing. of gonidia.

gonimium (gŏnim'ium) n. [Gk. gonimos, productive.] One of the bluishgreen gonidia of certain lichens.

gonimoblasts (gŏn'imöblăsts) n. plu. [Gk. gonimos, productive; blastos, bud.] Filamentous outgrowths of a fertilised carpogonium of certain algae.

gonimolobe (gŏn'ĭmölōb) n. gonimos, productive; lobos, lobe.] A group of carposporangia borne on a gonimoblast.

gonion (gōnī'ŏn) n. [Gk. gonia, angle.] The angle point on the lower jaw.

gonoblast (gŏn'öbläst) n. [Gk. gonos, offspring; blastos, bud.] A reproductive cell in animals.

gonoblastid (gŏn'öblas'tid) n. gonos, offspring; blastos, bud; A blastostyle of idion, dim.] Hydrozoa; gonoblastidium.

gonocalyx (gŏn'ökā'lĭks) n. [Gk. gonos, offspring; kalyx, cup.] The bell of a medusiform gonophore.

gonocheme (gŏn'ö $k\bar{e}m$) n. [Gk. gonos, offspring; ochema, support.] A medusoid bearing sex-cells, in Hydrozoa.

gonochorism (gŏn'ökō'rĭzm) n. [Gk. gonos, offspring; chorismos, separation.] The history or development of sex differentiation; sex determination; dioecism.

gonochoristic (gŏn'ökōrĭs'tĭk) [Gk. gonos, offspring; choristos, separated.] Having the sexes separate; producing distinct males and females; dioecious.

gonocoel (gŏn'ösēl) n. [Gk. gone, seed; koilos, hollow.] The cavity

containing the gonads.

gonocytes (gŏn'ösīts) n. plu. [Gk. gone, seed; kytos, hollow.] Sexual cells of sponges; mother-cells of ova and spermatozoa.

gonodendron (gŏn'öděn'drŏn) [Gk. gonos, offspring; dendron, tree.] A branching blastostyle in Physalia.

gonoduct (gŏn'ödŭkt) n. [Gk. gonos, birth; L. ductus, led.] A genital duct leading from gonad to exterior. gonogenesis (gŏn'öjĕn'ësĭs) n. [Gk.

gone, seed; genesis, descent.l Gametogenesis.

gonomery (gŏnŏm'ërĭ) n. [Gk. gonos, descent; meros, part.] Theory that paternal and maternal chromosomes remain in separate groups throughout life; separate grouping of paternal and maternal chromosomes during cleavage stages of some organisms.

gononephrotome (gŏn'önĕf'röt \bar{o} m) n. [Gk. gone, seed; nephros, kidney; temnein, to cut.] Embryonic segment containing primordia of the urinogenital system.

gononucleus (gŏn'önū'klëŭs) n. [Gk. gonos. begetting; L. nucleus, kernel.] The generative nucleus or micronucleus of many Protozoa.

gonophore (gŏn'öfōr) n. [Gk. gone, seed; pherein, to bear.] An elongation of thalamus between corolla and stamens; a reproductive zooid in a hydroid colony.

gonoplasm (gŏn'öpläzm) n. gone, seed; plasma, mould.] generative part of protoplasm.

gonopodium (gŏn'öpō'dĭŭm) n. [Gk. gone, seed; pous, foot.] The modified anal fin serving as copulatory organ in male poeciliid fishes; gonopod or clasper of male myriopods and insects.

gonopore (gŏn'öpōr) n. [Gk. gone, seed; poros, channel.] Reproduc-

tive aperture.

- gonosome (gŏn'ösōm) n. [Gk. gone, seed; soma, body.] The reproductive zooids of a hydrozoan colony collectively.
- gonosphaerium (gŏn'ösfē'rĭŭm) n.
 [Gk. gone, seed; sphaira, globe.]
 An oosphere.
- gonostyle (gŏn'östīl) n. [Gk. gonos, birth; stylos, pillar.] The blastostyle; sexual palpon or siphon of Siphonophora; clasper of Diptera.
- gonotheca (gŏn'othē'kā) n. [Gk. gonos, birth; theke, cup.] A transparent protective expansion of the perisarc round a blastostyle or gonophore.

gonotokont,—an auxocyte, q.v.

- gonotome (gŏn'ŏtōm) n. [Gk. gonos, birth; temnein, to cut.] An embryonic segment containing the primordium of the gonad.
- gonotrema (gŏnötrē'mă) n. [Gk. gonos, offspring; trema, hole.] Genital aperture, as in Arachnida; gonotreme.
- gonozooid (gŏn'özō'oid) n. [Gk. gonos, birth; zoon, animal; eidos, form.] A gonophore or reproductive individual of a hydrozoan colony; a zooid containing a gonad. gonydial (gŏnid'iāl) a. [Gk. genys,

gonydial (gönid'iál) a. [Gk. geny. lower jaw.] Pert. a gonys.

gonys (gŏn'is) n. [Gk. genys, lower jaw.] Lower part or keel of bird's bill.

Graafian follicle [R. de Graaf,
Dutch appetemint] A vesicular

Dutch anatomist]. A vesicular capsule in ovary and surrounding an ovum; ovisac with developing ova.

Graber's organ,—a complex larval

organ, presumably sensory, in Tabanidae.

gracilis (grās'ilis) n. [L. gracilis, slender.] A superficial muscle on medial side of the thigh; a fasciculus of medulla oblongata; nucleus of grey matter ventral to clava. graduated (grād'ūātëd) a. [L. gradus, step.] Tapering; becoming longer or shorter by steps.

graft-hybrid,—an individual formed from graft and stock, and showing characteristics of both; graft chimaera.

grain (grān) n. [L. granum, grain.]
The caryopsis or seed of cereals;
a granular prominence on the back of a sepal.

graminifolious (grăm'inĭfō'liŭs) a. [L. gramen, grass; folium, leaf.]

With grass-like leaves.

graminivorous (grām'inĭv'örūs) a. [L. gramen, grass; vorare, to eat.] Grass-eating.

- Grandry's corpuscle [— Grandry, Belgian anatomist]. An end-organ of touch, in beak and tongue of birds.
- granellae (grăněl'ē) n. plu. [L. dim. of granum, grain.] Oval, refractile granules consisting chiefly of barium sulphate, found in the tubes of certain Sarcodina.
- granellarium (grăn'elā'riŭm) n. [L. dim. of granum, grann.] The system of granellae-containing tubes of Sarcodina.
- granose (grăn'ōs) a. [L. granum, grain.] In appearance like a chain of grains, like some insect antennae; moniliform.
- granular (grăn'ūlăr) a. [L. granum, grain.] Consisting of grains or granules; appearing as if made up of granules.
- granulation (grăn'ūlā'shŭn) n. [L. granum, grain.] A grain-like formation or eminence; appl. arachnoid elevations or Pacchionian glands on outer surface of dura mater.
- granule (grăn'ūl) n. [L. granulum, small grain.] A small particle of matter; a small grain.
- granule cells,—ovoid or spheroid cells formed of soft protoplasm containing basiphil granules.
- granule glands,—the prostate glands of flatworms; skin glands of amphibians.
- granules of Nissl,—see Nissl granules.

- granulocytes (grăn'ūlösīts) n. plu. [L. granulum, small grain; Gk. kytos, cell.] Granular white blood corpuscles or polymorphs; myeloid cells formed in bone marrow.
- graphiohexaster (graf'iöheksas'ter) n. [Gk. graphis, style; hex, six; aster, star.] A hexaster spicule with long outwardly-directed filamentous processes from four rays.

grater (grā'tër) n. [O.F. grater, to scrape.] A denticle of Eunice.

gravid (grav'id) a. [L. gravidus, loaded.] Appl. female with eggs,

or pregnant uterus.

graviperception (grav'ipërsep'shun) n. [L. gravis, heavy; percipere, to feel.] Irritability to gravity; geotropic reaction.

gravitational (grav'ītā'shönal) a. [L. gravis, heavy.] Appl. water in excess of soil requirements, which sinks under action of gravity and drains away.

gravity (grăv'iti) n. [L. gravitas, heaviness.] The force of attraction of all bodies towards each other; the tendency of terrestrial bodies to drawn towards the earth's

green glands,-the excretory antennary glands of certain Crustacea.

- gregaloid (greg'aloid) a. [L. grex, flock; Gk. eidos, form.] Appl. colony of protozoa of indefinite shape, usually with gelatinous base, formed by incomplete division of individuals or partial union of adults.
- gregarious (grega'rius) a. [L. grex, flock.] Tending to herd together; colonial; growing in clusters.

grey matter,—tissue abundantly supplied with nerve cells, of greyish colour, internal to white matter in spinal cord, external in cerebrum.

- nerve fibres, --- semitransparent, grey or yellowish-grey, gelatinous non-medullated nervefibres, comprising most of the fibres of the sympathetic system and some of the cerebro-spinal; amyelinate fibres.
- groin (groin) n. [A.S. grynde, depres-

sion.] The depressed part of body between abdomen and thigh.

groove (groov) n. [Dut. groef, channel.] Any channel, furrow, or depression, as carotid, costal, optic, primitive vertebral groove.

ground tissue,—see conjunctive parenchyma.

growing point,—a part of plant body at which cell-division is localised, generally terminal and composed of meristematic cells.

growth factor G,—vitamin B₂ or riboflavin, q.v.

growth hormones, — in animals, growth - promoting pituitary secretions; in plants, auxins.

grumose (groom'os) a. [L. grumus, hillock.] Clotted; knotted; collected into granule masses; grumous.

grumulus (groo'mūlŭs) n. [Dim. of L. grumus, hillock.] Polar organ or caudal cell cluster in insect embryo.

gryochrome (grī'ökrōm) a. [Gk. gry, morsel; chroma, colour.] With Nissl granules irregularly scattered; appl. neurones, as in spinal ganglia.

guanidine (gwân'idin) n. [Peruvian, huano, dung.] A substance produced by oxidation of guanin, whose metabolism is regulated by parathyroids; CH₅N₃.

guanine (gwân'ĭn) n. [Peruvian, huano, dung.] A purine base found in some plants, teleosts, mammals, etc.; $C_5H_5ON_5$.

guanophore (gwân'öfor) n. [Peruvian, huano, dung; Gk. pherein, to bear.] A yellow pigment-bearing cell; an iridocyte.

guanylic (gwân'ilik) a. [Peruvian, huano, dung.] Appl. a nucleic acid, yielding guanin, found in pancreas and liver.

guard (gârd) n. [O.F. guarder, to guard.] Rostrum of a belemnite.

guard cells, -- cells surrounding stomata of aerial epidermis of plant tissue.

gubernacular (gū'bërnăk'ūlăr) a. [L. gubernaculum, rudder.] Pert. the gubernaculum.

- gubernaculum (gū'bërnăk'ūlŭm) n. [L. gubernaculum, rudder.] A cord stretching from epididymis to scrotal wall; mesocardial ligament; tissue between gum and dental sac of permanent teeth; strands of blastostylar ectoderm between gonophore and gonotheca in Hydromedusae; a posterior flagellum functioning as a rudder.
- guest insect,—an insect living or breeding in the nest of another.
- gula (gū'lă) n. [L. gula, gullet.]
 The upper part of throat; median ventral sclerite of insect head.
- gulamentum (gū'lămēn'tŭm) n. [L. gula, gullet; mentum, chin.] Plate formed by fusion of gula and submentum in insects.
- gular (gū'lār) a. [L. gula, gullet.]

 Pert. throat; appl. median and lateral plates between rami of mandible in Crossopterygii and Polypterini. n. An anterior unpaired horny shield on plastron of Chelonia.
- gullet (gŭl'ët) n. [O.F. goulet, from L. gula, gullet.] The oesophagus, a muscular canal extending from mouth cavity to stomach; the canal between cytostome and endoplasm of Ciliata.
- gum (gum) n. [L. gummi, gum.] An exudation of certain plants and trees; vegetable mucilage.
- gummiferous (gumif'ërus) a. [L. gummi, gum; ferre to carry.]
 Gum-producing or exuding.
- gummosis (gumö'sis) n. [L. gummi, gum.] Condition of plant tissue when cell-walls become gummy.
- **gums** (gumz) n. plu. [A.S. goma, jaws.] Dense fibrous tissues investing jaws; gingivae.
- gustatory (gūs'tātöri) a. [L. gustare, to taste.] Pert. sense of taste; appl. cells, hairs, pores, calyculus (q.v.), nerves.
- gut (gǔt) n. [A.S. gut, channel.]
 Intestine or part thereof, according to structure of animal.
- gutta (gūt'ā) n. [L. gutta, drop.] A small spot of colour on insect wing or elsewhere.

- guttate (gŭt'āt) a. [L. gutta, drop.] Having drop-like markings.
- guttation (guta'shun) n. [L. gutta, drop.] Formation of drops of water on plants from moisture in air; exudation of aqueous solutions, as through hydathodes, or by sporangiophores, or by nectaries.
- guttiform (gŭt'ifôrm) a. [L. gutta, drop; forma, shape.] Drop-like; in the form of a drop.
- guttulate (gŭt'ūlāt) a. [L. guttula, small drop.] In the form of a small drop, as markings.
- gymnanthous (jimnăn'thŭs) a. [Gk. gymnos, uncovered; anthos, flower.]
 With no floral envelope; achlamydeous.
- gymnetrous (jǐmnē'trŭs) a. [Gk. gymnos, naked; etron, abdomen.]
 Without an anal fin.
- gymnoarian (jim'nōā'riān) a. [Gk. gymnos, naked; oarion, small egg.] Appl. gonads when naked, or not enclosed in coelomic sacs; opp. cystoarian.
- gymnoblastic (jim'nöblăs'tik) a. [Gk. gymnos, naked; blastos, bud.] Without hydrothecae and gonothecae; appl. certain Coelenterata.
- gymnocarpic (jim'nökâr'pik) a. [Gk. gymnos, uncovered; karpos, fruit.] With naked fruit; appl. lichens with uncovered apothecia, mosses with expanded hymenium; gymnocarpous.
- gymnocidium (jim'nösĭd'ĭŭm) n. [Gk. gymnos, uncovered; oikos, house; idion, dim.] A basal swelling of certain moss capsules.
- gymnocyte (jim'nösīt) n. [Gk. gymnos, uncovered; kytos, hollow.] A cell without a defining cell-wall; opp. lepocyte.
- gymnocytode (jim'nösi'töd) n. [Gk. gymnos, naked; kytos, hollow; eidos, form.] Cytode without cell-wall or nucleus.
- gymnogenous (jǐmnōj'ënūs) a. [Gk. gymnos, naked; genos, offspring.] Naked when born; appl. birds.
- gymnogynous (jimnöj'inüs) a. [Gk. gymnos, naked; gyne, female.]
 With exposed ovary.

- gymnoplast (jim'nöpläst) n. [Gk. gymnos, naked; plastos, formed.] Protoplasm without definite formation or cell-wall.
- gymnopterous (jimnŏp'tërŭs) a. [Gk. gymnos, naked; pteron, wing.] Having bare wings, without scales; appl. insects.

gymnorhinal (jim'nöri'năl) a. [Gk. gymnos, naked; rhis, nose.] With nostril region not covered by feathers, as in some birds.

gymnosomatous (jǐm'nösŏm'ātŭs) a. [Gk. gymnos, naked; soma, body.] Having no shell or mantle, as certain molluscs.

gymnospermous (jimnosper'mus) a. [Gk. gymnos, uncovered; sperma, seed.] Having seeds not enclosed in a true ovary, as conifers.

gymnospore (jim'nöspōr) n. [Gk. gymnos, naked; sporos, seed.] A naked germ or spore not enclosed in a protective envelope.

- gymnostomatous (jim'nöstöm'ätüs)

 a. [Gk. gymnos, naked; stoma, mouth.] Naked-mouthed; having no peristome, appl. mosses; gymnostomous.
- gynaecaner (jin'ēkā'nër) n. [Gk. gyne, woman; aner, man.] A male ant resembling a female; a gynaecomorphic male.

gynaecium,—gynoecium, q.v.

gynaecoid (jine koid) n. [Gk. gyne, woman; eidos, form.] An egg-laying worker ant.

gynaecophore (jinē'köfōr) n. [Gk. gyne, woman; pherein, to carry.] Canal or groove of certain worms formed by inrolling of sides, in which the female is carried; gynaecophoric or gynaecophoral groove.

gynander (jřnăn'dër) n. [Gk. gyne, female; aner, male.] A gynandromorph.

gynandrism (jinan'drizm) n. [Gk. gyne, women; aner, man.] Hermaphroditism.

gynandromorph (jinan'drömôrf) n.
[Gk. gyne, woman; aner, man; morphe, form.] An individual exhibiting a spatial mosaic of male and female characters; cf. intersex.

- gynandromorphism (jřnăn'drömôrfizm). [Gk. gyne, woman; aner, man; morphe, form.] Condition of being a gynandromorph or manifesting a mosaic of male and female sexual characters, as having one side characteristically male, the other female.
- gynandrophore (jinăn'dröför) n. [Gk. gyne, woman; aner, man; pherein, to carry.] An axial prolongation bearing a sporophyll.
- gynandrosporous (jinăn'dröspō'rūs)
 a. [Gk. gyne, woman; aner, man; sporos, seed.] With androspores adjoining the oogonium, as in some algae.
- gynandrous (jīnān'drŭs) a. [Gk. gyne, woman; aner, man.] Having stamens fused with pistils, as in some orchids.
- gynantherous (jǐnăn'thërus) a. [Gk. gyne, woman; anthos, flower.]
 Having stamens converted into pistils.
- gynase (ji'nās) n. [Gk. gyne, woman.]
 A female-determining factor in the form of an enzyme or hormone.
- gynatrium (jinā'triūm) n. [Gk. gyne, woman; L. atrium, entrance-hall.] Female genital pouch or vestibulum, of certain insects.
- gyne (jī'nē) n. [Gk. gyne, woman.] A female ant.

gynecium,—gynoccium, q.v.

- gynetype (jin'étip) n. [Gk. gyne, woman; typos, pattern.] Type specimen of the female of a species.
- gynobase (jin'öbās) n. [Gk. gyne, woman; L. basis, base.] A gynoe-cium-bearing receptacle of certain plants; condition in which style appears to arise from ovary.
- **gynobasic style**,—a style arising from base of carpel.
- gynodioecious (jinödiē'siūs) a. [Gk. gyne, woman; dis, twice; oikos, house.] Appl. plants producing female or hermaphrodite flowers only.
- gynoecium (jinē'siŭm) n. [Gk. gyne, woman; oikos, house.] The pistils, carpels, or female organs of a flower.

gynogenesis (ji'nöjēn'esis) n. [Gk. gyne, woman; genesis, descent.]
Development from eggs penetrated by the spermatozoon but not embodying its nucleus, as in some nematodes.

gynogonidia (jš'nogonid'iă) n. plu. [Gk. gyne, woman; gonidion, small seed.] Female sexual elements formed after repeated division of parthenogonidia in Mastigophora.

gynomerogony (ji'nömerog'oni) n. [Gk. gyne, female; meros, part; gone, generation.] The development of an egg fragment, obtained before fusion with male nucleus, and containing maternal chromosomes only.

gynomonoecious (jl'nomonē'sīŭs) a. [Gk. gyne, woman; monos, alone; oikos, house.] Appl. plants with pistillate and hermaphrodite flowers only.

gynophore (ji'nofōr) n. [Gk. gyne, woman; pherein, to carry.] A stalk supporting the ovary; elongation of thalamus between stamens and pistil; female gonophore.

gynospore (jřínöspōr) n. [Gk. gyne, female; sporos, seed.] Female spore; megaspore; embryo-sac.

gynostegium (jǐn'östē'jǐum) n. [Gk. gyne, woman; stege, roof.] A protective covering for a gynoecium.

gynostemium (jin'östē'miŭm) n. [Gk. gyne, woman; stemon, warp.] The column composed of united pistil and stamens in orchids.

gypsophil (jĭp'söfil) a. [Gk. gypsos, chalk, gypsum; philein, to love.] Thriving in soils containing chalk or gypsum; gypsophilous; calcicolous, calciphil.

gypsophyte (jip'söfit) n. [Gk. gypsos. chalk, gypsum; phyton, plant.] A gypsophil plant; calcicole, calcipete, calciphile, calciphyte.

gyral (jī'rāl) a. [L. gyrus, circle.]
Pert. a gyrus; pert. circular or spiral movement.

gyration (jīrā'shun) n. [L. gyrare, to revolve.] Rotation, as of cells; a whorl of a spiral shell.

gyre (jīr) n. [Gk. gyros, L. gyrus,

circle.] Circular movement; spiral coiling, as of chromatids.

gyrencephalic (ji'rĕnkĕfāl'ik, -sĕf-) a. [Gk. gyros, circle; engkephalos, brain.] Having cerebral convolutions; gyrencephalous. ()pp. lissencephalic.

gyri,—plu. of gyrus.

gyrose (jī'rōs) a. [L. gyrare, to revolve.] With undulating lines; sinuous.

gyrus (jī'rūs) n. [L. gyrus, circle.] A cerebral convolution; a ridge winding between two grooves.

Η

habenula (hābēn'ūlă) n. [L. habena, strap.] A name appl. certain bandlike structures. a. Habenular; appl. a commissure of epithalamus.

habitat (hāb'ītāt) n. [L. habitare, to inhabit.] The locality or external environment in which a plant or animal lives.

habituation (hābīt'ūā'shūn) n. [L. habituare, to bring into a habit.] The adjustment, effected in a cell or in an organism, by which subsequent contacts of the same stimulus produce diminishing effects.

habitus (hāb'ītūs) n. [L. habitus, appearance.] The general appearance or conformation characteristic of a plant or an animal; constitutional tendency.

hadrocentric (hăd'rösĕn'trik) a. [Gk. hadros, thick; kentron, centre.] With phloem surrounding xylem.

hadrome (hād'rōm) n. [Gk. hadros, thick.] Conducting tissue of xylem; hadromestome.

Haeckel's law [E. H. Haeckel, German zoologist]. Biogenetic law; recapitulation theory, q.v.

haem (hēm) n. [Gk. haima, blood.]
A blood substance, oxidising to haematin; C₃₄H₃₂O₄N₄Fe.

haemachrome (hē'mākrōm) n. [Gk. haima, blood; chromos, colour.] Colouring matter found in blood.

haemacyte (hē'măsīt) n. [Gk. haima, blood; kytos, hollow.] A blood corpuscle. haemad (hē'măd) adv. [Gk. haima, blood.] Situated on same side of vertebral column as heart.

haemal (hē'māl) a. [Gk. haima, blood.] Pert. blood or bloodvessels; situated on same side of vertebral column as heart.

haemamoeba (hē'māmē'bā) n. [Gk. haima, blood; amoibe, change.] Protozoon with an amoeboid trophozoitic stage parasitic in a red blood-corpuscle.

haemangioblast (hēmān'jīoblāst) n. [Gk. haima, blood; anggeion, vessel; blastos, bud.] A blood island, q.v.

haemapoietic (hē'măpoiet'īk) a. [Gk. haima, blood; poiein, to form.]
Blood-forming; haemopoietic.

haemapophysis (hē'māpŏf'isis) n. [Gk. haima, blood; apo, from; phyein, to grow.] One of plate-like or spine-like processes growing from the latero-ventral surfaces of a vertebral centrum.

haematal (hěm'ātāl) a. [Gk. haima, blood.] Pert. blood or blood-vessels. haematid (hěm'ātīd) n. [Gk. haima, blood.] Red blood-corpuscle.

haematin (hēm'ātīn) n. [Gk. haima, blood.] A pigment formed by decomposition of haemoglobin, containing iron and having the property of carrying oxygen; protohaem; C₃₄H₃₅O₅N₄Fe.

haematobic (hěm'ātō'bĭk) a. [Gk. haima, blood; bios, life.] Living

in blood.

haematobium (hěm'átō'bĭŭm) n. [Gk. haima, blood; bios, life.] An organism living in blood.

haematoblast (hěm'ātöblāst) n. [Gk. haima, blood; blastos, bud.] Λ cell that will develop into a red blood-corpuscle; thrombocyte; blood platelet.

haematochrome (hěm'átökrōm) n.
[Gk. haima, blood; chroma, colour.]
A carotenoid red pigment of certain

haematocryal (hēm'ātökrī'āl) a. [Gk. haima, blood; kryos, cold.] Cold-blooded.

haematocyanin (hěm'ătösī'anin) n.

[Gk. haima, blood; kyanos, dark blue.] Haemocyanin, q.v.

haematocytozoon (hěm'ătösī'tözō'ŏn) n. [Gk. haima, blood; kytos,
hollow; zoon, animal.] An intracorpuscular blood parasite.

haematodocha (hēm'ātödōk'ā) n. [Gk. haima. blood; doche, receptacle.] A fibro-elastic bag at base of palpal organ in Araneae.

haematogen (hēm'ātöjēn) n. [Gk. haima, blood; genos, birth.] A nucleoprotein containing iron.

haematogenesis (hēm'ātöjēn'esis) n. [Gk. haima, blood; genesis, descent.] The formation of blood.

haematogenous (hēm'ātŏj'ēnŭs) a.
[Gk. haima, blood; genos, birth.]
Formed in blood; derived from blood.

haematoidin (hem'atoid'in) n. [Gk. haima, blood; eidos, form.] An iron-free derivative of haemoglobin, forming crystals in blood clots, and identical with bilirubin.

haematolysis (hēm'ātŏl'īsĭs) n. [Gk. haima, blood; lysis, loosing.] Haemolysis, q.v.

haematophagous (hēm'ātŏf'āgūs) a. [Gk. haima, blood; phagein, to eat.] Feeding on blood, or obtaining nourishment from blood.

haematophyte (hěm'ătöfit) n. [Gk. haima, blood; phyton, plant.] A vegetable micro-organism in blood.

haematoporphyrin (hēm'ātöpôr'fīrīn) n. [Gk. haima, blood;
porphyra, purple.] An iron-free
pigment formed by decomposition
of haematin; C₃₄H₃₃O₆N₄.

haematosis (hem'ato'sis) n. [Gk. haimatoein, to change to blood.]

Blood-formation.

haematothermal (hēm'ătöther'măl)

a. [Gk. haima, blood; thermos,
warm.] Warm-blooded.

haematozoon (hēm'ātözō'ŏn) n. [Gk. haima, blood; zoon, animal.] An

animal parasitic in blood. haemerythrin (hēm'ĕrĭth'ı

haemerythrin (hēm'ērīth'rīn) n. [Gk. haima, blood; erythros, red.] A red respiratory pigment of corpuscles in body fluid of sipunculids and some annelids; haemoerythrin.

- haemic (hē'mik) a. [Gk. haima, blood.] Pert. blood.
- haemin (hē'min) n. [Gk. haima, blood.] Haem; c'ilor de formed in blood clot: C₃₄H₃₂O₄N₄FeCl.
- haemoblast (hē'moblāst) n. [Gk. haima, blood; blastos, bud.] A cell which gives rise to an erythroblast; haematoblast.
- haemochromes (hē'mökrōmz) n. plu. [Gk. haima, blood; chroma, colour.] Blood pigments, as haemoglobin, haemocyanin, chlorocruorin, erythrocruorin, haemoerythrin.
- haemochromogen (hē'mökrō'möjēn)
 n. [Gk. haima, blood; chroma, colour; genos, birth.] A chromoprotein produced by alkali and reduction of haemoglobin.

haemoclastic (hē'möklās'tĭk) a. [Gk. haima, blood; klastos, broken.] Breaking down blood cells, appl. tissues; opp. haemoplastic.

haemocoele (hē'mösēl) n. [Gk. haima, blood; korlos, hollow.] An expanded portion of the blood system which replaces the true coelom.

haemoconia (hē'mökŏn'iā) n. [Gk. haima, blood; konis, dust.] Minute fat droplets entering or leaving cells; chylomicrons, q.v.

haemocyanin (hē'mosi'anin) n. [Gk. haima, blood; kyanos, dark blue.]
A haemoglobin-like blood pigment containing copper instead of iron, in molluses. crustaceans, and some arachnids.

haemocyte (hē'mösīt) n. [Gk. haima, blood; kytos, hollow.] A blood cell, as in insects.

haemocytoblast (hē'mösī'töblăst) n. [Gk. haima, blood; kytos, hollow; blastos, bud.] Primitive stem cell from which all blood cells are derived; a lymphoid haemoblast; lymphoidocyte.

haemocytolysis (hē-mösītŏl'īsīs) n. [Gk. haima, blood; kytos, hollow; lyein, to dissolve.] Breaking up of red blood-corpuscles by solution

haemocytotrypsis (hē'mösī'tötrĭp'-sīs) n. [Gk. haima, blood; kytos,

- hollow; *tribein*, to rub.] Breaking up of blood-corpuscles by pressure.
- haemoerythrin (hē'möčrīth'rīn) n.
 [Gk. haima, blood; erythros, red.]
 A red respiratory pigment in certain invertebrates; haemerythrin.
- haemofuscin (hē'möfus'sīn) n. [Gk haima, blood; L. fuscus, tawny.]
 A yellow blood pigment deposited under various pathological conditions.
- haemoglobin (hē'möglō'bin) n.
 [Gk. haima, blood; globos, sphere.]
 The red respiratory pigment of blood
 of vertebrates, consisting of haematin united to globin.
- haemohistioblast (hē'möhīs'ttöblāst) n. [Gk. haima, blood; histion, tissue; blastos, bud.] A free macrophage in blood, especially of veins.
- haemoid (hē'moid) a. [Gk. haima, blood; cidos, form.] Resembling blood.
- haemolymph (hē'mölimf) n. [Gk. haima, blood; L. lympha, water.] A fluid found in coelom of some invertebrates, regarded as equivalent to blood and lymph of higher forms.
- haemolysin (hē'mŏlī'sin) n. [Gk. hama, blood; lyein, to dissolve.] A substance developed in or added to blood serum, capable of destroying red blood-corpuscles.
- haemolysis (hēmöl'īsīs) n. [Gk. haima, blood; lysis, loosing.] The lysis or solution of red blood-corpuscles; erythrocytolysis; laking.
- haemophilia (hěm'öfil'iă) n. [Gk. haima, blood; philos, loving.] Absence of ready coagulation of shed blood, a sex-linked hereditary characteristic.
- haemoplasmodium (hē'möplăzmō'-dīum, -plăs-) n. [Gk. haima, blood; plasma, mould.] A unicellular parasite of blood.
- haemoplastic (hē'möplăs'tĭk) a. [Gk. haima, blood; plastos, formed.]
 Blood-forming; haemopoietic.
 Opp. haemoclastic.

haemopoiesis (hē'möpoi'ēsis) n.
[Gk. haima, blood; poiesis, making.] The formation and development of blood cells.

HAE-

haemopoietic (hē'möpoiet'ik) a. [Gk. haima, blood; poietiikos productive.] Blood-forming; pert. haemopoiesis; haemoplastic.

haemopsonin (hēmöp'sönin) n. [Gk. haima, blood; opsonein, to cater.] An opsonin for erythrocytes.

haemorrhoidal (hēmöroid'āl) a. [Gk. haima, blood; rhein, to flow.] Rectal, appl. blood-vessels, nerve.

haemosiderin (hēmösīdē'rĭn) n. [Gk. haima, blood; sideros, iron.] A yellow pigment of blood giving an iron reaction.

haemostatic (hē-möstāt'īk) a. [Gk. haima, blood; statikos, causing to stand.] Appl. membrane crossing joint between trochanter and femur in autotomy of limb of some arthropods.

haemotoxin (hē'mötök'sin) n. [Gk. haima, blood; toxikon, poison.] A toxin which produces haemolysis.

haemotropic (hē'mötrŏp'īk) a. [Gk. haima, blood; trope, turn.] Affecting or acting upon blood.

haemozoin (hē'mözō'in) n. [Gk. haima, blood; zoon, animal.] Granules of a black pigment, the residue from digestion of haemoglobin by malarial parasites.

hair (hār) n. [A.S. haer.] Any epidermal filamentous outgrowth consisting of one or more cells, varied in shape; a thread-like or filamentous outgrowth of epidermis of animals; a setum, q.v.

hair cells,—sensory cells in organ of Corti.

hair follicle,—tubular sheath formed by invagination of epidermis and surrounding base of hair.

half-inferior,—having ovary but partially adherent to calyx.

half-spindle,—unipolar spindle, as in meiosis of some insects.

half-terete,—rounded on one side, flat on the other.

haliplankton,-haloplankton, q.v.

Haller's organ [G. Haller, German zoologist]. A tarsal chemoreceptor in ticks.

HAM-

hallux (hăl'ŭks) n. [L. hallux, great toe.] First digit of hind-limb.

halm,-haulm.

halobios (hăl'öbī'ŏs) n. [Gk. hals, sea; bios, life.] Sum total of organisms living in the sea.

halolimnic (hăl'ölim'nik) a. [Gk. hals, sea; limne, marsh.] Pert. marine organisms modified to live in fresh water.

halophilous (hălŏf'ilŭs) a. [Gk. hals, salt; philein, to love.] Salt-loving; thriving in presence of salt; halophilic.

halophyte (hăl'öfīt) n. [Gk. hals, salt; phyton, plant.] A shore plant; plant capable of thriving on salt, impregnated soils.

haloplankton (hăl'öplăng'ktŏn) n. [Gk. hals, sea; plangktos, wandering.] The organisms drifting in the sea; also haliplankton.

halosere (hăl'ösēr) n. [Gk. hals, salt; L. serere, to put in a row.] A plant succession originating in a saline area.

halteres (hăltē'rēz) n. plu. [Gk. halter, weight.] A pair of small capitate bodies representing rudimentary posterior wings in Diptera; balancers, poisers.

hamate (hā'māt) a. [L. hamatus, hooked.] Hooked or hook-shaped at the tip; hamose; uncinate.

hamatum (hāmā'tum) n. [L. hamatus, hooked.] The unciform bone in the carpus, probably corresponding to fourth and fifth distalia of a typical pentadactyl limb.

hamirostrate (hā'mirŏs'trāt) a. [L. hamus, hook; rostrum, beak.]
Having a hooked beak.

hamstrings,—tendons of insertion of the posterior femoral muscles, i.e., of semitendinosus, semimembranosus, and biceps.

hamula (hām'ūlă) n. [L. hamulus, little hook.] Retinaculum, of insects; fused ventral appendages acting with caudal furcula in springtails or Collembola; hamulus, q.v.

hamular (hăm'ūlăr) a. [L. hamus, hook.] Hooked; hook-like.

hamulus (hăm'ūlūs) n. [L. hamulus, little hook.] A hooklet, or hook-like process; minute hook-like process on distal barbules which aid in interlocking of feather barbs; retinaculum of Hymenoptera.

hamus (hā'mŭs) n. [L. hamus, hook.] Hooked part of uncus in

male Lepidoptera.

hapaxanthous (hăp'āksăn'thŭs) a. [Gk. hapax, once; anthos, flower.] With only a single flowering period; hapaxanthic; opp. pollakanthic.

haplobiont (hap löbi ont) n. [Gk. haploos, simple; bion, living.] An organism characterised by one kind of individual. Opp. diplobiont.

haplocaulescent (hāp'lokôlĕs'ent) a. [Gk. haploos, simple; L. caulis, stem.] With a simple axis, i.e. capable of producing seed on the main axis.

haplochlamydeous (hap'loklamid'eus) a. [Gk. haploos, simple; chlamys, cloak.] Having rudimentary leaves in connection with sporophylls.

haplo-diploid (hăp'lödīp'loid) a. [Gk. haploos, simple; diploos, double; eidos, form.] Appl. sexdifferentiation in which the male is haploid, the female diploid.

haplodont (hăp'lödönt) a. [Gk. haploos, simple; odous, tooth.]
 Having molars with simple crowns.

haploid (hāp'loid) a. [Gk. haploos, simple; eidos, form.] Having the number of chromosomes characteristic of mature germ-cells for the organism in question; appl. the typical gametic number of chromosomes after meiosis. n. Organism having one genome; cf. diploid.

haplomitosis (hāp'lömītō'sīs) n. [Gk. haploos, simple; mitos, thread.] Type of cell division where nuclear granules form chromospires which withdraw in two groups or divide transversely in the middle.

haplont (hap'lont) n. [Gk. haploos, simple; on, being.] An organism having haploid somatic nuclei.

haploperistomous (hăp'löperis'tömus) a. [Gk. haploos, simple; peri, around; stoma, mouth.] Having a peristome with a single row of teeth, appl. mosses.

haplopetalous (hăp'löpět'ălūs) a. [Gk. haploos, simple; petalon, petal.] With a single row of

petals.

haplophase (hāp'löfāz) n. [Gk. haploos, simple; phasis, aspect.] Stage in life-history of an organism when nuclei are haploid; gametophyte phase.

haploptile (hap'lötil, -ptil) n. [Gk. haploos, simple; ptilon, feather.] Single neossoptile, without rachis, formed by precocious development of the barbs of the teleoptile.

haplosis (hăplō'sĭs) n. [Ġk. haploos, simple.] Halving of the chromosome number during meiosis; reduction and disjunction.

haplostemonous (hăp'löstěm'önüs)

a. [Gk. haploos. simple; stemon,
warp.] Having one whorl of
stamens.

haplotype (hăp'lötīp) n. [Gk. haploos, simple; typos, pattern.]
The only species in a genus originally, and thereby becoming a genotype.

haplozygous (hăplŏz'igūs) a. [Gk. haploos, simple; zygon, yoke.] Appl. genes in haploid organisms;

hemizygous.

haptera (hap'tëră) n. plu. [Gk. haptein, to fasten.] Holdfasts, special disc-like outgrowths from the stem-like portion of certain algae, which serve as organs of attachment.

haptic (hăp'tik) a. [Gk. haptein, to touch.] Pert. touch; appl. stimuli and reactions.

haptogen (hăp'töjën) a. [Gk. haptein, to fasten; -genes, producing.] Appl. a limiting membrane of solidified protein which prevents miscibility.

haptomonad (hap'tömön'ad) n. [Gk. haptein, to fasten; monas, unit.] An attached form of certain parasitic Flagellata; cf. nectomonad. haptophores (hăp'töförz) n. plu. [Gk. haptein, to fasten; pherein, to carry.] The combining qualities of the molecule of a toxin, lysin, opsonin, precipitin, or agglutinin: cf. toxophores.

haptotropic (hăp'tötrŏp'ĭk) a. [Gk. haptein, to touch; trope, turn.]
Appl. curvature of a plant organ due to contact stimulus; thigmotropic.

haptotropism (hăptŏt'ropizm) n. [Gk. hapteın, to touch; trope, turn.] Response to contact stimulus, as in tentacles, tendrils, stems.

haptotype (hăp'totip) n. [Gk. haptein, to touch; typos, pattern.]
An icotype collected with the holotype but possibly taken from another plant.

Harderian gland [J. J. Harder, Swiss anatomist]. An accessory lacrimal gland of third eyelid or nictitating membrane.

harlequin lobe,—a testicular lobe with cells differing from those of other lobes, in certain Hemiptera.

harmonic suture,—an articulation formed by apposition of edges or surfaces, as between palatine bones.

harmosis (hârmō'sīs) n. [Gk. harmosis, fitting.] Arrangement and adaptation in response to a stimulus.

harmozone (hârmō'zōn) n. [Gk. harmozo, I arrange.] One of the hormones which influence growth and nutrition.

harpagones (hâr'păgō'nēz) n. plu. [L. harpago, hook.] Claspers or valves of certain male insects; a pair of sclerites between harpes and claspers in mosquitoes; harpes in Lepidoptera.

harpes (hâr'pēz) n. plu. [Gk. harpe, sickle.] Chitinous processes between the claspers of mosquitoes; claspers or valves of Lepidoptera.

Hassall's concentric corpuscles
[A. H. Hassall, English physician].
Epithelial cell nests in medulla of thymus.

hastate (hās'tāt) a. [L. hasta, spear.]
Spear-shaped, more or less triangular with the two basal lobes divergent; appl. leaf markings.

Hatschek's nephridium [B. Hatschek, Austrian zoologist]. A nephridium between notochord and preoral pit in Cephalochorda.

Hatschek's pit,—a mucin-secreting gland in roof of oral cavity in Cephalochorda; preoral pit.

haulm (hôm) n. [A.S. healm.] The stem of such plants as peas; the stem of a grass.

haustellate (hôs'tëlāt) a. [L. haurire, to suck.] Having a proboscis adapted for sucking.

haustellum (hôstěľúm) n. [L. haurire, to suck.] A proboscis adapted for sucking.

haustoria,—plu. of haustorium.

haustorial (hôstō'rĭăl) a. [L. haurire, to drink.] Pert. or resembling a haustorium.

haustorium (hôstô'riùm) n. [L. haurire, to drink.] An outgrowth of stem, root, or hyphae of certain parasitic plants, which serves to draw food from the host plant; sucker; an outgrowth of embryosac which extends to nutritive tissue in certain non-parasitic plants.

haustra (hôs'tră) n. plu. [L. haustor, drawer.] Recesses of sacculations of the colon, between plicae semilunares. Sing. haustrum.

Haversian canals [C. Havers, English anatomist]. Small canals in bone, in which lie bloodcapillaries, nerve, and lymph-space.

Haversian system,—a Haversian canal, the surrounding concentric lamellae and lacunae with canaliculi.

H-disc,—lighter region in anisotropic band of myofibrillae; Hensen's disc.

head-cap,—apical part or galea of head of spermatozoon; perforatorium.

head-case,—the outer hard covering of insect head.

head-cell,—one of the cells on manubrium of antheridium of Chara.

- head-kidney,—the pronephric portion of kidney, in vertebrates usually represented only in embryo; a nephridium usually developed in cephalic segment of invertebrates.
- heart (hârt) n. [A.S. heorte.] A hollow muscular organ with varying number of chambers which by rhythmic contraction keeps up circulation of blood; core or central portion of a tree or fruit.

heart-wood,—the darker, harder, central wood of trees; duramen.

heat (hēt) n. [A.S. haetu.] A kind of energy manifested in various ways; the sensation of warmth produced by stimulation of special organs; the period of sexual desire.

heat spot,—a special area on the skin at which nerve endings sensitive to

heat are found.

hectocotylus (hěk'tökŏt'ílŭs) n. [Gk. hekaton, hundred; kotylos, cup.]
One of the arms of a male cephalopod, specialised to effect transference of sperms.

hedonic (hēdŏn'ĭk) a. [Gk. hedone. pleasure.] Appl. skin glands of certain reptiles, which secrete musk and are specially active at rutting season.

heel (hēl) n. [A.S. hela.] Hinder or posterior tarsal portion of foot; talon or talonid of a tooth; a spinule at base of tibia in Hymenoptera.

hekistotherm (hē'kistothërm) n. [Gk. hekistos, least; therme, heat.] A plant that thrives with the minimum of heat, as alpine plants.

helices, -plu. of helix.

helicine (hel'isin) a. [Gk. helix, spiral.] Spiral; convoluted; hoisted; appl. certain convoluted and dilated arteries in penis; pert. outer rim of pinna.

helicoid (hĕl'ĭkoid) a. [Gk. helix, spiral; eidos, like.] Spiral; shaped like a snail's shell; pert. type of sympodial branching in which sympodium consists of fork branches of same side. helicoid cyme,—an inflorescence produced by suppression of successive axes on same side, thus causing the sympodium to be spirally twisted.

helicorubin (hěl'íköroob'ín) n. [L. helix, spiral; ruher, red.] A red pigment of gut of pulmonate

gastropods.

helicotrema (hěl'íkotrē'mă) n. [Gk helix, spiral; trema, hole.] A small opening near summit of cochlea by which the scalae vestibuli and tympani communicate.

heliophil (he'liofil) a. [Gk. helios, sun; philein, to love.] Adapted for relatively high intensity of light; heliophilic, heliophilous. Opp. heliophobic, skiophil.

heliophobic,—skiophil, q.v.

heliophyte (hē'lĭofīt) n. [Gk. helios, sun; phyton, plant.] A sun plant, opp. shade plant or skiaphyte.

heliosis (hē'liō'sis) n. [Gk. helios, sun.] Production of discoloured spots or markings on leaves through concentration of sun on them; solarisation.

heliotaxis (hē'liŏtāk'sis) n. [Gk. helios, sun; taxis, arrangement.] Locomotor or other response to stimulus of sunlight.

heliotropism (hē'liŏt'röpīzm) n. [Gk. helios, sun; trepein, to run.] Curvature of organisms or certain parts in response to the stimulus of sunlight.

helix (hē'liks) n. [Gk. helix, spiral.] A spiral; the coiled spiral arrangement of certain structures in invertebrates; the outer rim of external car.

helmet (hel'met) n. [O.F. helmet, head-covering.] The process of bill of hornbills; the bony plates covering head of certain extinct fishes; the galea of flowers and of insects.

helminthology (hěl'minthöl'öjí) n.
[Gk. helmins, worm; logos, discourse.] The study of the natural history of worms; the study of parasitic flatworms and round-

- helophyte (hěl'öfit) n. [Gk. helos, marsh; phyton, plant.] A marsh plant; a cryptophyte growing in soil saturated with water.
- helotism (hěl'ŏtizm) n. [Gk. heilotes, serf, from Helos, Laconian town.] Symbiosis in which the one organism enslaves the other and forces it to labour in its behalf, e.g. in lichens, in some species of ants.

hema-,—see haema-.

hemelytron (hěměl'ítrŏn) n. [Gk. hemi, half; elytron, sheath.]
Proximally hardened forewing of certain insects; elytron of certain worms; hemelytrum.

hemera (hēmēr'ā) n. [Gk. hemera, day.] The time during which fossiliferous strata constituting a zone of sedimentary rocks were deposited.

hemeranthous (hēmeran'thus) a. [Gk. hemera, day; anthos, flower.] Flowering by day; hemeranthic.

hemerophyte (hē'mërofīt) n. [Gk. hemeros, tame; phyton, plant.]
A cultivated plant.

hemibasidium (hěm'ībăsīd'īum) n. [Gk. hemi, half; basis, base; idion, dim.] The promycelium of the Ustilaginales.

hemibathybial (hĕm'ībăthīb'īāl) a. [Gk. hemi, half; bathys, deep; bios, life.] Pert. plankton between littoral and bathybial zones.

hemibranch (hěm'íbrángk) n. [Gk. hemi, half; brangchia, gills.] Gill with gill filaments on one side only; half-gill.

hemicellulase (hěm'ísěl'ūlās) n. [Gk. hemi, half; L. cellula, small cell.] An enzyme which effects hydrolysis of a hemicellulose, occurring in endosperm, fungi, and certain invertebrates.

hemicellulose (hĕm'isĕl'ūlōs) n. [Gk. hemi, half; L. cellula, small cell.]
One of several polysaccharides, chemically unrelated to cellulose, occurring as cell wall constituents in cotyledons, endosperms, and woody tissues, and serving as reserve food.

- hemicephalous (hěm'íkěf'álůs, -sěf-).

 a. [Gk. hemi, half; kephale, head.]

 Appl. insect larvae with reduced head.
- hemichlamydeous (hem'iklamid'eus)

 a. [Gk. hemi, half; chlamys, cloak.] Having ovuliferous scale inverted and bearing nucellus.
- hemichordate (hěm'íkôr'dāt) a. [Gk. hemi, half; chorde, string.] Possessing a rudimentary notochord.
- hemicryptophyte (hem'ikrip'tofit) n. [Gk. hemi, half; kryptos, hidden; phyton, plant.] A plant with dormant buds in the soil surface, the aerial shoots surviving for a season only.
- hemicyclic (hem'isi'klik) a. [Gk. hemi, half; kyklos, round.] With some floral whorls cyclic, others spiral.

hemielytron,—hemelytron, q.v.

hemiepiphyte (hěm'iěp'ífit) n. [Gk. hemi, half; epi, upon; phyton, plant.] A plant whose seeds germinate on another plant, but later send roots to the ground.

hemigamy (hemig'ami) n. [Gk. hemi-, half; gamos, marriage.]
Activation of ovum by male nucleus without nuclear fusion; semigamy.

hemignathous (hĕmīg'nāthūs) a. [Gk. hemi, half; gnathos, jaw.] Having one jaw shorter than the other, as in some birds.

hemikaryon (hĕm'ĭkăr'ĭŏn) n. [Gk. hemi, half; karyon, kernel.] A nucleus with gametic or haploid number of chromosomes; a pronucleus.

hemikaryotic (hěm'ikåriŏt'ik) a. [Gk. hemi, half; karyon, kernel.] Pert. hemikaryon; haploid.

hemimetabolic (hēm'imētābōl'ik) a. [Gk. hemi, half; metabole, change.] Having an incomplete or partial metamorphosis, as certain insects.

hemiparasitic (hēm'īpār'āsīt'īk) a. [Gk. hemi, half; para, beside; sitos, food.] Pert. a plant which is capable of carrying on photosynthesis, but not sufficiently to supply all food material.

hemipenis (hěm'īpē'nīs) n. [Gk. hemi, half; L. penis, penis.] One of the paired copulatory organs in lizards and snakes.

hemipneustic (hěm'īnū'stīk, -pnū-) a. [Gk. hemi, half; pnein, to breathe.] With one or more pairs of spiracles closed.

hemipterygoid (hěm'īptēr'īgoid) n. [Gk. hemi, half; pieryx, wing; eidos, form.] In neognath birds, part of pterygoid which fuses with palatine.

hemisaprophyte (hēm'īsāp'röfīt) n. [Gk. hemi, half; sapros, decayed; phyton, plant.] A plant living partly by photosynthesis, partly by obtaining food from humus.

hemisome (hem'isōm) n. [Gk. hemi, half; soma, body.] The symmetrical half of an animal about a median vertical plane.

hemisphere (hěm'īsfēr) n. [Gk. hemi, half; sphaira, globe.] One of the cerebral or cerebellar hemispheres.

hemisystole (hēm'īsīs'tölē) n. [Gk. hemi, half; syn, with; stellein, to set.] Contraction of one ventricle of the heart.

hemitropous (hemit'röpüs) a. [Gk. hemi, half; trope, turn.] Turned half round, having an ovule with hilum on one side and micropyle, etc., opposite in a plane parallel to placenta.

hemixis (hēm'īksīs) n. [Gk. hemi, half; mixis, mingling.] Fragmentation and reorganisation of macronucleus without involving micronucleus, in Paramecium.

hemizygous (hēmīz'īgūs) a. [Gk. hemi, half; zygon, yoke.] Appl. genes in haploid organisms; appl. genes without alleles in normal diploid organisms; haplozygous.

hemo-,-see haemo-.

Henle's layer [F. G. J. Henle, German anatomist]. Outermost stratum of nucleated cubical cells in inner root-sheath of a hair follicle.

Henle's loop,—loop of a kidney tubule within apical portion of pyramid.
Henle's sheath,—perineurium, or its prolongation surrounding branches of a nerve.

Hensen's line [V. Hensen, German histologist]. A disc dividing the darker portion of a sarcomere into two parts; mesophragma, Q line. Hensen's node,—the primitive node,

q.v.
Hensen's stripe,—a band of interlacing fibrils on under surface of tectorial membrane of Corti's organ.

hepar (hē'par) n. [Gk. hepar, liver.] Liver, or an organ having a similar function.

heparin (hē'părīn) n. [Gk. hepar, liver.] Substance present in liver and some other tissues, which inhibits formation, or action, of thrombin.

hepatic (hepat'ik) a. [Gk. hepar, liver.] Pert., like, or associated with the liver; pert. liverworts.

hepatocolic (hĕp'ātökŏl'ĭk) a. [Gk. hepar, liver; colon, large intestine.] Pert. liver and colon.

hepatocystic (hĕp'átösĭs'tīk) a. [Gk. hepar, liver; kystis, bladder.] Pert. liver and gall-bladder.

hepatoduodenal (hĕp'ātödū'ödē'nāl) a. [Gk. hepar, liver; L. duodeni, twelve each.] Pert. liver and duodenum.

hepatoenteric (hěp'ătöěntěr'ĭk) a. [Gk. hepar, liver; enteron, gut.] Of or pert. liver and intestine.

hepatogastric (hěp'ātögăs'trīk) a. [Gk. hepar, liver; gaster, stomach.] Pert. liver and stomach.

hepatopancreas,—digestive gland in many invertebrates, supposed to perform a function similar to that of liver and of pancreas in higher forms.

hepatoportal (hěp'átöpôr'tăl) a. [Gk. hepar, liver; L. porta, gate.]
Pert. or designating portal circulation of liver.

hepatorenal (hěp'átörē'năl) a. [Gk. hepar, liver; L. renes, kidneys.]
Pert. liver and kidney.

hepatoumbilical (hep'atoumbil'ikal)
a. [Gk. hepar, liver; L. umbilicus,
navel.] Joining liver and umbili-

- heptagynous (hěptăj'înŭs) a. [Gk. hepta, seven; gyne, female.] With seven pistils.
- heptamerous (hěptăm'erŭs) a. [Gk. hepta, seven; meros, part.] Having whorls of flowers in sevens.
- heptandrous (hěptăn'drus) a. [Gk. hepta, seven; aner, man.] Having seven stamens.
- heptarch (hĕp'târk) a. [Gk. hepta, seven; arche, beginning.] Having seven initial groups of xylem.
- heptastichous (hěptás'tikůs) a. [Gk. hepta, seven; stichos, row.] Arranged in seven rows; appl. leaves. herb (hěrb) n. [M.E. herbe.] A

herb (herb) n. [M.E. herbe.] A seed plant without woody stem.

- herbaceous (hěrbā'shŭs) a. [L. herbaceus, grassy.] Pert. or resembling a herb, or similarly formed.
- Herbst's corpuscle [E. F. Herbst, German anatomist]. A simple type of Pacinian corpuscle, in birds.
- hercogamy (hěrkog'ámi) n. [Gk. herkos, barrier; gamos, union.] The condition in which self-fertilisation is impossible; also herkogamy.
- hereditary (hered'itari) a. [L. hereditas, heirship.] Transmissible from parent to offspring, as characteristics, physical or mental.

heredity (hërëd'iti) n. [L. hereditas, heirship.] The organic relation between successive generations; germinal constitution.

hermaphrodite (hermaf'rödit) n. [Gk. hermaphroditos, combining both sexes.] An organism with both male and female reproductive organs. a. Hermaphroditic.

hermaphroditism (hermaf'roditizm)

n. [Gk. hermaphroditos, combining both sexes.] The condition of having both male and female reproductive organs in one individual.

herpetology (her'petol'öji) n. [Gk. herpeton, reptile; logos, discourse.]
That part of zoology dealing with the structure, habits and classification of reptiles.

hesperidium (hes'perid'ium) n. [Gk. Hesperides, sisters guarding the golden apples given by Gaea.] A

superior, many-celled, few-seeded indehiscent fruit, having epicarp and mesocarp joined together, and endocarp projecting into interior as membranous partitions which divide the pulp into chambers. *E.g.* orange.

heteracanthous (hět'ërăkăn'thůs) a. [Gk. heteros, other; akantha, spine.] Having the spines in dorsal fin asymmetrical.

heteractinal (hēt'ērāk'tīnāl) a. [Gk. heteros, other; aktis, ray.] Pert. nail-like spicules having disc of six to eight rays in one plane, and a stout ray at right angles to these.

heterandrous (hět'ërăn'drŭs) a. [Gk. heteros, other; aner, man.] With stamens of different length or shape.

heterauxesis (het'erôksē'sis) n. [Gk. heteros, other; auxesis, growth.] Irregular or asymmetrical growth of organs; relative growth rate of parts of an organism; heterogonic or allometric growth; bradyauxesis and tachyauxesis, q.v.

heterauxin,—heteroauxin, q.v.

heteraxial (hět'erāk'sĭăl) a. [Gk. heteros, other; axis, axis.] With three unequal axes.

heteroagglutinin (het'eröägloot'inin)

n. [Gk. heteros, other; L. agglutinare, to glue to.] Fertilisin or agglutinin of eggs which reacts on sperm of different species; cf. isoagglutinin.

heteroauxin (hět'erôôk'sĭn) n. [Gk. heteros, other; auxein, to grow.]
A growth-promoting hormone, extracted from fungi; heterauxin; β-indolyl-acetic acid, C₁₀H₉O₂N.

heteroblastic (hět'ëröblăs'tĭk) a. [Gk. heteros, other; blastos, bud.] With indirect development.

- heterobrachial (hět'ëröbrā'kiăl) a. [Gk. heteros, other; L. brachium, arm.] Pert. chromosome arms on either side of centromere; pericentric.
- heterocarpous (hět'ërökâr'pŭs) a. [Gk. heteros, other; karpos, fruit.] Bearing two distinct types of fruit.

- heterocellular (hět'ërösěl'ūlăr) a. [Gk. heteros, other; L. cellula, small cell. | Composed of cells of more than one type. Opp. homocellular.
- heterocephalous (hět'ërökěf'ālūs. [Gk. heteros, other: -sĕf-) a. kephale, head.] Having pistillate flowers on separate heads from staminate.
- heterocercal (hět'ërosěr'kăl) a. [Gk. heteros, other; kerkos, tail.] Having vertebral column terminating in upper lobe of caudal fin, which is usually larger than lower.

heterocercy (hět'érösěr'sĭ) n. heteros, other; kerkos, tail.] Condition of having a heterocercal tail.

- heterochlamydeous (hět'eroklamid'-[Gk. heteros, other: čŭs) a. Having a calyx chlamys, cloak.] differing from corolla in colour, texture, etc.
- heterochromatic (hěť erokromať ik) [Gk. heteros, other; chroma, colour.] Pert. heterochromatin; appl. chromosomal regions liable to become heteropycnotic. euchromatic.
- heterochromatin (hět'ërökrō'mătĭn) n. [Gk. heteros, other; chroma, colour]. Chromatin retaining a high nucleic acid content and regulating nucleic acid metabolism in nucleus and cytoplasm.
- heterochromatism (hěť erokro mătĭzm) n. [Gk. heteros, other; chroma, colour. Change of colour, as seasonal colour change in an inflorescence.

heterochromaty,-differential stain-

heterochromia (hěť erökrô myň) n. [Gk. heteros, other; chroma, colour.] Difference in colour of parts normally of one colour, as of irides of a pair of eves.

heterochromosome (hět'ërokrō'mö-[Gk. heteros, other; n. chroma, colour; soma, body.] A chromosome other than an ordinary or typical one; sex-chromosome; allosome. Opp. autosome, euchro-

mosome.

- heterochromous (hět'ërökrō'mŭs) a. [Gk. heteros, other; chroma, colour.] Having disc and marginal florets differently coloured.
- heterochronism (hět'erők'rönizm) n. [Gk. heteros, other; chronos, time.] Departure from typical sequence in time of formation of organs; heterochrony.
- heterochrosis (hět'ërökrō'sĭs) n. [Gk. heteros, other; chrosis, colouring.] Abnormal coloration.
- heteroclinous (hět'ëröklī'nŭs) a. [Gk. heteros, other; kline, bed.] Heterocephalous, q.v.
- heterocoelous (hět'érösē'lŭs) a. [Gk. heteros, other; koilos, hollow.] Pert. vertebrae with saddle-shaped articulatory centra; concavoconvex.
- heterocysts (hět'érösists) n. plu. [Gk. heteros, other; kystis, bladder.] Clear cells occurring at intervals filaments of certain algae, marking limits of hormogonia.

heterodactylous (hět'erödăk'tilus) a. [Gk. heteros, other; daktylos, digit.] With the first and second toes turned backwards.

heterodont (hět'érodont) a. [Gk. heteros, other; odous, tooth.] Having the teeth differentiated for various purposes. Opp. homodont.

- heterodromous (hět'ërŏd'römŭs) a. [Gk. heteros, other; dramein, to Having genetic spiral of stem leaves turning in different direction to that of branch leaves.
- heteroecious (hět'ërē'sĭŭs) a. [Gk. heteros, other; oikos, house.] Passing different stages of life history in different hosts; exhibiting heteroecism; metoecious; metoxenous.
- heterogamete (hět'ërögămēt') [Gk. heteros, other; gametes. spouse.] One of dissimilar conjugating gametes; an anisogamete.
- heterogametic (hět'ërögămět'ik) a. heteros, other; gametes, spouse.] Elaborating two kinds of gametes in equal numbers; having unequal pair of sex chromosomes, XY or WZ; appl. sex that is heterozygous; reproducing sexually; digametic; cf. homogametic.

heterogamous (hět'ërög'amus) a. [Gk. heteros, other; gamos, mar-With unlike gametes; having two types of flowers; having indirect pollination methods.

heterogamy (hět'erog'ami) n. [Gk. heteros, other; gamos, offspring.] Alternation of two sexual generations, one being true sexual, the other parthenogenetic; condition of having, or union of, gametes of different size and structure; anisogamy.

heterogangliate (het'erogang'gliat) a. [Gk. heteros, other; gangglion, ganglion.] With widely separated and asymmetrically placed nerve-ganglia.

heterogenesis (hět'ëröjěn'esis) n. heteros, other; ſGk. descent.] Spontaneous generation;

alternation of generations.

heterogenetic (hěť eröjeněť ik) a. other; [Gk. heteros, genesis, descent.] Descended from different ancestral stock; pert. heterogenesis.

heterogeny (hět'ëroj'ëni) n. heteros, other; genos, generation.] Having several distinct generations succeeding one another in a regular series.

heterogonic (hět'ërögŏn'ík) a. [Gk. heteros, other; gonos, produce.] Differing in developmental growth rate; allometric.

heterogonous (hĕt'ërŏg'önŭs) [Gk. heteros, other; gonos, birth.] *Pert.* heterogenesis, or heterogony.

heterogony (hět'ërog'oni) n. [Gk. heteros, other; gonos, birth.] Condition of having two, or three, kinds of flowers differing in length of stamen; alternation of generations; allometry, q.v.

heterogynous (hět'ërŏj'inŭs) a. [Gk. heteros, other; gyne, woman.] With

two types of females.

heterokaryon (hĕt'ërökår'ĭŏn) IGk. heteros, other; karyon, nucleus. | An individual having heterokaryotic cells.

heterokaryosis (hět'erökăriō'sis) n. [Gk. heteros, other; karyon, nucleus.] Presence of genetically dissimilar nuclei within individual cells: heterokaryotic condition.

heterokaryote (hět'érökår'iōt) a. [Gk. heteros, other; karyon, nucleus.] Having two distinct types of nuclei.

heterokaryotic (hěť erökariŏt'ik) a. [Gk. heteros, other; karyon, nucleus.] Having genetically dissimilar nuclei, in a multinucleate cell, or in different cells of a hypha; heterokaryote.

heterokinesis (hĕt'ërökinē'sis) [Gk. heteros, other; kinein, to move.] Qualitative or differential division of chromosomes.

heterolecithal (het'eroles'ithal) a. [Gk. heteros, other; lekithos, volk.] Having unequally distributed deuto-

plasm.

heterologous (hět'ërŏl'ögŭs) a. [Gk. heteros, other; logos, relation.] Of different origin; derived from a different species; appl. various substances, e.g. agglutinins, affecting other than species of origin: cf. homologous.

heterology (hět'ërŏl'ojĭ) n. heteros, other; logos, relation.] Noncorrespondence of parts owing to different origin or different elements.

heteromallous (hět'ëromal'us)a. [Gk. heteros, other; mallos, lock of wool.] Spreading in different directions.

heteromastigate (hět'ërömăs'tigāt) [Gk. heteros, other; mastix, lash.] Having two different types of flagella; heteromastigote.

heteromerous (hět'ërŏm'ërŭs) [Gk. heteros, other; meros, part.] Having, or consisting of, an unequal number of parts, appl. whorls, tarsi, etc.; having a stratified thallus.

heterometabolic (hět'ërömětábol'ík) [Gk. heteros, other; metabole, change.] Having incomplete metamorphosis.

heteromorphic (hět'ërömôr'fik) a. [Gk. heteros, other; morphe, shape.] Having different forms at different chromosomes times: appl. different size and shape, chromosome pairs differing in size; appl. alternation of diploid and haploid phases in morphologically dissimilar generations, antithetic.

heteromorphism (het'eromôr'fizm) n. [Gk. heteros, other; morphe. shape.] The state or quality of

being heteromorphic.

heteromorphosis (hět'ërömôr'fōsis) n. [Gk. heteros, other; morphosis, shaping.] Production of a part in an abnormal position; regeneration, when the new part is different from that removed; cf. homoeosis.

heteromorphous (hět'ërömôr'fus) a. [Gk. heteros, other; morphe, shape.] Pert. an irregular structure, or departure from the normal.

heteronereis (hēt'erönē'rēis) n. [Gk. heteros, other; Nereis, Nereid.] A free-swimming dimorphic sexual stage of Nereis and other marine worms.

heteronomous (hět'ërŏn'ömus) a. [Gk. heteros, other; nómos, law.] Subject to different laws of growth; specialised on different lines. [Gk. heteros, other; nomós, department.] Appl. segmentation into dissimilar segments.

heteropelmous (hět'ëröpěl'můs) a. [Gk. heteros, other; pelma, sole of foot.] Having flexor tendons of toes bifid.

heteropetalous (hět'ëröpět'ălŭs) a. [Gk. heteros, other; petalon, petal.] With dissimilar petals.

heterophagous (hĕt'ërŏf'ăgŭs) [Gk. heteros, other; phagein, to Having young in altrices eat.] condition.

heterophil (hět'ëröfĭl) [Gk. heteros, other; philos, loving.] Appl. non-specific antigens and antibodies present in an organism, affording natural immunity; appl. granular leucocytes which show interspecific differences in their reaction to stains. n. Polymorphonuclear leucocyte.

(hĕt'ëröfīăd'ik) heterophyadic [Gk. heteros, other; phyas, shoot.] Producing separate shoots, vegetative, one reproductive.

heterophyllous (het'eröfil'us) a. [Gk. heteros, other; phyllon, leaf.] Bearing foliage leaves of different shape on different parts of the same plant.

heterophytic (hět'ëröff'tik) a. [Gk. heteros, other; phyton, plant.] With two kinds of spores, borne by different sporophytes; cf. homophytic.

heteroplasia (hět'éröplā'sĭă) n. [Gk. heteros, other; plassein, to mould.] The development of one tissue from another of a different kind.

heteroplasm (hět'eroplazm) n. [Gk. heteros, other; plasma, mould.] Tissue formed in abnormal places.

heteroplasma (hět'ëröplăzmă) n. [Gk. heteros, other; plasma, mould.] Plasma from a different species used as a medium for tissue culture; cf. autoplasma, homoplasma.

heteroplastic (hět'ëröplăs'tik)a. [Gk. heteros, other; plastos, formed.] Appl. grafts of unrelated material; appl. transplantation between individuals of different species or genera, opp. homoioplastic; cf.

xenoplastic.

heteroploid (hět'ëröploid) a. [Gk. heteros, other; haploos, onefold.] Having an extra chromosome through non-disjunction of a pair in meiosis; not having a multiple of the basic haploid number of chromosomes. n. An organism having heteroploid nuclei.

heteroproteose (hět'ëröprō'tëōs) n. [Gk. heteros, other; protos, first.] One of primary products formed by action of gastric juices on other hydrolysing agents on proteins;

propeptone.

heteropycnosis (hět'ëröpiknō'sis) n. [Gk. heteros, other; pyknos, dense.] Condensation of sex-chromosome during growth - period stages of gonia and cytes; condition of chromosome region or of chromosomes synthesising more or less nucleic acid than remainder of chromosome set.

heteropycnotic (hěť eröpíknoť ík) a. [Gk. heteros, other; pyknos, dense.] chromosome manifesting Appl.

heteropycnosis.

heterorhizal (hět'ërörī'zăl) a. [Gk. heteros, other; rhiza, root.] With roots coming from no determinate point.

- heterosexual (hět'ërösčk'sūăl) a. [Gk. heteros, other; L. sexus. sex.] Of, or pert. the opposite sex; appl. hormones, etc.
- heterosis (hēt'erō'sīs) n. [Gk. heteros, other.] Cross-fertilisation; hybrid vigour, result of heterozygosis
- heterosomal (hět'ërösō'măl) a. [Gk. heteros, other; soma, body.] Occurring in, or pert., different bodies; appl. rearrangements in two or more chromosomes of a set.
- heterosome (hěť'ērösōm) n. [Gk. heteros, other; soma, body.] A heterochromosome.
- heterosporangic (hěť erösporan'jík)

 a. [Gk. heteros, other; sporos, seed; anggeion, vessel.] Bearing two kinds of spores in separate sporangia.
- heterosporous (hět'ërŏs'pörŭs, hětëröspō'rūs) a. [Gk. heteros, other; sporos, seed.] Producing two kinds of spores; heterosporic.
- heterospory (hět'ërŏs'pöri) n. [Gk. heteros, other; sporos, seed.] The condition of being heterosporous; the production of megaspores and microspores.
- heterostemonous (hět'éröstěm'önűs)

 a. [Gk. heteros, other; stemon, stamen.] With unlike stamens.
- heterostrophy (hět'ërŏs'tröfi) n. [Gk. heteros, other; strophe, turning.]
 The condition of being coiled in a direction opposite to normal.
- heterostyled (het'erösti'ld) a. [Gk. heteros, other; stylos, pillar.] Having unlike or unequal styles; heterostylic.
- heterostyly (hět'ëröstî'lĭ) n. [Gk. heteros, other; stylos, pillar.] Condition of being heterostyled.
- heterosynapsis (hět'ërösĭnăp'sīs) n. [Gk. heteros, other; synapsis, union.] Pairing of two dissimilar chromosomes; cf. homosynapsis.
- heterotaxis (hēt'erötāk'sīs) n. [Gk. heteros, other; taxis, arrangement.]
 Abnormal or unusual arrangement of organs or parts.
- heterothallic (het'erothal'ik) a. [Gk. heteros, other; thallos, young

- shoot.] Requiring branches of two distinct mycelia to form a zygospore; appl. moulds; cf. homothallic.
- heterotic (hět'érŏt'īk) a. [Gk. heteros, other.] Pert. heterosis; appl. vigour.
- heterotomy (hět'erŏt'ömĭ) n. [Gk. heteros, other; temnein, to cut.] Condition of having parts of perianth whorls unequal or dissimilar; irregular dichotomy in Crinoidea.
- heterotopy (hět'ërŏt'öpĭ) n. [Gk. heteros, other; topos, place.] Displacement; abnormal habitat.
- heterotrichous (hět'ērŏt'rĭkŭs) a. [Gk. heteros, other; thrix, hair.] Having two types of cilia; having thallus consisting of prostrate and erect filaments, as certain algae.
- heterotrophic (het'erötröf'ik) a. [Gk. heteros, other; trophe, nourishment.] Getting nourishment from organic substances; appl. parasitic plants; opp. autotrophic.
- heterotropic chromosome, sexchromosome, q.v.
- heterotropous (hět'erŏt'röpŭs) a. [Gk. heteros, other; trepein, to turn.] Pert. ovule with hilum and micropyle at opposite ends in a plane parallel to placenta; heterotropic.
- heterotype (het'erötip) n. [Gk. heteros, other; typos, pattern.]
 First meiotic division.
- heterotypic (hěť erötíp'ik) a. [Gk. heteros, other; typos, pattern.]
 Pert. mitotic division in which daughter chromosomes remain united and form rings; appl. first or reduction division in meiosis; cf. homeotypic.
- heterotypical (hět'ërötĭp'ĭkăl) a. [Gk. heteros, other; typos, pattern.] Appl. a genus comprising species that are not truly related.
- heteroxenous (hět'ërözěn'ŭs) a. [Gk. heteros, other; xenos, host.] Occurring on or infesting more than one kind of host; heteroecious.
- heterozygosis (hět'ërözigō'sis) n. [Gk. heteros, other; zygon, yoke.] Descent from two different species, varieties, or races.

heterozygote (hět'ërozī'gōt) n. [Gk. heteros, other; zygon, yoke.] An organism having alternative forms of a gene; an impure dominant; a heterozygous organism.

heterozygous (hětěroz'īgŭs) a. [Gk. heteros, other; zygon, yoke.] Bearing two dissimilar alternative genetical features.

ical factors.

hexacanth (hěk'săkănth) a. [Gk. hex, six; akantha, thorn.] Having six hooks; appl. embryo of certain flat-worms.

hexactinal (hěk'sāk'tīnāl) a. [Gk. hex, six; aktis, ray.] With six rays. hexactine (hěk'sāk'tīn) n. [Gk. hex, six; aktis, ray.] A spicule with six equal and similar rays meeting at right angles.

hexactinian (hěk'säktin'iăn) a. [Gk. hex, six; aktis, ray.] With tentacles or mesenteries in multiples of

six.

hexacyclic (hěk'săsī'klĭk) a. [Gk. hex, six; kyklos, circle.] Having floral whorls consisting of six parts. hexagynous (hěksăj'ĭnŭs) a. [Gk. hex, six; gyne, woman.] Having six pistils.

hexamerous (hěksăm'crūs) a. [Gk. hex, six; meros, part.] Occurring in sixes, or arranged in sixes.

hexandrous (hěksān'drůs) a. [Gk hex, six; aner, man] Having six stamens.

hexapetaloid (hěk'săpět'áloid) a. [Gk. hex, six; petalon, petal; eidos. form.] With petaloid perianth of six parts.

hexapetalous (hěk'săpět'ălŭs) a. [Gk. hex, six; petalon, petal.] Having six petals.

hexaphyllous (hěk'săfil'üs) a. [Gk. hex, six; phyllon, leaf.] Having six leaves.

hexaploid (hěk'săploid) a. [Gk. hex, six; haploos, simple; eidos, form.] With six sets of chromosomes. n. An organism having six times the monoploid chromosome number.

hexapod (hěk'săpŏd) a. [Gk. hex, six; pous, foot.] Pert. animal with six legs. n. An insect.

hexapterous (hěksăp'terūs) a. [Gk. hea, six; pteron, wing.] Having six wing-like processes or expansions.

hexarch (hěk'sârk) a. [Gk. hex, six; arche, beginning.] Having six radiating vascular strands; appl. roots.

hexasepalous (hěk'sāsĕp'álŭs) a. [Gk. hex, six; F. sépale, sepal.] Having six sepals.

hexaspermous (hěk'săspěr'mŭs) a. [Gk. hex, six; sperma, seed.] Having six seeds.

hexastemonous (hēk'săstěm'onŭs) a. [Gk. hex, six; stemon, stamen.] Having six stamens; hexandrous.

hexaster (heksas'ter) n. [Gk. hex, six; aster, star.] A variety of hexactine in which the rays branch and produce star-shaped figures.

hexastichous (hěksǎs'tǐkǔs) a. [Gk. hex, six; stichos, row.] Having the parts arranged in six rows.

hexicology (hěk'sĭkŏl'ojĭ) n. [Gk. hexis, habit; logos, discourse.] Bionomics.

hexuronic acid,—ascorbic acid or vitamin C.

hiatus (hīā'tŭs) n. [L. hiare, to gape.] Any large opening.

hibernaculum (hī'bernāk'ūlūm) n.
[L. hibernaculum, winter-quarters.]
A winter bud; specially modified winter bud in fresh-water Polyzoa.

hibernate (hī'bërnāt) v. [L. hibernus, wintry.] To pass the winter in a resting state.

hibernating glands, -lymph glands of richly vascularised fatty tissue occurring in some rodents and insectivores.

hidrosis (hidro'sis) n. [Gk. hidros, sweat.] Excretion of sweat; perspiration.

hiemal (hī'ēmăl) a. [L. hiems, winter.] Pert. winter; appl. aspect of a community.

Highmore's antrum [N. Highmore, English surgeon]. The maxillary sinus, which communicates with the middle meatus of the nose.

Highmore's body,—corpus highmoreanum, mediastinum testis.

- hiliferous (hilif'ërus) a. [L. hılum, trifle; ferre, to carry.] Having a hilum.
- hilum (hī'lum) n. [L. hilum, trifle.]
 Scar on ovule where it was attached to placenta; eye of seed; nucleus of starch grain; small notch, opening, or depression, usually where vessels, nerves, etc., enter, of kidney, lung, spleen, etc.; also hilus.
- hind brain,—rhombencephalon, that portion of brain derived from third embryonic vesicle.
- hind gut,—diverticulum of yolk-sac extending into tail-fold in human embryo; posterior portion of alimentary tract.
- hinge-cells,—large epidermal cells which, by changes in turgor, control rolling and unrolling of a leaf.
- hinge-joint,—a joint in which articulatory surfaces are so moulded as to permit motion in one plane only; ginglymus.
- hinge-ligament,—the tough elastic substance joining the two valves of a bivalve shell.
- hinge-line,—the line of articulation of the two valves in a bivalve shell.
- hinge-tooth,—one of the projections found on the hinge-line in bivalves.
- hip-joint,—the ball-and-socket joint between femur and hip-girdle.
- hippocampal (hip'ökam'pal) a. [Gk. hippos, horse; kampe, bend.] Pert. the hippocampus.
- hippocampus (hip'ökäm'pus) n. [Gk. hippos, horse; kampe, bend.] Part of rhinencephalon forming an eminence extending throughout length of floor of inferior cornu of lateral ventricle.
- hippuric (hipū'rik) a. [Gk. hippos, horse; ouron, urine.] Obtained from horse's urine; appl. acid synthesised by kidney and present in urine of herbivorous animals.
- hirsute (hīrsūt') a. [L. hirsutus, hairy.] Covered with hair-like feathers, appl. birds; having stiff, hairy bristles or covering.

- hirudin (hǐrū'din) n. [L. hirudo, leech] A substance, obtained in solution from buccal glands of leech, which prevents clotting of blood by inhibiting action of thrombin on fibrinogen.
- His' bundle | W. His, German anatomist]. Band of muscle fibres, with nerve fibres, connecting auricles and ventricles of heart; atrioventricular or auriculoventricular bundle; Gaskell's bridge.
- hispid (his'pid) a. [L. hispidus, rough.] Having stiff hairs, spines, or bristles.
- histamine (his'tamin) n. [Gk. histos, tissue; ammoniakon, resinous gum.] Product of the basic amino acid and food constituent histidine, in ergot and animal tissues, stimulates autonomic nervous system, gastric juice secretion, and capillary dilatation; C₅H₉N₃.
- histiocyte (his'tiösīt) n. [Gk. histion, tissue; kytos, hollow.] A primitive blood cell giving rise to a monocyte; a monocyte of reticular origin, or a clasmatocyte derived from endothelium, a reticulo-endothelial cell; fixed macrophage in loose connective tissue; adventitial cell; rhagiocrine cell.
- histiomonocyte (his'tiömŏn'ösīt) n.
 [Gk. histion, tissue; monos, alone; kytos, hollow.] An endothelial cell of certain capillaries and associated with the histiocytic metabolic system.
- histiotypic (hīs'tiötīp'īk) a. [Gk. histion, tissue; typos, pattern.] Appl. uncontrolled or unorganised growth of cells, in tissue culture. Opp. organotypic.
- histoblast (his töbläst) n. [Gk. histos, tissue; blastos, bud.] A unit of tissue; imaginal disc.
- histochemistry (his'tökem'istri) n. [Gk. histos, tissue; chemeia, transmutation.] The chemistry of animal tissues.
- histocyte (hīs'tösīt) n. [Gk. histos, tissue; kytos, hollow.] Tissue cell as distinguished from germ cell.

histogenesis (hĭs'töjĕn'ësĭs) n. [Gk. histos, tissue; genesis, descent.] Formation and development of tissue.

histogenic (hĭs'töjĕn'īk) a. histos, tissue; -genes, producing.] Tissue-producing; appl. the separate merismatic layers in a stratified growing point.

histogens (his'töjenz) n. plu. [Gk. histos, tissue; gennaein, to produce.] Tissue-producing zones or layers: plerome, periblem, derma-

togen, and calyptrogen.

histohaematin (his'töhēm'ātin) n. [Gk. histos, tissue; haima, blood.] An intracellular haemin compound; a cytochrome.

histology (histol'oji) n. [Gk. histos, logos, discourse.] tissue; science which treats of the detailed structure of animal or plant tissues; microscopic morphology; morphology.

histolysis (histol'isis) n. [Gk. histos, tissue; lyein, to dissolve.] The dissolution of organic tissues; process by which most of pupal internal organs dissolve into creamy fluid, except certain cells round which new imaginal tissues are formed.

histone (his'ton) n. [Gk. histos, tissue.] A protein constituent of cell nuclei, thymus, blood corpuscles, and lymph glands.

histophyly (histöfī'li) n. [Gk. histos, tissue; phyle, tribe.] Phylogenetic history of a group of cells.

histoteleosis (hĭs'tötělē'ōsĭs) n. [Gk. histos, tissue; teleios, full-grown.] The completion of functional differentiation of tissue cells.

histotrophic (hĭs'tötrŏf'ĭk) a. [Gk. histos, tissue; trephein, to nourish.] *Pert.* or connected with tissue formation or repair.

histozoic (his'tözō'ik) a. [Gk. histos, tissue; zoon, animal.] Living within tissues; appl. trophozoitic stage of certain Sporozoa.

histozyme (his'tözīm) n. [Gk. histos, tissue; zyme, leaven.] An enzyme found in kidneys of certain animals, capable of decomposing hippuric acid.

hock (hok) n. [A.S. hoh, heel.] The tarsal joint, or its region; hough.

holandric (hŏlăn'drĭk) a. [Gk. holos, whole; aner, male.] Pert. holandry; transmitted from male to male through the Y-chromosome; appl. sex-linked characters.

holandry (hŏlăn'drĭ) n. [Gk. holos, whole; aner, male.] The condition of having full number of testes, as two pairs in Oligochaeta; cf. meroandry.

holarctic (hŏlârk'tĭk) a. [Gk. holos, whole; Arktos, Great Bear.] Appl. or pert. a zoogeographical region including northern parts of the Old and New Worlds or palaearctic and nearctic sub-regions, q.v.

holard (hölârd') n. [Gk. whole; ardo, I water.] Total water content of soil; cf. chresard, echard.

holaspidean (hŏl'aspid'ean) a. [Gk. holos, whole; aspis, shield.] With single series of large scales on posterior aspect of tarso-metatarsus.

holcodont (hŏl'ködŏnt) a. holkos, furrow; odous, tooth.] Having the teeth in a long continuous groove.

holdfast,--the sucker or disc on the thallus of certain algae; hapteron. holobasidium (hŏl'öbăsĭd'ĭŭm) n. [Gk. holos, whole; basis, base; A basidium not idion, dim.] divided by septa.

holobenthic (hŏl'obĕn'thĭk) a. [Gk. holos, whole; benthos, depths.] Living in depths of sea throughout life.

holoblastic (hŏl'öblăs'tĭk) a. holos, whole; blastos, bud.] Pert. eggs with total cleavage.

holobranch (hol'obrangk) n. holos, whole; brangchia, gills.] A gill in which gill filaments are borne on both sides.

holocarpic (hŏl'ökâr'pĭk) a. [Gk. holos, whole; karpos, fruit.] Appl, fungi without rhizoids or haustoria. living in host cell, as certain Phycomycetes; cf. eucarpic.

Holocene (hŏl'ösēn) a. [Gk. holos, whole; kainos, recent.] Recent geological epoch following Pleistocene; postglacial age.

holochlamydate (hŏl'öklām'idāt) a. [Gk. holos, whole; chlamys, cloak.] Having no notch on mantle margin.

holochroal (hŏl'ökrō'āl) a. [Gk. holos, whole; chros, close.] Having eyes with globular or biconvex lenses closely crowded together, so that cornea is continuous over whole eye.

holocrine (hŏl'ökrin) a. [Gk. holos, whole; krinein. to separate.]

Appl. glands in which secretory cells disintegrate and form part of secretion, as sebaceous glands;

cf. apocrine, merocrine.

holocyclic (hŏl'ösšk'lĭk) a. [Gk. holos, whole; kyklos, circle.] Pert. or completing alternation of sexual and parthenogenetic generations.

hologametes (hŏl'ögamēts') n. plu. [Gk. holos, whole; gametes, spouse.] Fully developed protozoa taking part in syngamy. Opp. merogametes.

hologamy (hölög'āmĭ) n. [Gk. holos, whole; gamos, marriage.] Macrogamy; condition of having gametes similar to somatic cells.

hologastrula (hŏl'ögas'troolă) n.
[Gk. holos, whole; gaster, stomach.]
Gastrula formed from holoblastic egg.

holognathous (hŏlŏg'năthŭs) a. [Gk. holos, whole; gnathos, jaw.] Having the jaw in a single piece.

hologynic (hölöjĭn'ik) a. [Gk. holos, whole; gyne, woman.] Transmitted direct from female to female; appl. sex-linked characters.

holomastigote (höl'ömäs'tigōt) a. [Gk. holos, whole; mastix, whip.] Having one type of flagellum scattered evenly over the body.

holometabolic (höl'ömětábol'ik) a. [Gk. holos, whole; metabole, change.] Having complete metamorphosis.

holometabolism (hŏl'ömětăb'ölizm)

n. [Gk. holos, whole; metabole,
change.] State of having complete

metamorphosis. Opp. hemimetabolism.

holomorphosis (hŏl'ömôr'fōsĭs) n. [Gk. holos, whole; morphe, shape.] Regeneration in which the entire part is replaced.

holonephridia,—meganephridia, q.v. holonephros (hŏl'önĕf'rŏs) n. [Gk. holos, whole; nephros, kidney.] The hypothetical continuous excre-

tory organ.

holophytic (höl'öfit'ik) a. [Gk. holos, whole; phyton, plant.] Obtaining the whole of its food after the manner of a plant; phototrophic. Opp. holozoic.

holoplankton (höl'öplängktön) n. [Gk. holos, whole; plangktos, wandering.] The marine or fresh-water organisms which complete their life cycle while drifting with the surrounding water.

holoplanktonic (höl'öplängktön'îk)

a. [Gk. holos, whole; plangktos,
wandering.] Living near the surface of sea, or of lake, throughout
life; pert. holoplankton.

holopneustic (hŏl'önū'stĭk) a. [Gk. holos, whole; pnein, to breathe.] With all spiracles open for respiration.

holoptic (hŏlŏp'tĭk) a. [Gk. holos, whole; ops, eye.] Having eyes of two sides meeting in a coadapted line of union.

holorhinal (hŏl'örī'năl) a. [Gk. holos, whole; rhis, nose.] Having nares with posterior margin rounded. Opp. schizorhinal.

holoschisis (hŏlŏs'kisis) n. [Gk. holos, whole; schizein, to cut.] Amitosis.

holosericeous (hŏl'ösĕrīsh'ūs) a. [Gk. holos, whole; L.L. sericeus, silken.] Completely covered with silky hairlike structures; having a silky lustre or sheen.

holostomatous (hŏl'östŏm'ătŭs) a. [Gk. holos, whole; stoma, mouth.] With margin of aperture entire.

holostyly (hol'östi'li) n. [Gk. holos, whole; stylos, pillar.] Primitive condition of jaw suspension in some fishes.

- holosystolic (hŏl'ösĭstŏl'ĭk) a. [Gk. holos, whole; systole, contraction.] Pert. complete systole.
- holotrichous (hŏlŏt'rĭkŭs) a. holos, whole; thrix, hair.] Having a uniform covering of cilia over the
- holotype (hŏl' \ddot{o} t \ddot{i} p) n. [Gk. holos, whole; typos, pattern.] The single specimen chosen for designation of a new species.
- holozoic (hŏl'özō'ik) a. [Gk. holos, whole; zoon, animal.] Obtaining the whole of its food after the manner of animals; ingulfing proteids. Opp. holophytic.

homacanth (hom'akanth) a. ſGk. homos, same; akantha, spine.] Having spines of dorsal fin symmetrical.

- (hömăk'sŏn) homaxon [Gk. homos, same; axon, axis.] up around equal axes; axonic.
- homeo-, also homoeo-, homoio-. homeokinesis (hom'eokine'sis) n. [Gk. homoios, alike; kinein, to

move.] Mitosis with equal division of chromatinic elements to daughter

homeostasis (homeos'tasis) n. [Gk. homoios, alike; stasis, standing.] The constancy of the internal environment of the body, as in birds and mammals.

homeosynapsis,—homosynapsis, q.v. homeotely (hom'eot'eli) n. homoios, alike; telos, end.] Evolution from homologous parts, but with less close resemblance.

homeotypic (hom'eotip'ik) a. [Gk. homoros, alike; typos, character.] Appl. second division in meiosis, similar to typical mitosis;

heterotypic.

homeozoic (hom'ēozo'ik) a. [Gk. homoios, alike; zoon, animal.] Pert. a region or series of regions with identical fauna.

homoblastic (hom'oblas'tik) a. [Gk. homos, same; blastos. bud.] Having direct embryonic development; arising from similar cells.

homobrachial (hŏm'öbrā'kĭǎl) [Gk. homos, same; L. brachium,

- arm.] Pert. the same chromosome arm; paracentric.
- homocarpous (hom'okar'pus) a. [Gk. homos, same; karpos, fruit.] Bearing one kind of fruit.
- homocellular (hom'osel'ular) a. [Gk. homos, same; L. cellula, small Composed of cells of one cell. type only. Opp. heterocellular.
- homocercal (hom'oser'kal) a. [Gk. homos, same; kerkos, tail.] Having a tail with equal or nearly equal lobes, and axis ending near middle of base.
- homocerebrin (hom'oser'ebrin) n. [Gk. homos, same; L. cerebrum. brain.] A substance identical with cerebrin.
- homochlamydeous (hom'oklamid'eŭs) a. [Gk. homos, same; chlamys, cloak.] Having outer and inner perianth whorls alike.
- homochromous (hom'okro'mus) a. [Gk. homos, same; chroma, colour.] Of one colour; appl. capitular
- homochronous (hömŏk'rönŭs) [Gk. homos, same; chronos, time.] Occurring at the same age or period, in successive generations.

homodermic (hom'oder'mik) a. [Gk. homos. same; derma, skin.] Sprung from same embryonic layer.

- homodont (hō'mödŏnt) a. homos, same; odous, tooth.] Having the teeth all alike, not differentiated. Opp. heterodont.
- homodromous (hömŏd'römŭs) [Gk. homos, same; dramein, to run.] Having genetic spiral alike in direction in stem and branches.
- homodynamic (hom'odinam'ik) a. homos, same; dynamis, power.] Developing without resting stages; appl. insects not requiring a diapause for further development; pert. homodynamy.
- homodynamy (hom'odi'nami) homos. same; dynamis. power.] Metameric homology.
- homoeandrous (homean'drus) [Gk. homoios, alike; aner, male.] Having uniform stamens.

homoeo-,-also homeo, homoio.

homoeologous (homeol'ogus) a. [Gk. homoios, like; logos, relation.] Appl. chromosomes having in part the same sequence of genes; partly homologous.

homoeologue, — a homoeologous

chromosome.

homoeomorphic (hom'ēomôr'fik) a. [Gk. homoios, like; morphe, form.] Resembling in shape or structure;

exhibiting convergence.

homoeosis (homē'osis) n. [Gk. homoiosis, likeness.] Assumption by one part of likeness to another part, as modification of antenna into foot, or of petal into stamen; metamorphy.

homoeotype (hom'ēotīp) n. [Gk. homoios, alike: typos, pattern.] A specimen authoritatively stated to be identical with the holotype, lectotype, paratypes, or syntypes of its species.

homoeozoic (hö'mēözō'ik) a. [Gk. homoios, alike; zoe. life.] Characterised by similar forms of life;

appl. areas or zones.

homogametic (hŏm'ögămět'ĭk) a. [Gk. homos, same; gametes, spouse.] Having homogametes or gametes of one type; appl. sex possessing two X-chromosomes; cf. heterogametic, digametic.

homogamous (hömög'ämŭs) a. [Gk homos, same; gamos, marriage.] Characterised by homogamy.

homogamy (hömög'āmĭ) a. [Gk. homos, same; gamos, marriage.] Inbreeding due to some type of isolation; condition of having flowers all alike, having stamens and pistils mature at same time.

homogangliate (hŏm'ögăng'liāt) a. [Gk. homos, same; ganglion, knot.] Having ganglia of nerve loops symmetrically arranged.

homogen (hō'möjĕn) n. [Gk. homos, same; genos, race.] One of a group having a common origin; one of a series of identically derived parts.

homogenesis (hom'öjen'esis) n. [Gk. homos, same; genesis, descent.]

The type of reproduction in which like begets like.

homogenetic (hŏm'öjĕnĕt'īk) a. [Gk. homos, same; genesis, descent.]
Pert. homogenesis; appl. pairing of homologous chromosomes.

homogenous (hömőj'enűs) a. [Gk. homos, same; genos, race.] More or less alike owing to descent from common stock; appl. graft from another animal of same species.

homogeny (hömöj'eni) n. [Gk. homos, same; genos, race.] Correspondence between parts due to common descent; the same genotypical structure; homogeneity.

homogony (homog'öni) n. [Gk. homos, same; gonos, offspring.] Condition of having one type of flower with equally long stamens and pistil.

homoio-,—also homeo-, homoeo-.

homoiomerous (homoiom'erus) a. [Gk. homoios, like; meros, part.] Having algae distributed equally through fungoid mycelium in a lichen.

homoioplastic (hömoi'öpläs'tĭk) a. [Gk. homoios, like; plastos, formed.] Appl. transplantation between individuals of the same species; homeoplastic.

homoiosmotic (hömoi'ōsmŏt'ik) a. [Gk. homoios, like; osmos, impulse.] Appl. organisms with constant internal osmotic pressure; euryhaline, q.v.

homoiothermal (hömoi'öther'mäl) a. [Gk. homoios, like; thermos, hot.] Having a more or less constant body temperature; warm-blooded; homoeothermal, homoiothermic, homothermal, homothermic, homothermous.

homoiotransplantation,—transplantation of tissue or organ from one organism to another, possibly unrelated; cf. autotransplantation.

homolecithal (hŏm'ölĕs'īthăl) a. [Gk. homos, same; lekithos, yolk.] Having little deutoplasm, which is equally distributed.

homologous (hömöl'ögüs) a. [Gk. homologos, agreeing.] Resembling in structure and origin; appl.

various substances, e.g. agglutinins affecting organisms of same species only; appl. chromosomes with the same sequence of genes; appl. genes determining the same character, e.g. eye colour. Cf. heterologous, antithetic.

homologue (hom'olog) n. [Gk. homologos, agreeing.] One of a series of structures similar in structure and origin; a homologous

agent.

homology (hömöl'öji) a. [Gk. homologia, agreement.] Similarity in structure and development of organ or parts.

homomallous (hŏm'ömäl'ŭs) a. [Gk. homos, same; mallos, lock of wool.] Curving uniformly to one side; appl. leaves.

homomorphic (hŏm'ömôr'fīk) a. [Gk. homos, same; morphe, form.] Of similar size or structure; pert., or exhibiting, homomorphism; appl. chromosome pairs; cf. heteromorphic.

homomorphism (höm'ömôr'fizm) n. [Gk. homos, same; morphe, shape.] The condition of having perfect flowers of only one type; hemimetabolism; similarity of larva and adult.

homomorphosis (hŏm'ömôr'fōsĭs) n. [Gk. homos, same; morphosis, shaping.] Condition of having a newly regenerated part like the

part removed.

homonomic (hō'mönōm'īk) a. [Gk. homos, same; nomos, law.] Having the same behaviour; appl. affinity, as of tissues combining, e.g. vascular anastomoses, or complementary affinity, as in adrenal medulla and cortex; homonomous, q.v.

homonomous (hömön'ömüs) a. [Gk. homos, same; nomós, department.] Appl. segmentation into similar segments. [Gk. nómos, law.] Following same stages or process, as of development or growth.

homonomy (hömön'ömi) n. [Gk. homos, same; nómos, law.] The homology existing between parts

arranged on transverse axes; homodynamy.

homonym (hŏm'önim) n. [Gk. homos, same; onyma, name.] A name preoccupied, and therefore unsuitable according to law of priority.

homopetalous (hŏm'öpĕt'ālūs) a. [Gk. homos, same; petalon, petal.] Having all the petals alike.

homophyadic (hŏm'öfīād'īk) a.
[Gk. homos, same; phyas, shoot.]
Producing only one kind of shoot.

homophylic (hom'ofil'ik) a. [Gk. homos, same; phyle, race.] Resembling one another owing to a common ancestry.

homophyllous (hom'ofil'us) a. [Gk. homos, same; phyllon, leaf.] Bear-

ing leaves all of one kind.

homophytic (hom'ofit'ik) a. [Gk. homos, same; phyton, plant.] With two kinds of spores, or one bisexual type, borne by a single sporophyte; cf. heterophytic.

homoplasma (hom'oplaz'ma) n. [Gk. homos, same; plasma, mould.] Plasma from another animal of same species used as a medium for tissue culture; cf. autoplasma, heteroplasma.

homoplast (hom'oplast) n. [Gk. homos, same; plastos, moulded.]
An organ or organism formed of similar plastids; coenobium; catallact.

homoplastic (hŏm'oplăs'tik) a. [Gk. homos, same; plastos, moulded.] Pert. homoplasty; appl. graft made into individual of same species; cf. autoplastic.

homoplasty (hŏm'öplăs'tĭ) n. [Gk. homos, same; plastos, moulded.] Convergence; resemblance in form of structure between different organs or organisms due to evolution along similar lines; also homoplasy; isotely.

homopterous (hömop'tërus) a. [Gk. homos, same; pteron, wing.] Hav-

ing the wings alike.

homorhizal (hŏm'örī'zăl) a. [Gk. homos, same; rhiza, root.] Not having an antiapical root, as Pteridophyta. Opp. allorhizal.

- homosomal (hŏm'ösō'măl) a. [Gk. homos, same; soma, body.]
 Occurring in, or pert., the same body; appl. rearrangements restricted to a single chromosome.
- homosporangic (hom'osporan'jik) a. [Gk. homos, same; sporos, seed; anggeion, vessel.] Bearing spores of one kind or of two kinds in one sporangium.
- homosporous (hŏmöspō'rŭs) a. [Gk. homos, same; sporos, seed.] Producing only one kind of spore; homosporic; isosporous.

homostyled (höm'ostīld) a. [Gk. homos, same; stylos, pillar.] With uniform styles; homogonous.

homosynapsis (hom'osinap'sis) n. [Gk. homos, same; synapsis, union.] Pairing of two homologous chromosomes; cf. heterosynapsis.

homotaxial (hom'otak'sial) a. [Gk. homos, same; taxis, arrangement.] Containing the same assemblage of species, appl. fossiliferous deposits.

homotaxis (hŏm'ötāk'sīs) n. [Gk. homos, same; taxis, arrangement.] Similar assemblage or succession of species or types in different regions or strata, not necessarily contemporaneous.

homothallic (hom'othal'ik) a. [Gk. homos, same; thallos, young shoot.] Forming zygospores from two branches of the same mycelium; appl. moulds. Opp. heterothallic.

homothermous, -- homoiothermal, q.v.

homotropous (hömŏt'röpŭs) a. [Gk. homos, same; trope, turn.] Erect; having micropyle and chalaza at opposite ends; appl. ovules.

homotypic (hōmōtip'ik) a. [Gk. homos, saine; typos, pattern.] Homeotypic, q.v.; pert. or exhibiting homotypy.

homotypy (hōm'ötīpī) n. [Gk. homos, same; typos, pattern.] Equality of structures along main axis of body; serial homology; reversed symmetry; enantiomorphic condition, q.v.

homoxylous (hom'ozi'lus) a. [Gk. homos, same; xylon, wood.] Appl.

wood without vessels and consisting of tracheids.

homozygosis (hom'ozigo'sis) n. [Gk. homos, same; zygon, yoke.] Condition of having a given genetical factor in the duplex condition, and producing gametes of only one kind as regards that factor.

homozygote (höm'özī'gōt) n. [Gk. homos, same; zygon, yoke.] An organism in which characters are stable, resulting from union of gametes bearing similar genes.

homozygous (hŏm'özī'gūs) a. [Gk. homos, same; zygon, yoke.] Having identical genes for a given character; exhibiting or pert. homozygosis; pert. homozygote.

homunculus (hömun'kūlūs) n. [L. homunculus, little man.] The small miniature of human foetus supposed to be in spermatozoon, according to Animalculists; homonculus.

honey-dew, — a sugary exudation found on leaves of many plants; a viscous fluid secreted by mycelium of ergot; a sweet secretion produced by certain insects, e.g., by aphids.

hooded (hood'ëd) a. [A.S. hōd.]
Bearing a hood-like petal; cucullate; rolled up like a cone of paper, as certain leaves; having head conspicuously and differently coloured from rest of body; having crests on head; having wingshaped expansions on neck, as in cobra.

hook-glands,—paired longitudinal glands uniting anteriorly to form head gland in Pentastomida.

hordeaceous (hôr'deā'shūs) a. [L. hordeum, barley.] Pert. or resembling barley.

horiodimorphism (hō'rĭödīmôr'fĭzm)

n. [Gk. horios, in season; dis,
twice; morphe, shape.] Seasonal
dimorphism.

horizon (höri'zön) n. [Gk. horizon, bounding.] Soil layer of a more or less well-defined character; a layer of deposit characterised by definite fossil species and formed at a definite time.

horizontal (hŏr'Izŏn'tăl) a. [Gk. horizon. bounding.] Growing in a plane at right angles to primary axis.

horme (hôr'mē) n. [Gk. horme, impetus.] Purposive behaviour; conation; urge or drive in living cells or organisms; élan vital.

hormocyst (hôr'mosīst) n. [Gk. hormos, chain; kystis, bladder.]
A modified thick-walled hormogonium, in some blue-green algae.

hormogonium (hôr'mögō'nĭŭm) n. [Gk. hormos, chain; gone, generation.] That portion of an algal filament between two heterocysts, which, breaking away, acts as a reproductive body; hormogone.

hormones (hôrmōnz) n. plu. [Gk. hormaein, to excite.] Substances normally produced in cells and necessary for the proper functioning of other distant cells to which they are conveyed and of the body as a whole; internal secretions of ductless glands which pass into blood vessels by osmosis; exciting agents, opp. chalones; internal secretions in plants, as auxins, q.v.

hormonic (hôrmon'īk) a. [Gk. hormaein, to excite.] Pert. hormones; appl excitatory internal secretions, opp. chalonic.

hormoproteins (hôr'möprō'teinz) n.
plu. [Gk. hormaein, to excite; protos,
first.] Proteins or protein derivatives secreted by endocrines.

horn (hôrn) n. [A.S. horn.] The process on head of many animals; any projection resembling a horn; anterior part of each uterus when posterior parts are united to form median corpus uteri; a tuft of feathers as in owl; a spine in fishes; a tentacle in snails; an awn; any pointed projection or process in plants; cornu.

horn core,—the os cornu, fusing with frontal bone, over which fits hollow horn of ruminants.

horodimorphism, — horiodimorphism, q.v.

horotelic (hōrötěl'ik) a. [Gk. hora,

right time; telos, fulfilment.] Evolving at the standard rate; cf. bradytelic, tachytelic.

host (hōst) n. [L. hostis, stranger.]
Any organism in which another organism spends part or the whole of its existence, and from which it derives nourishment or gets protection; an organism which receives grafted or transplanted tissue.

house (hows) n. [A.S. hūs.] The external gelatinous - like covering secreted by certain tunicates.

Houston's valves [J. Houston, Irish surgeon]. Semilunar transverse folds of mucous membrane in the rectum; plicae transversales recti.

humeral (hū'mērăl) a. [L. humerus, shoulder.] Pert. shoulder region; pert. the anterior basal angle of insect wing, appl. a cross vein; one of horny plates on plastron of chelonians.

humerus (hū'mėrūs) n. [L. humerus, shoulder.] The bone of the upper

humistratous (hū'mistrā'tŭs) a. [L. humus, soil; sternere, to spread.] Spreading over surface of ground.

humoral (hū'morāl) a. [L. humor, moisture.] Appl. theory of immunity ascribing to body fluids the power to resist infection.

humour (hū'mŏr) n. [L. humor, moisture.] Any fluid or juice; the fluid of the eye.

humus (hū'mūs) n. [L. humus, earth.] A dark material formed by

decomposition of vegetable or animal matter and constituting organic part of soils.

husk (husk) n. [M.E. huske.] The outer coating of various seeds.

Huxley's layer [T. H. Huxley, English zoologist]. The middle layer of polyhedral cells in inner rootsheath of hair.

hyaline (hī'ālīn) a. [Gk. hyalos, glass.] Clear; transparent; free from inclusions.

hyalodermis (hī'ālödēr'mīs) n. [Gk. hyalos, glass; derma, skin.] Tissue of large, empty and absorptive cells in Sphagnum.

- hyalogen (hī'ālöjěn) n. [Gk. hyalos, glass; -genes, producing.] Any of substances found in animal tissues which are insoluble and related to mucoids.
- hyaloid (hi'aloid) a. [Gk. hyalos, glass; eidos, form.] Glassy; transparent.
- hyaloid artery,---from central artery of retina through hyaloid canal to back of lens, in foetal eye.
- canal,—through vitreous body of eye, from optic nerve to back of lens.
- hyaloid fossa,—anterior concavity of vitreous body, receptacle of crystalline lens.
- hyaloid membrane,-delicate membrane enveloping vitreous body of
- hyalomere (hī'ālömēr) n. [Gk. hyalos, glass; meros, part.] clear homogeneous part of a blood platelet, opp. chromomere.
- hyalomucoid (hī'ălömū'koid) n. [Gk. hyalos, glass; L. mucus, mucus; Gk. eidos, like.] One of the nonphosphorised gluco - proteids in vitreous humour.
- hyaloplasm (hī'ālöplāzm) n. hyalos, glass; plasma, mould.] Ground substance of cell as distinguished from microsomes, or from reticulum or spongioplasm, ectoplasm or peripheral zone in plant cells; also hyaloplasma.

hyalopterous (hī'ālŏp'tërŭs) a. [Gk. hyalos, glass; pteron, wing.] Hav-

ing transparent wings.

hyalosome (hī'ālösōm) [Gk. hyalos, glass; soma, body.] nucleolar-like body in a cell-nucleus, only slightly stainable by nuclear or plasma stains.

hyalosporous (hī'alöspo'rus, hī'alos'pörüs) a. [Gk. hyalos, glass; sporos, seed.] Having colourless spores or conidia.

hybrid (hī'brid) n. [L. hibrida, cross.] Any cross-bred animal or plant; heterozygote. a. Cross-bred; heterozygous.

incapacitation, - hybrid hybrid sterility and inviability, inclusively. hybridisation (hī'bridīzā'shun) n. [L. hibrida, cross.] Act or process of hybridising; state of being hybridised; cross-fertilisation.

hybridise (hī'brīdīz) v. [L. hibrida, cross.] To interbreed, to cross, to

produce hybrids.

- hybridism (hī'brīdizm) n. hibrida, cross.] The state or quality of being a hybrid.
- hydathode (hī'dăthōd) hydatos, of water; hodos, way.] An epidermal structure specialised for secretion, or for exudation, of water; water stoma.
- hydatid (hī'dătid) n. [Gk. hydatis, watery vesicle.] Any vesicle or sac filled with clear watery fluid; sac containing encysted stages of larval tapeworms; vestige of Müllerian duct constituting appendix of testis, hydatid of Morgagni; stalked appendix of epididymis.

hydatiform (hī'dătĭfôrm) a. hydatis, watery vesicle; L. forma, shape.] Resembling a hydatid.

- hydatigenous (hī'dătĭj'ënŭs) a. [Gk. hydatis, watery vesicle; -genes, producing] Producing or forming hydatids.
- hydranth (hī'dranth) n. [Gk. hydor, water; anthos, flower. | A nutritive zooid in a hydroid colony.
- hydrarch (hī'drark) n. [Gk. hydor, water; arche, beginning.] Appl.seres progressing from hydric towards mesic conditions.
- hydric (hī'drīk) a. [Gk. hydor. water.] Characterised by an abundant supply of moisture, appl. plants, environment.
- hydrocaulis (hī'drökôl'is) n. [Gk. hydor, water; L. caulis, stalk.] The branching vertical portion of coenosarc in a hydroid col-
- hydrocircus (hī'drösēr'kūs) n. [Gk. hydor, water; kirkos, circle.] The hydrocoelic ring surrounding mouth in echinoderms.
- hydrocladia (hī'drö'klā'dĭā) n. plu. [Gk. hydor, water; klados, shoot.] The secondary branches of a hydrocaulis.

- hydrocoel (hī'drösēl) n. [Gk. hydor, water; koilos, hollow.] The watervascular system in echinoderms.
- **hydrocryptophyte**, hydrophyte, q.v.
- hydrocyst (hī'drösist) n. [Gk. hydor, water; kystis, bladder.] A dactylozooid.
- hydroecium (hidrē'siŭm) n. [Gk. hydor, water; oikos, house.] A closed tube at upper end of a siphonophore; an infundibulum.
- hydroid (hī'droid) n., a. [Gk. hydor, water; eidos, form.] Elongated empty cell in central cylinder of mosses; a tracheid; the polyp of Hydrozoa.
- hydrolysis (hīdrŏl'īsīs) n. [Gk. hydor, water; lyein, to dissolve.] The reaction between a chemical compound and the hydrogen and hydroxyl ions of water.
- hydrome (hi'drōm) n. [Gk. hydor, water; mestos, full.] Any tissue that conducts water; also hydrom
- hydromegatherm (hī'dromeg'atherm)

 n. [Gk. hydor, water; megas,
 great; therme, heat.] A plant
 which must have much moisture
 and heat to develop fully.
- hydromorphic (hīdromôr fik) a. [Gk. hydor, water; morphe, form.] Structurally adapted to an aquatic environment, as organs of water plants.
- hydronasty (hī'drönăs'tĭ) n. [Gk. hydor, water; nastos, closepressed.] Plant movement induced by changes in atmospheric humidity.
- hydrophilous (hī'drŏf'īlŭs) a. [Gk. hydor, water; philein, to love.]
 Pollinated through agency of water.
- hydrophoric (hī'dröfðr'ik) a. [Gk. hydrophoros, carrying water.] Appl. canal, the madreporic or stone canal, q.v.
- hydrophyllium (hī'dröfil'iūm) n. [Gk. hydor, water; phyllon, leaf.] One of leaf-like transparent bodies arising above and partly covering the sporosacs in a siphonophore.

hydrophyte (hī'dröfīt) n. [Gk. hydor, water; phyton, plant.] An aquatic plant; cf. hygrophyte.

hydrophyton (hīdrofī'tŏn) n. [Gk. hydor, water; phyton, plant.] A complete hydroid colony, root-like organ, stem and branches.

hydroplanula (hī'droplān'ūlă) n [Gk. hydor, water; I.. planus, flat.] Stages between planula and actinula in larval history of coelenterates.

- hydropolyp (hī'dropŏl'ĭp) n. [Gk. hydor, water; F. polype, polyp.] A polyp of a hydroid colony; a hydrula.
- hydroponics (hī'dropon'īks) n. [Gk. hydor, water; ponos, exertion.]
 The science and art of crop production in liquid culture media; cf. water culture.
- hydropore (hī'dropōr) n. [Gk. hydor, water; poros, opening.] The opening into right hydrocoel in echinoderm larvae.
- hydropote (hī'dröpōt) n. [Gk. hydropotes, water drinker.] A cell or cell-group, in some submerged leaves, easily permeable by water and salts.
- hydropyle (hī'dröpīlē) n. [Gk. hydor, water; pyle, gate.] Specialised area in cuticular membrane of embryo, for passage of water, as in grasshoppers.
- hydrorhabd (hī'drörābd) n. [Gk. hydor, water; rhabdos, rod.] A rhabdosome, in graptolites.
- hydrorhiza (hī'drorī'ză) n. [Gk. hydor, water; rhiza, root.] The creeping root-like portion of coenosarc of a hydroid colony.
- hydrosere (hī'drösēr) n. [Gk. hydor, water; L. serere, to put in a row.] A plant succession originating in a wet environment.
- hydrosome (hī'drösōm) n. [Gk. hydor, water; soma, body.] The conspicuously hydra-like stage in a coelenterate life-history; hydrosoma.
- hydrospire (hī'dröspīr) n. [Gk. hydor, water; L. spira, coil.] The folds on the stereom of blastoids, being respiratory structures.

- hydrostatic (hī'drostāt'īk) a. [Gk. hydor, water; statikos, causing to stand.] Appl. organs of flotation, as air sacs in aquatic larvae of insects.
- hydrostome (hī'dröstōm) n. [Gk. hydor, water; stoma, mouth.] The mouth of a hydroid polyp.

hydrotaxis (hī'drötāk'sĭs) n. [Gk. hydor, water; taxis, arrangement.] Response of organisms to stimulus of moisture.

hydrotheca (hī'dröthē'kā) n. [Gk. hydor, water; theke, cup.] Cup-like structure into which the polyp may withdraw, in many coelenterates.

hydrotropic (hī'drotrop'īk) a. [Gk. hydor, water; trope, turn.] Appl. curvature of a plant organ towards a greater degree of moisture.

hydrotropism (hī'drŏt'röpĭzm) n.
[Gk. hydor, water; trepein, to
turn.] Response to stimulus of
water.

hydrula (hī'droolă) n. [Gk. hydor, water.] Hypothetical simple polyp. hyetal (hī'etāl) a. [Gk. hyetos, rain.] Pert. rain; pert. precipitation.

hygiene (hi'jēn) n. [Gk. hygieinos, healthful.] That part of biology dealing with health preservation.

hygric (hi'grik) a. [Gk. hygros, wet.] Humid; tolerating, or adapted to, humid conditions. Οφρ. xeric.

hygrochasy (hī'grökā'si) n. [Gk. hygros, wet; chasis, separation.]
Dehiscence of seed vessels when induced by moisture; cf. xerochasy.

hygrokinesis (hī'grökinē'sīs) n. [Gk. hygros, wet; kinesis, movement.] Movement in response to change in humidity.

hygrophilous (hīgrōf'īlūs) a. [Gk. hygros, wet; philein, to love.]
Inhabiting moist or marshy places.

hygrophyte (hī'gröfīt) n. [Gk. hygros, wet; phyton, plant.] A plant which thrives in plentiful moisture; cf. hydrophyte.

hygroplasm (hī'gröplāzm) n. [Gk. hygros, wet; plasma, mould.] The more liquid part of protoplasm, opp. stereoplasm.

hygroscopic (hī'gröskŏp'īk) a. [Gk. hygros, wet; skopein, to regard.] Sensitive to, or retaining, moisture. hylion (hī'lĭön) n. [Gk. hyle, wood; on, being.] Forest climax; hylium.

hylophagous (hilof'agus) a. [Gk. hyle, wood; phagein, to eat.] Eating wood; appl. certain insects.

hylophyte (hī'löfit) n. [Gk. hyle, wood; phyton, plant.] A plant growing in woods.

hylotomous (hī'lot'ömus) a. [Gk. hyle, wood; temnein, to cut.] Wood-cutting; appl. certain insects. hymen (hī'men) n. [Gk. hymen, membrane.] Thin fold of mucous

membrane at orifice of vagina. hymenial (hīmē'nĭăl) a. [Gk. hymen,

skin.] Pert. hymenium.

hymeniferous (hī'mēnif'érūs) a. [Gk. hymen, skin; L. ferre, to carry.] Having a hymenium.

hymenium (hīmē'nīum) n. [Gk. hymen, skin.] The outermost layer of mushroom lamellae, or of other fungi, consisting of spore-producing cells interspersed with barren cells or paraphyses.

hymenophore (hī'mënöfōr) n. [Gk. hymen, skin; pherein, to carry.] The hymenial portion of sporophore of a fungus.

hymenopterous (hī'mēnŏp'tērūs) a. [Gk. hymen, skin; pteron, wing.] Having membranous wings; appl. certain insects.

hyobranchial (hī'öbrăng'kĭăl) a. [Gk. Y; brangchia, gills.] Pert. hyoid and branchial arches.

hyoepiglottic (hī'öĕpīglot'ik) a. [Gk. Y; epi, upon; glotta, tongue.] Connecting hyoid and epiglottis.

hyoglossal (hī'öglös'ăl) a. [Gk. Y; glossa, tongue.] Pert. tongue and hyoid; appl. membrane and muscle.

hyoid (hī'oid) a. [Gk. hyoeides, Y-shaped.] Pert. or designating a bone or series of bones lying at base of tongue and developed from hyoid arch of embryo; appl. a sclerite enclosing pharynx in some insects; hyoidean. n. The hyoid bone.

hyoidean (hīoid'ēan) a. [Gk. hyoeides, Y-shaped.] Pert. or associated with the hyoid arch or bone; appl. a branch of first efferent branchial vessel, or of lingual artery; appl. nerve, the posterior post-trematic nerve.

hyoideus (hīoid'eŭs) n. [Gk. hyoeides, Y-shaped.] A nerve which supplies mucosa of mouth and muscles of hyoid region.

hyomandibular (hī'omăndib'ūlăr) a. [Gk. Y; L. mandibulum, jaw.] Pert. hyoid and mandible; pert. dorsal segment of hyoid arch in fishes. hyomental (hī'omēn'tāl) a. [Gk. Y; L. mentum, chin.] Pert. hyoid and

chin.

hyoplastron (hī'öplăs'trŏn) n. [Gk.

Y; F. plastron, breast-plate.] The
second lateral plate in plastron of

Chelonia.

hyostapes (hī'ostā'pēz) n. [Gk. Y; L.L. stapes, stirrup.] Lower portion of columellar primordium which gives rise to part of columella in some reptiles.

hyosternum (hī'östĕr'num) n. [Gk. Y; sternon, breast.] Hyoplastron. hyostylic (hī'östīl'īk) a. [Gk. Y; stylos, pillar.] Having jaw articulated to skull by hyomandibular or corresponding part; exhibiting hyostyly; cf. autostylic.

hyosymplecticum (hī'ösimplěk'tīkŭm) n. [Gk. Y; symplektos, plaited together.] The cartilaginous primordium from which hyomandibular and symplecticum are

derived.

hyothyroid (hī'öthī'roid) a. [Gk. Y; thyreos, shield; eidos, like.] Pert. hyoid bone and thyroid cartilage of larynx; appl. ligaments, membrane.

hypallelomorph (hip'ālēl'omôrf) n. [Gk. hypo, under; allelon, of one another; morphe, form.] Allelomorphs which under certain conditions are themselves compound.

hypandrium (hǐpăn'drium) n. [Gk. hypo, under; aner, male.] Subgenital plate or ninth abdominal sternite of certain insects.

hypanthium (hǐpăn'thiŭm) n. [Gk. hypo, under; anthos, flower.] Any enlargement of the torus.

hypanthodium (hip'anthō'dium) n. [Gk. hypo, under; anthodes, like flowers.] An inflorescence with concave capitulum on whose walls the flowers are arranged.

hypantrum (hǐpăn'trum) n. [Gk. hypo, under; antron, cave.] Notch on vertebrae of certain reptiles for articulation with hyposphene.

hypapophysis (hip'apof'isis) n. [Gk. hypo, under; apo, upon; phyein, to grow.] A ventral process on a vertebral centrum.

hyparterial (hip'arte'riăl) a. [Gk. hypo, under; L. arteria, artery.] Situated below an artery; appl. branches of bronchi below pulmonary artery.

hypaxial (hǐpāk'siǎl) a. [Gk. hypo, under; I.. axis, axis.] Ventral; below vertebral column; appl. muscles.

hyperapophysis (hī'përăpŏf'isis) n. [Gk. hyper, above; apo, from; phyein, to grow.] A postero-lateral process of dorsal side of vertebra.

hyperchromasy (hī'përkrō'măsĭ) n. [Gk. hyper, above; chroma, colour.] A relatively superabundant supply of chromatin to cytoplasm in a cell.

hyperchromatosis (hī'pērkrō'mătōsis) n. [Gk. hyper, above; chroma, colour.] Excess of nuclear substance in a cell previous to division.

hypercoracoid (hī'përkŏr'ākoid) a. [Gk. hyper, above; korax. crow; eidos, form.] Pert. or designating upper bone at base of pectoral fin in fishes.

hypercyesis (hī'pērsīē'sis) n. [Gk. hyper, above; kyesis, conception.] Superfoctation; additional fertilisation in a mammal already pregnant.

hyperfeminisation,—condition of a feminised male with female characteristics exaggerated, as in small size and weight.

hypergamesis (hī'pērgămē'sīs) n. [Gk. hyper, above; gamos, marriage.] Process of absorption by female of excess spermatozoa.

hypermasculinisation, — condition of a masculinised female with male characteristics exaggerated, as in large proportions, appearance of male secondary sexual characters.

hypermetamorphosis (hī'pērmēt'āmôr'fōsis) n. [Gk. hyper, above; meta, after; morphosis, shaping.] A protracted and thoroughgoing metamorphosis; metamorphosis involving two or more distinct types of larval instar, in certain insects.

hypermorphosis (hī'pērmôr'fōsīs) n.
[Gk. hyper, above; morphosis, shaping.] The development of additional characters, in comparison with the adult ancestral stage.

hyperparasite (hi'përpăr'ăsît) n. [Gk. hyper, above; para, beside; sitos, food.] A parasite which is parasitic on or in another parasite.

hyperphalangy (hī'pērfāl'ānjī) n. [Gk. hyper, above; phalangx, line of battle.] Condition of having digits with more than normal number of phalanges.

hyperpituitarism(hī'pērpītū'itārīzm)

n. [Gk. hyper, above; L. pituita,
phlegm.] Overaction of pituitary
gland, resulting in gigantism or
giantism; hyperhypophysism.

hyperplasia (hī'pērplā'siā) n. [Gk. hyper, above; plassein, to mould.]
Overgrowth; excessive or hyperplastic development due to increase in number of cells; cf. hypertrophy.

hyperploid (hī'përploid) a. [Gk. hyper, above; haploos, onefold; eidos, form.] Aneuploid with extra chromosomes, opp. hypoploid.

hyperpnoea (hī'përpnē'ă) n. [Gk. hyper, above; pnoe, breath.] Rapid breathing due to insufficient supply of oxygen.

hypersensitivity (hī'përsĕn'sĭtīv'ītī)

n. [Gk. hyper, above; L. sentire,
to feel.] A condition of being
unduly sensitive to a stimulus.

hypertely (hipěr'tëli) n. [Gk. hyper, above; telos, end.] Excessive imitation in colour or pattern, being of problematical utility; overdevelopment, as canines of Babirusa; hypertelia.

hypertensin,—angiotonin, q.v.

hypertonia (hī'pērtō'niā) n. [Gk. hyper, above; tonos, tone.] Excessive tonicity.

hypertrophy (hīpěr'tröfi) n. [Gk. hyper, above; trophe, nourishment.] Excessive growth due to increase in size of cells; cf. hyperplasia.

hypha (hī'fā) n. [Gk. hyphe, web.]
The thread-like element or filament of vegetative mycelium of a fungus; filamentous cell in medulla of an algal thallus.

hyphasma (hīfāz'mă) n. [Gk. hyphasma, thing woven.] A barren mycelium; an Agaricus thallus.

hyphodrome (hī'födrōm) a. [Gk. hyphe, web; dromos, course.]
Running throughout the tissues; appl. thick leaves where veins are not visible from surface.

hyphopodium (hīföpō'diŭm) n. [Gk. hyphe, web; pous, foot.] A hyphal branch with enlarged terminal cell or haustorium for attaching the hypha, as in some Ascomycetes.

hypnocyst (hip'nösist) n. [Gk. hypnos, sleep; kystis, bladder.]
 Cyst in which contained organism simply rests; dormant cyst.

hypnody (hip'nödi) n. [Ġk. hypnodia, sleepiness.] The long resting period of certain larval forms.

hypnogenic (hǐp'nöjĕn'ik) a. [Gk. hypnos, sleep; -genes, producing.] Sleep-inducing; appl. influences which tend to produce hypnosis.

hypnosperm (hĭp'nösperm) n. [Gk. hypnos, sleep; sperma, seed.] A hypnospore.

hypnosporangium (hĭp'nöspörăn'jiŭm) n. [Gk. hypnos, sleep; sporos, seed; anggeion, vessel.] A sporangium containing resting spores.

hypnospore (hip'nöspōr) n. [Gk. hypnos, sleep; sporos, sced.] A resting spore; a zygote that remains in a quiescent condition during winter.

hypnote (hǐp'nōt) n. [Gk. hypnos, sleep.] An organism in a dormant condition.

hypoachene (hī'pöǎkēn') n. [Gk. hypo, under; a, not; chainein, to

gape.] Achene developed from an inferior ovary.

hypoarion (hī'pōā'riŏn) n. [Gk. hypo, under; oarion, little egg.] A small lobe below the optic lobes of most teleosts.

hypobasal (hi'pöbā'săl) n. [Gk. hypo, under; basis. base.] The lower segment of a developing ovule, which ultimately gives rise to the root. Opp. epibasal.

hypoblast (hī'poblāst) n. [Gk. hypo, under; blastos, bud.] The inner

germ layer in a gastrula.

hypoblastic (hi'pöblăs'tik) a. [Gk. hypo, under; blastos, bud.] Pert., or derived from, the inner layer; endodermal.

hypobranchial (hi'pobrăng'kiăl) a. [Gk. hypo, under; brangchia, gills.] Pert. lower or fourth segment of branchial arch.

hypocentrum (hī'pösĕn'trŭm) n. [Gk. hypo, under; kentron, centre.] A transverse cartilage that arisebelow nerve cord and forms part of vertebral centrum.

hypocercal (hī'posĕr'kăl) a. [Gk hypo, under; kerkos, tail.] Having notochord terminating in lower lobe of caudal fin.

hypochilium (hī'pokī'liŭm) n. [Gk. hypo, under; cherlos, lip.] The lower portion of lip of an orchid.

hypochondrium (hī'pokŏn'drīum) n. [Gk. hypo, under; chondros, cartilage.] A region lateral to epigastric and above lumbar.

hypochord (hī'pökôrd) n. [Gk. hypo, under; chorde, cord.] A transitory subnotochordal rod in anamniotes.

hypochordal (hī'pökôr'dāl) a. [Gk. hypo, under; chorde, cord.] Below the notochord; appl. lower lobe of caudal fin; appl. bar of mesodermal tissue developing into ventral arch of atlas and amalgamating with fibrocartilages in other cervical vertebrae.

hypocleidium (hī'pöklīdī'um) n. [Gk. hypo, under; kleis, key; idion, dim.] The interclavicle.

hypocone (hī'pökōn) n. [Gk. hypo, under; konos, cone.] Postero-

internal cusp of upper molar; the part posterior to girdle in Dinoflagellata, opp. epicone.

hypoconid (hī'pokō'nĭd) n. [Gk. hypo, under; konos, cone.] Postero-buccal cusp of lower molar.

hypoconule (hīpokon'ūl) n. [Gk. hypo, under; konos, cone.] Fifth or distal cusp of upper molar.

hypoconulid (hi'pòkŏn'ūlid) n. [Gk. hypo, under; konos, cone.] Posteromesial cusp of lower molar.

hypocoracoid (hī'pokŏr'ākoid) a. [Gk. hypo, under; korax, crow; eidos, like.] Pert. lower bone at base of pectoral fin in fishes.

hypocotyl (hi'pokŏt'il) n. [Gk. hypo, under; kotyle, cup.] That portion of stem below cotyledons in an embryo.

hypocotyledonary (hi'pököt'ilē'dŏnări) a. [Gk. hypo, under; kotyledon, hollow.] Below the cotyledons.

hypocrateriform (hī'pokrātēr'Ifôrm)
a. [Gk. hypo, under; krater, bowl;
L. forma, shape.] Saucer-shaped;
having a gamopetalous corolla with long narrow tube, and limbs at right angles to tube;
hypocraterimorphous.

hypodactylum (hī'pòdāk'tīlūm) n. [Gk. hypo, under; daktylos, digit.] The under surface of a bird's toes.

hypoderma (hī'pödĕr'mă) n. [Gk. hypo, under; derma, skin.] Hypodermis or tissue just under epidermis in plants.

hypodermal (hī'pödēr'māl) a. [Gk. hypo, under; derma, skin.] Pert. hypoderma; pert. hypodermis.

hypodermalia (hī'pödērmā'liā) n.
plu. [Gk. hypo, under; derma, skin.]
Sponge spicules situated just below
the derma or skin.

hypodermic (hī'pödēr'mīk) a. [Gk. hypo, under; L. dermis, skin.] Pert. parts just under the skin.

hypodermis (hī'pödēr'mis) n. [Gk. hypo, under; L. dermis, skin.] The cellular layer lying beneath and secreting the cuticle of Annulata, Arthropoda, etc.; hypoblast, q.v.; hypoderma, q.v.

hypodicrotic (hī'pödikröt'ik) a. [Gk. hypo, under; di, two; krotein, to beat.] Having two arterial beats for the one cardiac.

hypogastric (hip'ögás'trik) a. [Gk. hypo, under; gaster, stomach.]
Pert. lower median region of abdomen; appl. artery, vein, plexus, etc.

hypogastrium (hip'ögäs'triüm) n. [Gk. hypo, under; gaster, stomach.] Lower median region of abdomen. hypogastroid,—hypoischium, q.v.

hypogeal (hī'pöjē'ál) a. [Gk. hypo, under; ge, earth.] Undergound; appl. stems, etc.; also hypogean.

hypogenesis (hī'pöjĕn'ësĭs) n. [Gk. hypo, under; genesis, origin.] Development without occurrence of alternation of generations.

hypogenous (hīpōj'ënŭs) a. [Gk. hypo, under; -genes, produced.] Growing on lower surface of anything.

hypogeous (hī'pojē'us) a. [Gk. hypo, under; ge, earth.] Growing or maturing under earth's surface.

hypoglossal (hī'pöglös'āl) n. [Gk. hypo, under; glossa, tongue.] The twelfth paired cranial nerve, distributed to base of tongue.

hypoglottis (hī'pöglöt'is) n. [Gk. hypo, under; glotta, tongue.] The under part of tongue; a division of labium of beetles.

hypognathous (hīpōg'nāthūs) a. [Gk. hypo, under; gnathos, jaw.] Having the lower jaw slightly longer than the upper.

hypogynium (hī'pöjīn'iŭm) n. [Gk. hypo, under; gyne, female.] Structure supporting ovary in such plants as sedges.

hypogynous (hīpōj'īnŭs) a. [Gk. hypo, under; gyne, female.] Inserted below the gynoecium, and not adherent; immediately below oogonium, appl. antheridium, as in some Peronosporales.

hypohyal (hī'pöhī'ál) n. [Gk. hypo, under; hyoeides, Y-shaped.] The hyoid element lying between ceratohyal and basihyal.

hypoischium (hī'pöis'kium) n. [Gk. hypo, under; ischion, hip.] A

small bony rod passing backwards from ischiadic symphysis and supporting ventral cloacal wall; hypogastroid; os cloacae.

hypolemmal (hī'pölēm'āl) a. [Gk. hypo, under; lemma, peel.] Beneath the sarcolemma; appl. arborisation of an axis cylinder in a motor plate.

hypolimnion (hī'pölimnī'ŏn, -lim'nyŏn) n. [Gk. hypo, under;
limnaios, stagnant; limne, marshy
lake.] The water between the
thermocline and bottom of lakes.
Opp. epilimnion.

hypomeral (hi'pom'ëral) a. [Gk. hypo, under; meros, part.] Hypomeric; appl. slender bones among lower trunk muscles in some fishes.

hypomere (hī'pömēr) n. [Gk. hypo, under; meros, part.] Lower or lateral plate zone of coelomic pouches.

hypomeron (hīpom'ēron) n. [Gk. hypo, under; meros, part.] The lateral inflexed side of a coleopterous prothorax.

hypomorph (hī'pömôrf) n. [Gk. hypo, under; morphe, form.] A gene having a smaller effect than its wild-type allelomorph.

hyponasty (hī'pönāstī) n. [Gk. hypo, under; nastos, close-pressed.] The state of growth in a flattened structure in which the under surface grows more vigorously than the upper.

hyponeural (hī'pönū'răl) a. [Gk. hypo, under; neuron, nerve.] Appl. system of radial and transverse motor nerves in echinoderms.

hyponome (hī'ponōm) n. [Gk. hyponomos, underground passage.] The funnel of Cephalopoda.

hyponychium (hī'pönīk'ĭŭm) n. [Gk. hypo, under; onyx, nail.] Layer of epidermis on which nail rests.

hypopetalous (hī'pöpēt'ālūs) a. [Gk. hypo, under; petalon, petal.] Having corolla inserted below, and not adherent to, gynoecium.

hypophamine,—see pitocin, pitressin. hypophare (hī'pöfār) n. [Gk. hypo, under; pherein, to bear.] Lower part of sponge, in which there are no chambers; cf. spongophare.

hypopharyngeal (hi'pöfärin'jëäl) a. [Gk. hypo, under; pharyngx, pharynx.] Pert. or situated below or on lower surface of pharynx.

hypopharynx (hī'pöfār'ingks) n.
[Gk. hypo, under; pharyngx, pharynx.] The lingua of many insects; in mosquitoes, an outgrowth from base of labium which bears the salivary groove or duct.

hypophloeodal (hī'pöflē'ödăl) a. [Gk. hypo, under; phloios, bark.] Living or growing under bark.

hypophyllium (hī'pöfil'ium) n. [Gk. hypo, under; phyllon, leaf.] A scale-like leaf below a cladophyll; base of stipulate leaf, forming abscission layer.

hypophyllous (hī'pöfil'ŭs) a. [Gk. hypo, under; phyllon, leaf.]
Located or growing under a leaf.

hypophysectomy (hī'pöfisek'tömi) n. [Gk. hypo, under; physis, growth; ek, out; temnein, cut.] Excision or removal of the pituitary gland.

hypophysial (hī'pöfiz'iāl) a. [Gk. hypo, under; physis, growth.]
Pert. the hypophysis.

hypophysin (hīpòf'īsīn) n. [Gk. hypo, under; physis, growth.]
Pituitary extract.

hypophysis (hīpŏf'īsĭs) n. [Gk. hypo, under; physis, growth.] The pituitary body; also hypophyseos; the olfactory pit in the lancelet; the last cell of the suspensor; the cell from which root-tip arises in dicotyledons.

n. [Gk. hypo, under; L. pituita, phlegm.] Deficiency of pituitary gland, resulting in a type of infantilism; hypohypophysism.

hypoplasia (hī'pöplā'siă) n. [Gk. hypo, under; plasis, formation.]
Developmental deficiency; hypoplastic development; deficient growth.

hypoplastron (hī'pöplăs'trŏn) n. [Gk. hypo, under; F. plastron, shield.]
The third lateral bony plate in plastron of Chelonia.

hypopleuron (hī'pöploor'ŏn) n. [Gk.

hypo, under; pleura, side.] Region below metapleuron in insects.

hypoploid (hī'pöploid) a. [Gk. hypo, under; haploos, onefold; eidos, form.] Aneuploid with fewer chromosomes; lacking one chromosome of the complement. Opp, hyperploid.

hypopodium (hī'pöpō'dium) n. [Gk. hypo, under; podion, little foot.]
Basal portion of a leaf, including stalk; style of carpel.

hypoproct (hī'pöprŏkt) n. [Gk. hypo, under; proktos, anus.] Medial prolongation of terminal abdominal segment beneath the anus, in Diplopoda and some Insecta.

hypopteron (hî'pŏp'tërŏn) n. [Gk. hypo, under; pteron, feather.]

Axillary feather in birds.

hypoptilum (hī'pŏp'tīlūm) n. [Gk. hypo, under; ptilon, down.] The aftershaft, q.v.

hypopus (hī'popus) n. [Gk. hypo, under; pous, foot.] Cyst-like stage of Tyroglyphidae.

hypopyge (hī'popī'jē) n. [Gk. hypo, under; pyge, rump.] Clasping organ of male dipterous insect; also hypopygium.

hyporhachis (hī'pörā'kĭs) n. [Gk. hypo, under; rhachis, spine.] The stem of aftershaft of a feather.

also hyporachis.

hyposkeletal (hī'pöskēl'étăl) a. [Gk. hypo, under; skeletos, hard.] Lying beneath or internally to endoskeleton.

hyposphene (hī'pösfēn) n. [Gk. hypo, under; sphen, wedge.] A wedge-shaped process on neural arch of vertebra of certain reptiles which fits into hypantrum.

hypostasis (hipŏs'tăsis) n. [Gk. hypo, under; stasis, standing.] Sediment or deposit, as of blood; recessiveness of non-allelomorphic

characters; cf. epistasis.

hypostatic (hi'pöstät'ik) a. [Gk. hypo, under; stasis, standing.] Appl. the recessive of two characters whose genes are not at the same time allelomorphs; exhibiting condition of hypostasis; cf. epistatic; pert. a sediment.

hypostereom (hi'pöstěr'eŏm) n. [Gk. hypo, under; stereos, solid.] The third or inner layer of thecal plates, of Cystidea; the inner layer of integument, of Crinoidea.

hypostoma (hīpōs'tömā) n. [Gk. hypo, under; stoma, mouth.] The fold bounding posterior margin of oral aperture in crustaceans; labrum or median preoral plate in trilobites; oral projection or manubrium of a hydrozoan; anteroventral region of insect head; ventral mouth part of ticks; hypostome.

hypostomatic (hī'pöstŏmāt'ik) a. [Gk. hypo, under; stoma, mouth.] Situated beneath stomata of plant epidermis; appl. chamber or cavity.

hypostomatous (hī'pöstŏm'ātūs) a. [Gk. hypo, under; stoma, mouth.] Having stomata on under surface; having mouth placed on lower or ventral side.

hypotarsus (hī'pötâr'sŭs) n. [Gk. hypo, under; L. tarsus, ankle.] The calcaneum of a bird.

hypothalamus (hī'pöthāl'āmŭs) n. [Gk. hypo, under; thalamos, chamber.] Region below thalamus, and structures forming greater part of floor of third ventricle.

hypothallus (hī'pöthāl'ūs) n. [Gk. hypo, under; thallos, young shoot.] The marginal hyphal outgrowth; a membranous case bearing sporangia.

hypotheca (hīpöthē'kā) n. [Gk. hypo, under; theke, cup.] Theca covering hypocone in Dinoflagellata; younger half of frustule in diatoms.

hypothecium (hīpöthē'siūm, -shīūm)
n. [Gk. hypo, under; theke, cup.]
The layer of dense hyphal threads below the thecium in lichens; upper layer of ascoma.

hypothenar (hī'pöthěn'ăr) a. [Gk. hypo, under; thenar, palm of hand.] Pert. the prominent part of palm of hand above base of little finger.

hypothetical units, — the ultimate component parts of protoplasm;

ultracellular units ranking between the molecule and the cell; also called variously, physiological units, pangens, gemmules, biophores, bioblasts, somacules, idiosomes, plasomes, micellae, plastidules, inotagmata, idioblasts, biogens, gemmae, microzymas, genes, gens, primordia.

hypotonic (hī'pötön'ik) a. [Gk. hypo, under; tonos, tension.] Having a lower osmotic pressure than that of another fluid, as of serum.

hypotrematic (hi'pötrēmāt'īk) a. [Gk. hypo, under; trema, pore.] Appl. the lower lateral bar of branchial basket of lamprey.

hypotrichous (hīpŏt'rīkŭs) a. [Gk. hypo, under; thrix, hair.] Having cilia mainly restricted to under surface; with deficient hair.

hypotrochanteric (hi'pötrők'ántěr'ik) a. [Gk. hypo, under; trochanter, runner.] Beneath the trochanter.

hypotympanic (hi'pötimpăn'ik) a. [Gk. hypo, under; L. tympanum, drum.] Situated below the tympanum; pert. quadrate.

hypotype (hi'pötip) n. [Gk. hypo, under; typos, pattern.] Any specimen described or figured in order to amplify or correct the identification of a species; plesiotype, q.v.

hypovalve (hī'pövālv) n. [Gk. hypo, under; L. valva, fold.] The antapical part of envelope in certain Dinoflagellata; hypocone.

hypoxanthine (hi'pözăn'thīn) n. [Gk. hypo, under; xanthas, yellow.] A crystalline nitrogenous substance found in glandular and muscle tissue and in some seeds; C₆H₄ON₄.

hypozygal (hi'pözi'gäl) n. [Gk. hypo, under; zygon, yoke.] Lower ossicle of a syzygial pair bearing no pinnule.

hypsodont (hip'södönt) a. [Gk: hypsos, height; odous, tooth.] Pert. or designating teeth with high crowns and short roots; hypselodont.

hypsophyll (hip'söfil) n. [Gk. hypsi, high; phyllon, leaf.] Any leaf beneath the sporophylls; bract, bracteole. hypural (hľpū'răl) a. [Gk. hypo, under; oura, tail.] Pert. a bony structure, formed by fused haemal spines of last few vertebrae, which supports caudal fin in certain fishes.

hysteranthous (histërăn'thus) a. [Gk. hysteros, coming after; anthos, flower.] Leafing after appearance of flowers.

hysterectomy (his'tĕrĕk'tömi) n. [Gk. hystera, womb; ek, out; temnein, cut.] Excision of the uterus.

hysteresis (hīstěr'ēsīs) n. [Gk. hysteresis, late arrival.] Lag in one of two associated processes or phenomena; lag in adjustment of external form to internal stresses, as in chromosome during spiralisation.

hysterogenic (hĭs'tëröjčn'ĭk) a. [Gk. hysteros, later; genos, birth.] Of later development or growth.

hysterosoma (his'tëröso'mă) n. [Gk. hysteros, after; soma, body.] Part of body posterior to proterosoma and comprising metapodosoma and opisthosoma in Acarina.

hysterotely (his'tëröt'eli) n. [Gk. hysteros, after; telos, completion.] The retention or manifestation of larval characters in pupa or imago, or of pupal characters in imago; metathetely, opp. prothetely.

hyther (hīth'ēr) n. [Gk. hy(dor), water; ther(me), heat.] Combined effect of moisture and temperature

on an organism.

I

- ichnite (ik'nīt) n. [Gk. ichnos, track.] A fossil footprint; ichnolite.
- ichthyic (ik'thiik) a. [Gk. ichthys, fish.] Pert. or characteristic of fishes; ichthyoid.
- ichthyodorulite (ik'thiödŏr'ūlīt) n. [Gk. ichthys, fish; dory, spear; lithos, stone.] A fossil dermal or fin spine of fish.

ichthyoid,-ichthyic.

ichthyolite (ik'thiölit) n. [Gk. ichthys, fish; lithos, stone.] A fossil fish or part of one.

ichthyology (ikthiol'oji) n. [Gk. *ichthys*, fish; *logos*, discourse.] The study of fishes.

ichthyopterygia (ik'thĭŏptërij'iă) n. plu. [Gk. ichthys, fish; pteryx, wing or fin.] Paired fish fins.

iconotype (īkō'nötīp) n. [Gk. eikon, image; typos, pattern.] Representation, drawing or photograph, of a type.

icosandrous (ī'kosan'drus) a. [Gk. eikosi, twenty; aner, man.] Having

twenty or more stamens.

icotype (ī'kōtīp) n. [Gk. eikon, image; typos, pattern.] A representative specimen used for identification of a species.

id (id) n. [Gk. idios, distinct.] A hypothetical structural unit; the chromomere.

idant (Idant) n. [Gk. idios, distinct.] A unit resulting from an aggregation of ids; the chromosome.

identical (īděnt'íkăl) a. [L. idem, the same.] Appl. progeny having the same genes, as monozygotic twins.

ideoglandular (īdēöglān'dūlār) a. [Gk. idein, to see; L. glandula, small acorn.] Pert. glandular activity induced by a mental image.

ideomotor (idēömō'tŏr) a. [Gk. idein, to see; L. movere, to move.]
Pert. unwilled movement in response to a mental image.

ideotype (īdēötīp) n. [Gk. idein, to see; typos, pattern.] Specimen, other than a topotype, named by the author who has described the species to which it belongs.

ideovascular (idēövăs'kūlār) a. [Gk. idein, to see; L. vasculum, small vessel.] Pert. circulatory changes induced by a mental image.

idioandrosporous (ĭd'iöăndröspō' rūs) a. [Gk. idios, distinct; aner, male; sporos, seed.] With androspores formed on filaments that do not bear oogonia.

idiobiology (Id'iöbīŏl'oji) n. idios, personal; bios, life; logos, discourse.] Biology of an individual

organism; autobiology.

idioblast (ĭd'Iöblăst) n. [Gk. idios, distinct; blastos, bud.] A hypothetical unit, q.v.; plant cell containing oil, gum, calcium carbonate, or other product and which differs from the surrounding parenchyma.

idiocalyptrosome (id'iökălip'trösōm) n. [Gk. idios, distinct; kalyptra, covering; soma, body.] Outer zone derived from idiosphaerosome in

sperm cells.

- idiochromatin (ĭd'ĭökrō'mătĭn) n. idios. [Gk. distinct; chroma, colour.] Temporarily dormant generative chromatin; cf. trophochromatin.
- idiochromidia (ĭd'iökrōmĭd'iă) n. plu. [Gk. idios, distinct; chroma, colour.] Sporetia; generative chromidia; cf. trophochromidia.
- idiochromosome (ĭd'iökrō'mösōm) n. idios, distinct; chroma. colour; soma, body.] chromosome.
- idiocryptosome (ĭd'ĭökrĭp'tösōm) n. [Gk. idios, distinct; kryptos, hidden; soma, body.] Inner zone derived from idiosphaerosome in sperm cells.
- idiomuscular (ĭd'ĭömŭs'kūlár) [Gk. idios, peculiar; L. musculus, muscle.] Appl. contraction of a degenerated muscle artificially stimulated.
- idiophthartosome (Id'iofthar'tösom) n. [Gk. idios, distinct; phthartos, transitory; soma, body.] idiozome remnant.
- idioplasm (ĭd'iöplazm) n. [Gk. idios, distinct; plasma, mould.] Chromatin; the generative or germinal part of a cell; cf. trophoplasm.
- idiosoma (ĭdĭösō'mā) n. [Gk. idios, distinct; soma, body.] The body. prosoma and opisthosoma Acarina.
- idiosome (id'iosom) n. [Gk. idios, distinct; soma, body.] A hypothetical unit, q.v.; sphere or region

of cytoplasm differing in viscosity from remainder of cell and surrounding the centriole or centrosome; idiozome, q.v.

idiosphaerosome (ĭd'Iösfē'rösōm) n. [Gk. idios, distinct; sphaira, globe; soma, body.] Acrosome; central

granule of idiosphaerotheca.

idiosphaerotheca (ĭd'ĭösfē'röthē'kă) n. [Gk. idios, distinct; sphaira, globe; theke, case.] Acroblast; vesicle containing acrosome sperm cells.

idiothalamous (ĭd'iöthăl'ămŭs) a. [Gk. idios, distinct; thalamos. Appl. lichens in which various parts are differently coloured

from thallus.

idiothermous (ĭd'iöther'mŭs) a. [Gk. idios, personal; thermos, hot.] Warm-blooded; homoiothermal.

- idiotype ($id'i\ddot{o}t\ddot{i}p$) n. [Gk. idios, personal; typos, pattern.] vidual genotype.
- idiovariation,—mutation, q.v. idiozome (ĭd'ĭözōm) n. [Gk. idios, distinct; zoma, girdle.] In spermatogenesis a separated portion of archoplasm which ultimately becomes head-cap of spermatozoon;
- cytes containing the centrioles. I-disc,—singly refracting or isotropic band in myofibrillae.

centrotheca; a cell-body of auxo-

- idorgan (ĭd'ôrgăn) n. [Gk. idios, distinct; organon, instrument.] A purely morphological multicellular unit which does not possess the features of a soma.
- ileocaecal (ĭl'eösē'kăl) a. [L. ileum, groin; caecus, blind.] Pert. ileum and caecum; appl. fossae, folds.
- ileocolic (ĭl'eökŏl'ĭk) a. [L. ileum, groin; Gk. kolon, colon.] Pert. ileum and colon; appl. artery, lymph-glands.
- ileum (il'eum) n. [L. ileum, groin.] Lower part of small intestine.
- iliac (ĭl'ĭăk) a. [L. ilia, flanks.] Pert. ilium, a pelvic bone; appl. artery, fossa, furrow, tuberosity, vein, etc.; appl. muscle, iliacus, from upper part of iliac fossa to side of tendon of psoas major; appl. processes of

ischiopubic plate; forming base for pelvic fins.

iliocaudal (il'iökô'dăl) a. [L. ilia, flanks; cauda, tail.] Connecting ilium and tail; appl. muscle.

iliococcygeal (ĭl'ĭökŏksĭj'ëăl) a. [L. ilia, flanks; Gk. kokkyx, cuckoo.] Pert. ileum and coccyx; appl. a muscle.

iliocostal (ĭl'ĭokŏs'tăl) a. [L. ilia, flanks; costa, rib.] In region of ilia and ribs; appl. several muscles.

iliofemoral (ĭl'ĭöfĕm'örăl) a. [L. ilia, flanks; femur, thigh.] Pert. ilium and femur; appl. a ligament.

iliohypogastric (ĭl'ĭöhī'pögăs'trĭk) a. [L. ilia, flanks; Gk. hypo, under; gaster, stomach.] Pert. ilium and lower anterior part of abdomen; appl. a nerve.

ilio-inguinal (ĭl'ĭöĭng'gwĭnăl) a. [L. ilia, flanks; inguen, groin.] In the region of ilium and groins; appl. a

nerve.

ilio-ischiadic (ĭl'ĭoĭskĭăd'ĭk) a. [L. ilia. flanks; Gk. ischion, hip.] Appl. fenestra between ilium and ischium when these are fused at both ends.

iliolumbar (ĭl'ĭölŭm'băr) a. [L. ilia, flanks; lumbus, loin.] In region of ilium and loins; appl. artery,

ligament, vein.

iliopectineal (ĭl'ĭöpĕktĭn'eăl) a. [L. ilia, flanks; pecten, crest.] Appl. an eminence marking point of union of ilium and pubis; appl. fascia.

iliopsoas (ĭl'ĭösō'ăs, -psō'ăs) n. ilia, flanks; Gk. psoa, loins.] Iliacus and psoas major considered as one muscle.

iliotibial (ĭl'ĭötĭb'ĭăl) a. [L. ilia, flanks; tibia, shin.] Appl. tract or band of muscle at lower end of

iliotrochanteric (ĭl'ĭötrōkăntěr'ĭk) a. [L. ilia, flanks; Gk. trochanter, runner.] Uniting ilium and trocharter of femur; appl. a ligament.

ilium (il'ium) n. [L. ilium, flank.] That part of hip-bone supporting the flank; dorsal bone of pelvic arch.

illuvial (ĭlū'vlăl) a. [L. in, into;

luere, to wash.] Appl. lower soil layers or B horizon.

imaginal (Imāj'īnāl) a. [L. imago, image.] Pert. an imago; appl. larval discs, patches of cells from which new organs develop.

imago (ĭmā'gŏ) n. [L. imago, image.] The last or adult stage n insect metamorphosis; the perfect

insect.

imbibition (im'bibish'un) n. [L. in, into; bibere, to drink.] Absorption of fluids, as of water by roots.

imbricate (ĭm'brikāt) a. [L. imbricare, to tile.] Having parts overlying each other like roof-tiles; appl. scales, plates, bud-scales, bracts.

imitative (ĭm'ĭtātĭv) a. [L. imitari, to imitate.] Appl. form, structure, habit, colouring, etc., assumed for protection or aggression.

immaculate (imak'ūlāt) a. [L. in, not; macula, spot.] Without spots or marks of different colour.

immune body,—heat-stable antibody or lysin; amboceptor, q.v.

immunise (ĭmūnīz', ĭm'ūnīz) v. [L. immunis, free.] To render invulnerable to a toxin, usually by injecting the toxin in small quantities at short intervals, without appearance of severe symptoms.

immunity (imū'niti) n. [L. immunis, An organism's resistance, natural or acquired, to the onset of pathological conditions from infection, natural or artificial, by microorganisms or their products.

impar (im'pâr) a. [L. impar, unequal.] Not paired; not existing

in pairs; azygous.

imparidigitate (impăr'idij'itāt) a. [L. impar, unequal; digitus, finger.] Having an odd number of digits.

imparipinnate (impăr'ipin'āt) a. [L. impar, unequal; pinna, wing.] Unequally pinnate; pinnate with an odd terminal leaflet.

imperfect (impër'fěkt) a. [L. imperfectus, unfinished.] Incomplete; appl. fungi lacking the sexual spore stage.

- imperforate (imper'förāt) a. [L. in, not; per, through; foratus, bored.]
 Not pierced; appl. foraminiferous shells without fine pores in addition to principal opening.
- impervious (Imper'vius) a. [L. in, not; pervius, passable.] Not permeable; appl. nostrils with septum between nasal cavities.
- implant (im'plant) n. [L. in, into; plantare, to plant.] An organ or part transplanted to an abnormal position; a graft.
- implantation cone,—cone of origin, q.v.
- implex (im'pleks) n. [L. in, into; plicare, to fold.] Endoplica or infolding of integument for muscle attachment in insects.
- importation (impôrtā'shun) n. [L. importare, to carry into.] Ingestion by sinking of food into protoplasm of captor, as in certain protozoa.
- impregnation (im'pregnā'shun) n. [L. impraegnare, to fertilise.] Transference of spermatozoa from male to body of female; insemination.
- impressio (impres'iō) n. [L. impressio, impression.] Impression or concavity in one organ or structure where in contact with another, as of surface of liver in contact with stomach, etc.
- impuberal (impū'berāl) a. [L. impubes, under age.] Prepubertal; sexually immature.
- impulse (Im'puls) n. [L. impulsus, driven.] Self-propagated disturbance induced by excitation.
- inantherate (ĭnăn'therāt) a. [L. in, not; Gk. antheros, flowering.] Without anthers; anantherous.
- inarticulate (inartik'ūlāt) a. [L. in,
 not; articulatus, jointed.] Not
 segmented; not jointed.
- inaxon (ľnāk'sŏn) n. [Gk. is, fibre; axon, axis.] A nerve-cell with axiscylinder branching at a distance from it.
- inbreeding,—breeding through a succession of parents belonging to the same stock, or very nearly related.

- Inca bones, distinct portions of interparietal, found in skulls of former Peruvians; os interparietale.
- incaliculate (înkălik'ūlāt) a. [L. in, not; caliculus, small flower-cup.] Wanting a calicle.
- incasement theory, preformation theory, q.v.
- incisal (insi'săl) a. [L. incidere, to cut into.] Cutting, as edge of a tooth.
- incised (insizd') a. [L. incisus, cut
 into.] With deeply notched margin.
- incisiform (Insī'zīfôrm) a. [L. incisus, cut into; forma, shape.]
 Incisor-shaped.
- incisive (insī'sīv) a. [L. ncisus, cut into.] Pert. or in region of incisors; appl. bones, foramina, fossa.
- incisor (Insi'sŏr) a. [L. incisus, cut into.] Adapted for cutting—appl. mammalian premaxillary teeth. n. A crest or ridge of palatine process of maxilla.
- incisura (ĭnsīsū'ră) n. [L. incidere, to cut into.] Notch, depression, or indentation, as in bone, stomach, liver. etc.
- included (inklood'ëd) a. [L. includere, to shut in.] Having stamens and pistils not protruding beyond corolla; not exserted.
- inclusion bodies,—intracellular particles, as pigment granules, mitochondria, Golgi bodies, microsomes, viruses, etc.
- incomplete metamorphosis, insect metamorphosis in which young are hatched in general adult form and develop without quiescent stage.
- incongruent (inköng'grooënt) a. [L. incongruens, not suiting.] Not suitable or fitting; appl. surface of joints which do not fit properly.
- incoordination (în'köôr'dinā'shūn) n.
 [L. in, not; cum, together; ordo, order.] Want of co-ordination; irregularity of movement due to loss of muscle control.
- incrassate (Inkräs'āt) a. [L. incrassare, to thicken.] Thickened; becoming thicker.
- incretion (inkrē'shun) n. [L. in,

- into; cretus, separated.] Internal secretion.
- incrustation (ĭnkrŭstā'shŭn) n. [I.. in, into; crusta, shell.] Fossilisation by encasement in mineral substance.
- incubation (in'kūbā'shūn) n. [L. incubare, to lie on.] The hatching of eggs by means of heat, natural or artificial; period between infection and appearance of symptoms induced by parasitic organisms.
- incubatorium (in'kūbătō'riŭm) n. [L. incubare, to lie on.] Temporary pouch surrounding mammary area, in which egg of Echidna is hatched.
- incubous (In'kūbŭs) a. [L. incubare, to lie on.] Appl. leaves so arranged that the base of each is covered by upper portion of next lower.
- incudal (ĭn'kūdăl) a. [L. incus, anvil.] Pert. the incus; appl. fold, fossa.
- incudate (ĭn'kūdāt) a. [L. incus, anvil.] Appl. type of rotifer mastax with large and hooked rami and reduced mallei.
- incumbent (inkům'bënt) a. [L. incumbere, to lie upon.] Lying upon; bent downwards to lie along a base; appl. cotyledons so folded that flat sides are next radicle; appl. hairs or spines applied lengthwise to their base; appl. insect wings resting on abdomen.
- incurrent (inkŭr'ënt) a. [L. in, into; currere, to run.] Leading into; afferent; appl. ectoderm-lined canals which admit water, in sponges; appl. inhalant siphons of molluscs.
- incurvate (ĭnkŭr'vāt) a. [L. incurvus, bent.] Curved inwards or bent back.
- incurvation (in'kūrvā'shūn) n. [L. incurvare, to curve.] The doubling back on itself of a structure or organ, as of a spirochaete about to divide.
- incus (ĭn'kŭs) n. [L. incus, anvil.]
 Part of a rotifer mastax; the anvilshaped ear ossicle of mammals.
- indeciduate (in'dësid'ūāt) a. [L. in,

not; decidere, to fall down.] Non-caducous; with maternal part of placenta not coming away at birth.

- indeciduous (ĭn'dësid'ūŭs) a. [L. in, not; decidere, to fall down.] Persistent; not falling off at maturity; everlasting; evergreen.
- indefinite (indef'init) a. [L. in, not;
 definitus, limited.] Not limited; not
 determinate; of no fixed number;
 racemose, q.v.
- indehiscent (ĭn'dēhīs'ent) a. [L. in, not; dehiscens, gaping.] Not splitting at maturity; appl. certain fruits.
- indeterminate growth,—growth of stem, branch or shoot not limited or stopped by development of a terminal bud; indefinite prolongation and subdivision of an axis.
- indeterminate inflorescence, growth of a floral axis by indefinite branching because unlimited by development of a terminal bud.
- index,—the forefinger or digit next to the thumb; a number or formula expressing ratio of one quantity to another.
- indicators,—species characteristic of climatic, soil and other conditions in a particular region or habitat; dominant species in a biotope.
- indigenous (ĭndĭj'ënŭs) a. [L. indigena, native.] Belonging to the locality; not imported; native.
- individual (ĭn'divid'ūāl) a. [L. in, not; dividuus, divisible.] Pert. a single example or unit, as individual variations of colour. n. A person or zooid of distinctive function of a hydrozoan colony.
- individualism,—symbiosis in which the two parties together form what appears to be a single organism.
- individuation,—development of interdependent functional units, as in colony formation; organisation of morphogenetic processes; regional or tissue differentiation; process of developing into an individual.
- indole-acetic acid,—see heteroauxin.
 induced movement, movement
 dictated and influenced by external
 stimulus, as plant curvature.

- induction (Induk'shun) n. [L. inducere, to lead in.] Act or process of causing to occur; process whereby a cell or tissue influences neighbouring cells or tissues; lowering by one reflex of the threshold of another, spinal induction.
- inductive stimulus,—an external stimulus which influences growth or behaviour of an organism.
- indumentum (In'dūměn'tūm) n. [L. indumentum, covering.] The plumage of birds; a hairy covering.
- induplicate (indu'plikāt) a. [L. in, in; duplex, double.] In vernation having bud-leaves bent or rolled without overlapping; in aestivation having bud sepals or petals folded inwards at points of contact.
- induplicative (indū'plikātiv) a. [L. in, in; duplex, double.] Appl. vernation or aestivation with induplicate foliage or floral leaves respectively.
- indusial (ĭndū'zĭāl) a. [L. induere, to put on.] Containing larval insect cases, as certain limestones; pert. the indusium.
- indusiate (ĭndū'ziāt) a. [L. induere, to put on.] Having an enveloping case, appl. insect larvae; having an indusium.
- indusiform (ĭndū'zĭfôrm) a. [L. induere, to put on; forma, shape.]
 Resembling an indusium.
- indusium (indū'ziŭm) n. [L. induere, to put on.] An outgrowth of plant epiderm covering and protecting a sorus; an insect larva case; the supracallosal gyrus of the rhinencephalon, indusium griseum.
- induviae (Indū'viē) n. plu. [L. induere, to put on.] Scale-leaves; leaves which remain attached to stem after withering.
- induviate (Indū'vĭāt) a. [L. induere, to put on.] Covered with scaleleaves or induviae.
- inequilateral (iněk'wilăt'êrăl) a. [L. in, not; aeguus, equal; latus, side.] Having two sides unequal; having unequal portions on either side of a line drawn from umbo to gape of a bivalve shell.

inequilobate (iněk'wilō'bāt) a. [L. in, not; aequus, equal; lobus, lobe.] With lobes of unequal size.

inequivalve (ĭněk'wĭvălv) a. [L. in, not; aequus, equal; valvae, folding doors.] Having two valves of shell unequal; appl. molluscs.

- inerm (inerm') a. [L. inermis, unarmed.] Without means of defence and offence; without spines; inermous.
- inert (inert') a. [L. iners, inactive.] Physiologically inactive; appl. heterochromatic region of chromosome with paucity of active genes.
- inferior (ĭnfe'rĭŏr) a. [L. inferior, lower.] Appl. lower placed of two, farther down axis; growing or arising below another organ.
- inferoanterior (ĭn'feröantē'riŏr) a. [L inferus, beneath; anterior, in front.] Below and in front.
- inferobranchiate (ĭn'fëröbrăng'kiāt)
 a. [L. inferus, beneath; Gk. brangchia, gills.] With gills under margin of mantle, as in certain molluscs.
- inferolateral (in'fërölăt'ërăl) a. [L. inferus, beneath; latus, side.]
 Below and at or towards the side.
- inferomedian (ĭn'ferome'dian) a. [L. inferus, beneath; medius, middle.] Below and about the middle.
- inferoposterior (In'féroposte'rior) a. [L. inferus, beneath; posterior, behind.] Below and behind.
- inferradial (ĭn'fērrā'diāl) n. [L. inferus, beneath; radius, radius.]
 Lower part of transversely bisected radials of certain fossil crinoids.
- inflected (inflek'têd) a. [L. inflectere, to bend in.] Curved or abruptly bent inwards or towards the axis; inflexed.
- inflorescence (In'flöres'ens) n. [L. inflorescere, to begin to blossom.] A flowering or putting forth blossoms; method in which flowers are arranged on an axis.
- influents (in'flooënts) n. plu. [L. influere, to flow into.] The animals present in a plant community, or those primarily dependent and acting upon the dominant plant species. infra-axillary (in'fraak'silari) a. [L.

- infra, below; axilla, armpit.] Branching off below the axil.
- infrabasal (ĭn'frābā'săl) n. [L. infra, below; basis, base.] One of a series of plates, perradial in position, below the basals in crinoids.
- infrabranchial (ĭn'frābrăng'kĭāl) a.
 [L. infra, below; branchiae, gills.]
 Below the gills; appl. part of pallial chamber.
- infracentral (in'frăsĕn'trăl) a. [L. infra, below; centrum, centre.]
 Below a vertebral centrum.
- infraclavicle (in'frāklāv'īkël) n. [L. infra, below; clavicula, little key.] Membrane bone occurring in pectoral girdle of some fishes.
- infraclavicular (In'frāklāvik'ūlār) a. [L. infra, below; clavicula, small key.] Beneath the clavicula; appl. branches of brachial plexus; appl. fossa or triangle between deltoid and pectoralis major.
- infracortical (in'frăkôr'tikăl) a. [L. infra, below; cortex, bark.] Beneath the cortex.
- infracostal (ĭn'frākŏs'tăl) a. [L. infra, below; costa, rib.] Beneath the ribs; appl. muscles.
- infradentary (In'fraden'tari) a. [L. infra, below; dens, tooth.] Beneath the dentary bone.
- infraglenoid (In'frăglē'noid) a. [L. infra, below; Gk. glene, socket; eidos, like.] Below glenoid cavity; appl. a tuberosity.
- infrahyoid (Infrahi'oid) a. [L. infra, below; Gk. hyoeides, Y-shaped.] Beneath the hyoid; appl muscles.
- infralabial (in'frălă'biăl) a. [L. infra, below; labium, lip.] Beneath the lower lip.
- inframarginal (inframar'jīnal) a. [L. infra, below; margo, margin.] Under the margin, or marginal structure; appl. a cerebral convolution; appl. certain plates on carapace of Chelonia below marginals; appl. lower of two series of plates round margin of stelleroid arms and discs.
- inframaxillary (ĭn'frămăksĭl'ărī) a. [L. infra, below; maxilla, jaw.] Beneath maxilla; appl. nerves.

- infranasal (ĭn'frănā'zăl) n. [L. infra, below; nasus, nose.] An additional nasal element in some Theromorpha.
- infraorbital (in'fraôr'bital) a. [L. infra, below; orbis, circle.] Beneath the orbit; appl. artery, canal, foramen, groove, nerve, glands, etc.
- infrapatellar (ĭn'frăpătël'ăr) a. [L. infra. below; patella, knee-cap.]
 Appl. pad of fat beneath patella.
- infrarostral (in'frărŏs'trăl) a. [L. infra, below; rostrum, snout.]
 Beneath a rostrum; appl. paired cartilages, derived from Meckel's cartilage, of lower part of suctorial mouth of tadpoles.
- infrascapular (ĭn'frăskăp'ūlăr) a. [L. infra, below; scapula, shoulder-blade.] Beneath the scapula; appl. artery.
- infraspecific (in'frăspesif'îk) a. [L. infra, below; species, particular kind.] Pert. a subdivision of a species, as subspecies and varieties.
- infraspinatous (ĭn'frăspinā'tŭs) a. [L. infra, below; spina, spine.] Beneath the spine; beneath scapular spine; appl. muscle, fossa; infraspinous.
- infrastapedial (In'frastape'dial) a. [L. infra, below; stapes, stirrup.] Beneath stapes of ear; appl. part of columella.
- infrasternal (in'fraster'nal) a. [L. infra, below; sternum, breast-bone.]
 Below the breast-bone; appl. notch superficially at lower end of sternum.
- infratemporal (in'fratem'poral) a. [L. infra, below; tempora, temples.] Beneath the temporal bone; appl. a crest and fossa.
- infratrochlear (in'frătrok'lëăr) a. [L. infra, below; trochlea, pulley.] Beneath the trochlea; appl. a nerve given off from nasociliary nerve.
- infructescence (in'früktes'ëns) n.
 [L. in, into; fructus, fruit.] An inflorescence matured into a fruit; a composite or confluent fruit.

- infundibular (In'fundib'ulăr) a. [L. infundibulum, funnel.] Funnel-shaped; appl. an abdominal muscle; appl. corolla; infundibuliform; choanoid, q.v.; pert. infundibulum.
- infundibulin (Infundib'ūlin) n. [L. infundibulum, funnel.] An extract of posterior pituitary lobe causing decrease in renal water excretion; antidiuretin; infundin.
- infundibulum (in'fūndīb'ūlūm) n. [L. infundibulum, funnel.] Any funnel-shaped organ or structure; appl. part of brain, of ethmoid bone, of right ventricle, etc.; a cephalopod siphon; part of bird's oviduct; flattened stomach-like cavity of ctenophore; septal funnel in Scyphozoa.

infuscate (ĭnfus'kāt) a. [L. in, into; fuscus, dark.] Tinged to appear

dark, as insect wings.

- infusoriform (In'fūsō'rifôrm) a. [L. infusus, poured into; forma, shape.] Resembling an infusorian; appl. embryonic forms of Coelentera; appl. male form of Dicyemidae.
- infusorigen (in'fūsō'rijēn) n. [L. in-fusus, poured into; genos, off-spring.] A gastrula-like phase in development of certain Mesozoa.

ingest (injest') v. [L. ingestus, taken in.] To convey food material into the alimentary canal or food-cavity.

- ingesta (ĭnjĕs'tă) n. plu. [L. ingestus, taken in.] The sum-total of substances taken in by the body. Opp. egesta.
- ingestion (injes'tion) n. [L. ingestus, taken in.] The swallowing or taking in of food-material.
- ingluvies (ingloov'iez) n. [L. ingluvies, crop.] The crop of a bird; a dilatation of oesophagus; the rumen.
- inguinal (Ing'gwinăl) a. [L. inguen, groin.] In region of groin.
- inguinal ring,—see abdominal ring. inguino-abdominal,—in region of abdomen and groin.
- inguino-crural,—in region of groin and leg.
- inhalant (Inhā'lant) a. [L. in, into; halare, to breathe.] Adapted for

inspiring or drawing in, as terminal pores of incurrent canals in sponges, or siphons in molluscs.

inhibition (in'hibish'ön) n. [L. in-hibere, to prohibit.] Prohibition, or checking, of an action or process.

- inhibitory (inhib'itori) a. [L. inhibere, to prohibit.] Appl. nerves which control movement or secretion.
- inion (ĭn'ĭŏn) n. [Gk. inion, back of head.] The external protuberance of occipital bone.
- initial (inish'ăl) n. [L. initium, beginning.] A cell which initiates differentiation of tissues, as in apical meristem, vascular cambium, etc.; histogen cell, primordial cell.
- ink sac,—in Sepia, a pear-shaped body in wall of mantle cavity which contains the ink gland, secreting a black substance, ink or sepia, ejection of which is a means of defence.
- innate (in'nāt) a. [L. in, in; gnatus, born.] Inherited; basifixed, appl. anther with filament attached only to base.
- innervation (ĭn'nërvā'shūn) n. [L. in, into; nervus, sinew.] Nervedistribution; vital nerve force.
- innidiation (in'nidiā'shūn) n. [L. in, into; nidus, nest.] Colonisation or development of cells or organisms in a part of the body to which they have been transferred by metastasis, q.v.
- innominate (Innom'ināt) a. [L. in, not; nomen, name.] Nameless; appl. various arteries and veins.
- innominate bone,—the hip-bone or lateral half of pelvic girdle; os coxae, os innominatum.
- innovation (in'övā'shun) n. [L. innovare, to renew.] A growth or shoot of mosses which develops into a new plant by dying-off of portion of parent-plant behind it; basal vegetative shoot of grasses.
- inocomma (Ĭn'ökom'ă) n. [Gk. is, fibre; komma, clause.] Portion of muscle fibril between telophragmata or Z discs; inokomma; sarcomere.

- inocular (ĭnŏk'ūlăr) a. [L. in, into; oculus, eye.] Appl. antennae inserted close to eye.
- inocyte (ľn'ösīt) n. [Gk. is, fibre; kytos, hollow.] Elongated cell of fibrous tissue.
- inogen (Ĭn'öjĕn) n. [Gk. is, fibre; genos, offspring.] A nitrogenous substance of muscle tissue.
- inophragma (Ĭn'öfrăg'mă) n. [Gk. is, fibre; phragma, fence.] The transverse membrane through adjacent myofibrillae; mesophragma and telophragma; M and Z lines bisecting A-and I-discs.
- inositol,—carbohydrate present in cells, the lipotropic anti-alopecia factor of bios, q.v.; C₆H₁₂O₆.
- inotagmata (in'otăg'mătă) n. plu. [Gk. is, fibre; tagma, arrangement.] Hypothetical units, q.v.
- inquiline (in'kwilin) n. [L. inquilinus, tenant.] Animal living in home of another and getting share of its food; partner in commensalism; an insect developing in gall produced by an insect of another species, being detrimental to the latter.
- inscriptions, tendinous, three fibrous bands crossing the rectus abdominis muscle.
- insectivorous (ĭn'sĕktīv'örŭs) a. [L. insectum, cut into; vorare, to devour.] Insect-eating; appl. certain animals and carnivorous plants.
- insemination (ĭn'sĕmĭnā'shūn) n. [1...
 in, in; seminatio, sowing.] The
 introduction of semen or spermatozoa into female genital tract.
- inserted (ĭnsĕr'tĕd) a. [L. in, in; serere, to join.] United by natural growth.
- insertion (inser'shun) n. [L. insertus, joined.] Point of attachment of organs, as of muscles, leaves; point on which force of a muscle is applied.
- insertional,—appl. translocation in which the portion between two breaks of a chromosome is transferred to a break in another chromosome; cf. shift.
- insessorial (ĭn'sĕsō'rĭăl) a. [L. in-

- sidere, to sit upon.] Adapted for perching.
- insistent (insis'tënt) a. [I.. insistere, to stand upon.] Appl. hind toe of certain birds whose tip only reaches the ground.
- insolation (ĭnsölā'shŭn) n. [L. in, into; sol, sun.] Exposure to sun's rays.
- inspiration (inspira'shun) n. [L. inspirare, to inhale.] The act of drawing air into the lungs; absorption of oxygen by plants.
- instaminate (Instăm'ināt) a. [L. in, not; stamen, thread.] Not bearing stamens.
- instar (in'stăr) n. [L. instar, form.] Insect at a particular stage between moults.
- insula (in'sūlă) n. [L. insula, island.] Island of Reil, a triangular eminence lying deeply in lateral fissure of temporal lobe; islet of Langerhans, q.v.; a blood island, q.v.
- insulin (ĭn'sūlin) n. [L. insula, island.] The anti-diabetic endocrine product of pancreas, formed in β-cells of islets of Langerhans; C₄₅H₆₉O₁₄N₁₁S. 3H₂O.
- integrifolious (integ'rifo'liŭs) a. [L. integer, whole; folium, leaf.] With entire leaves.
- integripallial (ĭntĕg'rīpāl'īāl), integripalliate (īntĕg'rīpāl'īāt) a. [L. integer, whole; pallium, mantle.] Having an unbroken pallial line; appl. shells of molluscs with small or no siphons.
- integument (Integ'ument) n. [L. integumentum, covering.] A covering, investing, or coating structure or layer; coat of ovule; integumentum.
- interacinous (In'tërăs'înŭs) a. [L. inter, between; acinus, grape.]
 Among alveoli of a racemose gland; interacinar.
- interalveolar (in'tërăl'vēölăr) a. [L. inter, among; alveolus, small cavity.]
 Among alveoli; appl. cell islets.
- interamb (Intëramb') n. [L. inter, between; ambulare, to walk.] Interambulacral area.

- interambulacral (in'tërămbūlā'krăl)
 a. [L. inter, between; ambulare, to
 walk.] Appl. area of echinoderm
 test between two ambulacral areas.
 n. A plate of that area.
- interambulacrum (in'tërambula'krum) n. [L. inter, between;
 ambulare, to walk.] The area
 between two ambulacral areas.
- interarticular (ĭn'tërârtik'ūlăr) a. [L. inter, between; articulus, joint.] Between articulating parts of bones; appl. certain ligaments and fibrocartilages.
- interatrial (in'tërā'triāl) a. [L. inter, between; atrium, hall.] Appl. groove separating the two atria of the heart.
- interauricular (ĭn'tërôrĭk'ūlăr) a. [L. inter, between; auricula, little ear.]
 Between auricles of heart.
- interaxillary (in'tërāk'silāri) a. [L. inter, between; axilla, armpit.] Placed between the axils.
- interbrachial (ĭn'tërbrā'kĭăl) a. [L. inter, between; brachium, arm.] Between arms, rays, or brachial plates.
- interbranchial (in'tërbrang'kial) a. [L. inter, between; branchiae, gills.] Appl. septum between successive gill slits.
- interbreed (in'tërbred') v. [L. inter, between; A.S. brod, brood.] To cross different varieties of plants or animals.
- intercalare (In'tërkăl'ărē) n. [L. intercalaris, inserted.] In many fishes and fossil amphibians, an additional element in the vertebra.
- intercalarium (ĭn'tërkălā'rĭūm) n.
 [L. intercalaris, inserted.] The third
 Weberian ossicle.
- intercalary (In'tërkāl'ārī, Inter'kālārī)

 a. [L. intercalaris, inserted.] Inserted; additional; appl. meristematic layers between masses of permanent tissue; appl. growth elsewhere than at growing point; appl. veins between main veins of insect wings; appl. plates in Dinoflagellata; appl. bands in diatoms; appl. cartilage between neural arches, interneural or interdorsal plate.

intercapitular (ĭn'tërkăpĭt'ūlăr) a.
[L. inter, between; capitulum, little head.] Between capitula; appl. veins of fingers and toes.

intercarotid (in'tërkărŏt'id) a. [L. inter, between; Gk. karos, deep sleep.] Between carotid arteries.

- intercarpal (în'tërkâr'păl) a. [L. inter, between; carpus, wrist.] Among or between carpal bones; appl. joints.
- intercarpellary (in'tërkâr'pëlări) a. [L. inter, between; Gk. karpos, fruit.] Between the carpels.
- intercartilaginous (In'tërkârtilăj'inŭs) a. [L. inter, between; cartilago, gristle.] Between cartilages.
- intercavernous (ĭn'tërkăv'ërnŭs) a.
 [L. inter, between; caverna, cavern.] Appl. sinuses connecting cavernous sinuses, part of ophthalmic veins.
- intercellular (in'tërsël'ūlăr) a. [L. inter, between; cellula, little cell.] Among or between cells, as spaces in meristem, biliary passages among liver-cells, plexus of dendrites between sympathetic ganglion cells, etc.
- intercentral (in'tërsën'trăl) a. [L. inter, between; centrum, centre.]
 Uniting, or between, two centra.
- intercentrum (in'tërsën'trum) n. [L. inter, between; centrum, centre.]
 A second central ring in an embolomerous vertebra.
- interchange,—mutual or reciprocal translocation, in chromosomes.
- interchondral (in'tërkôn'drăl) a. [L. inter, between; Gk. chondros, cartilage.] Appl. articulations and ligaments between costal cartilages.
- intercingular (in'tërsing'gūlăr) a. [L. inter, between; cingulum, girdle.] Appl. area of longitudinal groove between parts of a spiral girdle, in certain Dinoflagellata.
- interclavicle (in'tërklăv'ikl) n. [L. inter, between; clavicula, small key.] The episternum; a median ventral bone between clavicles.
- interclavicular (In'tërklăvik'ūlăr) a. [L. inter, between; clavicula, small key.] Between the clavicles; appl. a ligament.

- interclinoid (In'tërkli'noid) a. [L. inter, between; Gk. kline, bed; eidos, form.] Joining clinoid processes; appl. fibrous process or ligament.
- intercolumnar (ĭn'tërkölüm'năr) a.
 [L. inter, between; columna, column.] Between columnar structures, as certain abdominal muscle fibres.
- intercondyloid (ĭn'tërkŏn'dĭloid) a. [L. inter, between; Gk. kondylos, knob; eidos, form.] Between condyles; appl. an eminence of tibia, and fossae of femur and tibia.
- intercostal (ĭn'tërkŏs'täl) a. [L. inter, between; costa, rib.] Between the ribs, as arteries, glands, membranes, nerves, veins, muscles; between ribs of leaf, mericarp, etc.
- intercostobrachial (in'tërkos'tòbra'-kiăl) a. [L. inter, between; costa, rib; brachium, arm.] Appl. lateral branch of second intercostal nerve which supplies upper arm; intercostohumeral.
- intercoxal (In'tërkök'săl) a. [L. inter, between; coxa, hip.] Between the coxae or proximal limb-joints of arthropods; appl. plate, etc.
- intercrescence (in'terkres'ens) n.
 [L. inter, between; crescere, to grow.] A growing into each other, as of tissues.
- intercrural (in'tërkroo'răl) a. [L. inter, between; crus, leg.] Appl. intercolumnar tendinous fibres arching across external oblique muscles.
- intercuneiform (In'tërkūnē'Ifôrm) a.
 [L. inter, between; cuneus, wedge; forma, shape.] Connecting the three cuneiform bones of the ankle; appl. articulations and ligaments.
- interdeferential (In'tërdëfërën'shål)
 a. [L. inter, between; deferre, to carry down.] Between the vasa deferentia.
- interfascicular (ĭn'tërfăsīk'ūlăr) a. [L. inter, between; fasciculus, small bundle.] Situated between the fascicles or vascular bundles; appl. cambium.

- interfemoral (In'tërfëm'örăl) a. [I.. inter, between; femur, thigh bone.] Between the thighs.
- interference (in'tërfē'rĕns) n. [L. inter, between; ferire, to strike.] The lessened probability of crossing-over in the neighbourhood of a previous crossing-over.
- interfertile (in'tërfër'til) a. [L. inter, between; fertilis, fertile.] Able to interbreed.
- interfilamentar (in'tërfilamen'tăr) a.
 [L. inter, between; F. filament, from L. filum, thread.] Appl. junctions or horizontal bars connecting molluscan gill filaments.
- interfilar (in'tërfi'lar) a. [L. inter, between; filum, thread.] Appl. ground-substance of protoplasm, as opposed to reticulum.
- interfoliaceous (ĭn'tërfolĭa'shŭs) a.

 [L. inter, between; folium, leaf.]

 Situated or arising between two opposite leaves; interfoliar.
- interfrontal (in'terfrun'tal) n. [L. inter, between; frons, forehead.]
 An unpaired median bone between frontals and nasals in Eryops.
- interganglionic (in'tërgănggliŏn'īk)
 a. [L. inter, between; Gk. gangglion, little tumour.] Connecting two ganglia, as nerve cords or strands.
- intergeneric (ĭn'térjěnčr'īk) a. [L. inter, between; genus, kind.] Between genera; appl. hybridisation.
- intergenital (in'terjen'ital) a. [L. inter, between; genitus, produced.] Between the genitals; appl. certain echinoderm plates.
- interglacial (ĭn'tërglā'sĭāl) a. [L. inter, between; glacies, ice.] Appl. or pert. ages between glacial ages, particularly of the Pleistocene epoch.
- interglobular (interglob'ūlăr) a. [L. inter, between; globulus, small globe.] Appl. a series of spaces towards outer surface of dentine, due to imperfect calcification.
- intergular (in'tërgū'lăr) n. [L. inter, between; gula, gullet.] A paired or unpaired plate in front of gulars in Chelonia.

- interhyal (in'tërhī'āl) n. [L. inter, between; Gk. hyoeides, \tau-shaped.] A small bone between hyomandibular and rest of hyoid of some higher vertebrates.
- interkinesis (In'tërkinë'sis) n. [L. inter, between; Gk. kinesis, movement.] Interphase; resting stage between two mitotic divisions of a cell.
- interlamellar (ĭn'tërlămēl'ăr) a. [L. inter, between; lamella, thin plate.] Appl. vertical bars of tissue joining gill lamellae of molluscs; appl. spaces between lamellae or gills of agarics.
- interlaminar (ĭn'tërlăm'ĭnăr) a. [L. inter, between; lamina, thin plate.]
 Uniting laminae; between laminae.
- interlobar (ĭn'tërlō'băr) a. [L. inter, between; L.L. lobus, lobe.]
 Between lobes; appl. sulci and fissures dividing cerebral hemispheres into lobes.
- interlobular (ĭn'tërlöb'ūlăr) a. [L. inter, between; lobulus, small lobe.]
 Occurring between lobules; appl.
 kidney arteries, vessels of liver, etc.
- interlocular (ĭn'tërlŏk'ūlăr) a. [L. inter, between; loculus, compartment.] Between loculi.
- interloculus (ĭn'tërlŏk'ūlūs) n. [L. inter, between; loculus, compartment.] Space between two loculi.
- intermandibular (in'tërmăndib'ūlăr)
 a. [L. inter, between; mandibulum, jaw.] Between rami of mandibles.
- intermaxilla (in'tërmăksĭl'ă) n. [L. inter, between; maxilla, jaw.] Bone between maxillae; the premaxilla.
- intermaxillary (în'tërmăksîl'ărī) a. [L. inter, between; maxilla, jaw.] Between maxillae; pert. premaxillae; appl. gland in nasal septum of certain amphibians and reptiles.
- intermediary (in'tërmē'diări) a. [L. inter, between; medius, middle.] Acting as a medium; appl. nervecells receiving impulses from afferent cells and transmitting them to efferent cells.

intermediate (ĭn'tërmē'diāt) a. [L. inter, between; medius, middle.]
Occurring between two points or parts; appl. a nerve-mass, certain areas of brain, a disc in striated muscle, ribs, etc.

INT-

- intermediate host,—host intervening between two others in life-history of certain parasites, as Limnaea in life-history of Fasciola.
- intermedin (In'tërmē'din) n. [L. inter, between; medius, middle.]
 Hormone obtained from pars intermedia of the pituitary gland; B substance.
- intermedium (ĭn'tërmē'dĭŭm) n. [L. inter, between; medius, middle.] A small bone of carpus and tarsus.
- intermesenteric (In'tërmës'ëntër'îk, -mëz-) a. [L. inter, between; Gk. mesos, middle; enteron, gut.] Occurring between mesenteries; appl. spaces in sea-anemones.
- intermitotic (in'tërmitot'îk) n. [L. inter, between; Gk. mitos, thread.] A cell with individual life between mitoses causing its origin and division into daughter cells; cf. postmitotic.
- intermuscular (in'tërmus'kūlăr) a.
 [L. inter, between; musculus, muscle.] Between or among muscle fibres
- intermyotomic (in'tërmi'otŏm'îk) a. [L. inter, between; Gk. mys, muscle; tome, cutting.] Appl. vertebra formed of caudals of one somite and cranials of next posterior; cf. intrasegmental.
- internal (ĭntĕr'năl) a. [L. internus, within.] Located on inner side; nearer middle axis; located or produced within.
- internal secretion,—endocrine secretion; hormone.
- internal spiral,—coil within a single chromatid, as between prophase and anaphase.
- internasal (in'tërnā'zăl) a. [L. inter, between; nasus, nose.] Between nostrils; appl. plate, septum, gland.

- interneural (in'ternū'răl) a. [L. inter, between; Gk. neuron, nerve.] Between neural processes, arches or spines; appl. sharp bones attached to dorsal fin rays; appl. intercalary cartilages.
- interneurone (ĭn'tërnū'rŏn) n. [L. inter, between; Gk. neuron, nerve.] An internuncial neurone or relay cell.
- internodal (ĭn'tërnō'dăl) a. [L. inter, between; nodus, knot.] Pert. part between two nodes.
- internode (in'tërnōd) n. [L. inter, between; nodus, knot.] The part between two successive nodes or joints, as of plant stem, of medulated nerve fibre; non-genetic segment of a chromosome.

internodia (ĭn'tërnō'diă) n. plu. Phalanges.

- internum (Inter'num) n. [L. internus, inward.] Inner region or medulla of a mitochondrium or of Golgi apparatus.
- internuncial (in'tërnun'siăl) a. [L. inter, between; nuntius, messenger.] Intercommunicating, as paths of transmission or nerve fibres; appl. neurone interposed between afferent and efferent nerve cells, association neurone.
- interoceptor (ĭn'tërösĕp'tŏr) n. [L. internus, inside; capere, to take.] A receptor which receives stimuli from within the body; end-organ for visceral sensibility.

interocular (ĭn'tërök'ūlăr) a. [I.. inter, between; oculus, eye.] Placed between the eyes.

interoperculum (in'tërŏpěr'kūlūm) n.
[L. inter, between; operculum, lid.]
A membrane bone of operculum of
Teleostomi and Dipnoi, attached to
mandible; interopercle.

interoptic (ĭn'tërŏp'tĭk) a. [L. inter, between; Gk. optikos, pert. sight.]

Between optic lobes.

interorbital (in'tërôr'bītăl) a. [L. inter, between; orbis, circle.] Between the orbits; appl. septum of tropibasic skull; appl. sinus.

interosculant (ĭn'tërŏs'kūlănt) a. [L. inter, between; osculari, to kiss.]

Possessing characters common to two or more groups or species.

interosseous (ĭn'tërös'ëŭs) a. [L. inter, between; os, bone.] Occurring between bones; appl. arteries, membranes, muscles, nerves.

interparietal (In'tërpări'ëtăl) a. [L. inter, between; paries, wall.] In many vertebrates a bone arising between parietals and supraoccipital.

interpeduncular (in'tërpëdung'kūlăr)
a. [L. inter, between; pedunculus, little foot.] Appl. fossa between cerebral peduncles, and a ganglion.

interpetaloid (in'terpet'aloid) a. [L. inter, between; Gk. petalon, leaf; eidos, form.] Between petaloid areas of an echinoderm test.

interpetiolar (in'tërpët'iölăr) a. [L. inter, between; petiolus, little foot.] Situated between petioles or bases of opposite leaves.

interphalangeal (In'terfălăn'jeăl) a. [L. inter, between; Gk. phalangx, line of battle.] Appl. articulations between successive phalanges.

interphase (in'terfāz) n. [L. inter, between; Gk. phasis, aspect.] Resting stage between first and second mitotic divisions; interkinesis.

interplacental (in'tërplăsĕn'tăl) a. [L. inter, between; placenta, flat cake.] Between placentae.

interpleural (ĭn'tërploo'răl) a. [L. inter, between; Gk. pleuron, side.] Between pleurae.

interpositional growth,—of cells, by interposition between neighbouring cells without loss of contact; intrusive growth. Opp. sliding growth.

interpubic (in'tërpū'bik) a. [L. inter, between; pubes, mature.] Appl. the fibrocartilaginous lamina between pubic bones.

interracial (ĭn'tër-rā'sĭāl) a. [L. inter, between; radix, root.] Between races or breeds; appl. hybridisation, differences, etc.

interradial (in'tër-rā'diăl) a. [L. inter, between; radius, radius.]

Pert. an interradius.

- interradium (ĭn'tër-rā'diŭm) n. [I.. inter, between; radius, radius.] The area between two radii of any radially symmetrical animal.
- interradius (In'ter-ra'dius) n. [L. inter, between; radius, radius.]
 The radius of a radiate animal halfway between two perradii.
- interramal (ĭn'tër-rā'māl) a. [L. inter, between; ramus, branch.]
 Between branches or rami.
- interramicorn (ĭn'tër-rām'ĭkôrn) n.
 [L. inter, between; ramus, branch; cornu, horn.] A piece of a bird's bill beyond mandibular rami forming the gonys.

interrenal (ĭn'tër-rë'nāl) a. [L. inter, between; renes, kidneys.] Between the kidneys; appl. veins.

interrenal body,—a gland, situated between kidneys of elasmobranchs, representing the adrenal cortex of higher vertebrates.

interrupted (in'tër-rūp'tëd) a. [L. inter, between; rumpere, to break.]
With continuity broken; irregular; asymmetrical.

interruptedly pinnate, — pinnate with pairs of small leaflets occurring between larger ones.

- interscapular (In'tërskap'ūlar) a. [L. inter, between; scapula, shoulder-blade.] Between the shoulder-blades; appl. feathers; appl. brown fatty tissue, so-called hibernating gland, as in some rodents.
- intersegmental (ĭn'tërsĕgmĕn'tăl) a. [L. inter, between; segmentum, piece.] Between segments; between spinal segments, appl. axons, septa.
- intersegmentalia (In'tërsëg'mëntā'lĭā) n. plu. [L. inter, between; segmemtum, piece.] Sclerites between adjacent body segments in insects.
- interseptal (in'tërsëp'tăl) a. [L. inter, between; septum, fence.] Pert. spaces between septa or partitions.
- intersex (In'tërsëks) n. [L. inter, between; sexus, sex.] An organism with characteristics intermediate between typical male and typical female of its species; an organism first developing as a male or female,

then as an individual of the opposite sex; a sex mosaic in time; cf. gynandromorph.

interspecific (in terspesif'ik) a. [L. inter, between; species, kind.]
Between distinct species; appl. crosses, as mule, hinny, cattalo, tigron; appl. selection.

interspicular (ĭn'tërspĭk'ūlăr) a. [L. inter, between; spiculum, sharp point.] Occurring between spicules.

- interspinal (ĭn'tërspi'năl), interspinous (ĭn'tërspi'năs) a. [L. inter, between; spina, spine.] Occurring between spinous processes or between spines; appl. bones, muscles, ligaments.
- interstapedial (in'terstape'dial) a. [L. inter, between; stapes, stirrup.] Appl. a part of columella of ear.
- intersterility (in'tërstëril'iti) n. [L.
 inter, between; sterilis, unfruitful.]
 Incapacity for interbreeding.
- intersternal (in'terster'năl) a. [L. inter, between; sternum, breastbone.] Between the sterna; appl. ligaments connecting manubrium and body of sternum.
- intersternite (in'tërstër-nīt) n. [L. inter, between; sternum, breast-bone.] A sternal sclerite between thoracic segments of insects; intersegmental sternite; a furcasternite, q.v.
- interstitial (ĭn'tërstish'iăl) a. [L. inter, between; sistere, to set.] Occurring in interstices or spaces; appl. growth; appl. lamellae between Haversian systems; appl. cells within tissues, e.g. within gonad tissues.
- intertemporal (in'tërtëm'pörăl) n.
 [L. inter, between; tempora, temples.] A paired membrane bone, part of sphenoid complex, fusing with alisphenoids; dermosphenotic.
- intertentacular (In'tërtëntăk'ūlăr) a. [L. inter, between; tentaculum, feeler.] Between tentacles; appl. a ciliated tube opening at base of tentacles and connecting coelom and exterior, found in Molluscoidea.

- intertergal (interter'găl) a. | L. inter, between; tergum, back.] Between tergites or dorsal sclerites.
- intertidal (ĭn'tërtī'dăl) a. [L. inter, between; A.S. tid, time.] Appl. shore organisms living between high- and low-water marks.
- intertrabecula (In'tërtraběk'ūlă) n. [L. inter, between; trabecula, little beam.] A separate plate between the trabeculae anteriorly, in some birds.
- intertragic (intërtraj'ik) a. [L. inter, between; Gk. tragos, goat.] Appl. notch between tragus and antitragus.
- intertrochanteric (ĭn'tërtrökăntĕr'îk)
 a. [L. inter, between; Gk. trochanter, runner.] Between trochanters; appl. crest, line.
- intertrochlear (in'tërtrok'lëar) a. [L. inter, between; trochlea, pulley.] Appl. an ulnar ridge fitting into a groove of the humerus.
- intertubercular (in'tërtūbër'kūlār) a.
 [L. inter, between; tuberculum, small hump.] Appl. plane of body through tubercles of iliac crests; appl. sulcus between tubercles of humerus.
- intervarietal (în'tërvărī'ētăl) a. [L. inter, between; varius, diverse.] Appl. crosses between two distinct varieties of a species.
- interventricular (in'tërvëntrik'ūlăr)

 a. [L. inter, between; ventricula, small cavity.] Between ventricles; appl. foramen between third and lateral ventricles, foramen of Monro.
- intervertebral (ĭn'tërvĕr'tëbrăl) a.
 [L. inter, between; vertebra, vertebra.] Occurring between vertebrae; appl. discs, fibrocartilages, foramina, veins.
- intervillous (In'tërvil'us) a. [L. inter, between; villi, hairs.] Occurring between villi; appl. spaces in trophoblastic network filled with maternal blood.
- interxylary (in'tërzī'lārĭ) a. [L. inter, between; Gk. xylon, wood.]
 Between xylem strands; appl. phloem.
- interzonal (ĭn'tërzō'năl) a. [L. inter,

- between; zona, belt.] Between two zones; appl. spindle fibres uniting groups of daughter chromosomes in anaphase of mitosis.
- interzooecial (ĭn'tërzōē'sĭāl) a. [L. inter, between; zoon, animal; oikos, house.] Occurring among zooecia.
- intestinal (întěs'tīnăl, întëstī'năl) a.
 [L. intestinus, internal.] Pert.
 intestines; appl. glands, villi, etc.
- intestine (Intes'tin) n. [L. intestinus, internal.] Part of alimentary canal from pylorus to anus, or part corresponding to this.
- intextine (intěk'stǐn) n. [L. intus, within; exter, without.] An inner membrane of an extine.
- intima (ĭn'timā) n. [L. intimus, innermost.] The innermost lining membrane of a part or organ; tunica intima.
- intine (in'tin) n. [L. intus, within.]
 The inner covering membrane of a pollen grain, or of a spore. Opp. extine.
- intrabiontic (in'trăbion'tik) a. [L. intra, within; Gk. bios, life; on, being.] Appl. a process of selection occurring in a living unit.
- intrabulbar, —intragemmal, q.v. intracapsular (ĭn'trăkăp'sūlăr) a. [L. intra, within; capsula, small chest.]
- Contained within a capsule; appl. protoplasm of Radiolaria; appl. dendrites.
- intracardiac (ĭn'trăkâr'dĭăk) a. [L. intra, within; Gk. kardia, heart.] Endocardiac, q.v.
- intracartilaginous (ĭn'trăkâr'tĭlăj'ĭnŭs) a. [L. intra, within; cartilago, gristle.] Inside the cartilage; appl. ossification; endochondral.
- intracellular (In'trăsēl'ūlăr) a. [L. intra, within; cellula, little cell.] Within the cell.
- intracortical (in'trăkôr'tĭkăl) a. [L. intra, within; cortex, rind.] Within the cortex; uniting parts of brain cortex.
- intra-epithelial (ĭn'trăĕpĭthē'liăl) a. [L. intra, within; Gk. epi, upon; thele, nipple.] Occurring in epithelium; appl. glands, usually

- intrafascicular (ĭn'trăfăsĭk'ūlăr) a. [L. intra, within; fasciculus, little bundle.] Within a vascular bundle.
- intrafoliaceous (in'trăfoliă'shus) a.
 [L. intra, within; folium, leaf.]
 Appl. stipules encircling stem and forming a sheath; ochreate.
- intrafusal (ĭn'trăfū'zăl) a. [L. intra, within; fusus, spindle.] Appl. fasciculi and fibres connected respectively with neurotendinous and neuromuscular spindles.
- intragemmal (in'trăjĕm'āl) a. [L. intra, within; gemma, bud.] Within a taste-bud; appl. nerve fibres, spaces.
- intraglobular (in'trăglŏb'ūlăr) a. [L. intra, within; globulus, globule.]
 Occurring within a globule or corpuscle.
- intrajugular (in'trăjug'ūlăr) a. [L. intra, within; jugulum, throat.] Appl. a process in middle of jugular notch of occipital bone.
- intralamellar (ĭn'trălămel'ăr) a. [L. intra, within; lamella, thin plate.] Within a lamella; appl. trama of gill-bearing fungi.
- intralobular (ĭn'trălŏb'ūlăr) a. [L. intra, within; lobulus, small lobe.]
 Occurring within lobules; appl. veins draining liver lobules.
- intramembranous (in-trămēm'brănŭs) a. [L. intra, within; membrana, film.] Appl. bone developed in membrane.
- intramolecular (In'trămŏlěk'ūlăr) a. [L. intra, within; F. molécule, small particle.] Appl. plant respiration from splitting up of complex substances within the cell.
- intranuclear (în'trănū'kleār) a. [L. intra, within; nucleus, kernel.] Within the nucleus; appl. spindles, fibres, etc.
- intraparietal (In'trapari'etal) a. [L. intra, within; paries, wall.] Enclosed within an organ; within parietal lobe, as sulcus, etc.
- intrapetalous (ĭn'trăpět'ălŭs) a. [L. intra, within; Gk. petalon, leaf.] Situated in a petaloid area, in echinoderms.
- intrapetiolar (ĭn'trăpět'ĭölăr) a. [L.

intra, within; petiolus, little foot.] Within the petiole base expansion.

- intrapleural (in'trăploo'răl) a. [L. intra, within; Gk. pleuron, side.]
 Within the thoracic cavity.
- intrasegmental (in'trăsegmen'tăl) a. [L. intra, within; segmentum, part.] Appl. vertebra formed of cranial and caudal elements of same original myotome; cf. intermyotomic.
- intraselection (in'trăsēlěk'shun) n. [L. intra, within; selectio, choice.] Selection within an organ, of cells fittest to survive.
- intrasexual (in'trăsēk'sūāl) a. [L intra, within; sexus, sex.] Appl. selection of competing individuals of the same sex.
- intraspecific (in'trăspesif'ik) a. [L. intra, within; species, particular kind; facere, to make.] Within a species; appl. selection of individuals.
- intraspicular (ĭn'trăspik'ūlăr) a. [L. intra, within; spicula, small spike.] Having spicules completely embedded in spongin.
- intrastelar (in'trăstē'lăr) a. [L. intra, within; stela, pillar.] Within the stele of a stem or root; appl. ground tissue, bundles, etc.
- intratarsal (ĭn'trătâr'săl) a. [L. intra, within; tarsus, ankle.] Within the tarsus; appl. joint of reptilian limb between rows of tarsal bones.
- intrathyroid (ĭn'trăthī'roid) a. [L. intra, within; Gk. thyreos, shield; eidos, like.] Appl. a cartilage joining laminae of thyroid cartilage during infancy.
- intrauterine (in'trău'tërīn) a. [L. intra, within; uterus, womb.]
 Within the uterus.
- intravaginal (ĭn'trăvăj'ĭnăl) a. [L. intra, within; vagina, sheath.] Within vagina; contained within a sheath, as grass branches.
- intravascular (In'trăvăs'kūlăr) a. [L. intra, within; vasculum, small vessel.] Within blood-vessels.
- intraventricular (ĭn'trăvěntrīk'ūlăr)
 a. [L. intra, within; ventriculus,
 small cavity.] Within a ventricle;
 appl. caudate nucleus of corpus

striatum, seen within ventricle of

intravesical (ĭn'trăvěs'īkăl) a. [L. intra, within; vesica, bladder.] Within the bladder.

intravitelline (ĭn'trăvĭtěl'īn) a. [L. intra, within; vitellus, egg-yolk.] Within the yolk of an egg.

intraxylary (ĭn'trăzī'lărĭ) a. intra, within; Gk. xylon, wood.] Within the wood or xylem.

intrinsic (intrin'sik) a. [L. intrininwards.] Appl.muscles, as of tongue, of syrinx, appl. brightness sensation due to differential retinal response to different wave-lengths. extrinsic.

introitus (intro'itus) n. [L. introitus, entry.] An opening or orifice. intromittent (ĭn'trömĭt'ent) a. [L. intro, within; mittere, to send.] Adapted for inserting; appl. male copulatory organs.

introrse (ĭntrôrs') a. [L. introrsus, within.] Turned inwards or towards axis; of anthers, opening on side

next pistil.

introvert (ĭn'trövert) n. [L. intro, within; *vertere*, to turn.] which is capable of involution, as anterior region of body of certain zooids, of certain annulates, mouth of certain molluscs. extremity (ĭn'trövěrt') v. To turn, bend, or draw inwards.

intrusive growth,—see interposi-

tional growth.

intussusception (In'tŭssŭsĕp'shŭn) n. [L. intus, within; suscipere, to receive.] Growth in surface-extent or volume by intercalation of particles among those already present. Opp, accretion; cf. apposition.

inulase (ĭn'ūlās) n. [L. inula, elecam-A plant enzyme which hydrolyses inulin into laevulose.

inulin (ĭn'ūlĭn) n. [L. inula, elecampane.] A carbohydrate occurring in rhizomes and roots of many plants, and forming laevulose when hydrolysed; dahlia starch; $(C_6H_{10}O_5)_{\alpha}$.

invaginate (ĭnvăj'ĭnāt) υ. [L. in, into; vagina, sheath.] To involute or draw into a sheath; appl. insinking of wall of a cavity or vessel. a. Introverted; concave.

invagination (ĭnvăj'ĭnā'shŭn) n. [l.. in, into; vagina, sheath.] Involution; introversion; gastrulaformation by infolding of blastula ingestion by temporarily transformed periplast-like ectoplasm in certain Protozoa.

inversion (ĭnvër'shun) n. [L. invertere, to turn upside down.] Reversal in order of genes, or reversal of a chromosome segment, within the chromosome as a whole; a turning inward, inside out, or upside-down of a part.

invertase (invertas) n. [L. invertere, to turn into.] A plant enzyme which converts cane sugar into dextrose and laevulose; invertin; sucrase.

invertebrate (ĭnvĕr'tëbrāt) a. [L. in, not; vertebra, joint.] Back-boneless; without spinal column.

investing bones,—membrane bones. investment (invest'ment) n. [L. in, in; vestire, to clothe.] Outer covering of a part, organ, animal, plant.

involucel (ĭnvŏl'ūsĕl) n. [Dim. of L. involucrum, covering.] The small bracts at base of a secondary umbel; a partial involucre.

involucellate (invol'ūsěl'āt) a. | Dim. of L. involucrum, covering.] Bearing involucels.

involucellum,-involucel.

involucral (ĭn'vŏlū'krăl) a. [L. involucrum, covering.] Pert. or like an involucre.

involucrate (ĭn'vŏlū'krāt) a. [L. involucrum, covering.] Bearing involucres.

involucre (ĭn'vŏlū'kër) n. [L. involucrum, covering.] Bracts forming whorl at base of a condensed inflorescence, as of capitulum and umbel; a group of leaves surrounding antheridial and archegonial groups in bryophytes; involucrum. involucret,—involucel, q.v.

involucrum (ĭn'vŏlū'krŭm) n. [L. involucrum, covering.] In Hydromedusae, protective cup into which nematocysts can be spirally retracted; metanotum of Orthoptera; periosteal layer formed around dead portion of bone, in certain diseased conditions; an involucre, q.v.

involuntary (ĭnvŏl'ŭntărī) a. [L. in, not; voluntas, wish.] Not under control of will; appl. plain unstriped muscles, as of alimentary canal, and to their movements.

involute (In'vôlūt) a. [L. involutus, rolled up.] Of leaves, having the edges rolled inwards at each side;

of shells, closely coiled.

involution (in'volū'shūn) n. [L. involutus, rolled up.] Reduction to normal of enlarged, modified, or deformed conditions; shrinking of tissues in old age; appl. forms that have become deformed in structure, but not to such an extent as to be incapable of recovery; a rolling inwards, as of leaves; movement of cells to interior in a certain type of gastrulation; resting, appl. spores, stage, etc.

iodophilic (i'ōdöfil'ik) a. [Gk. ioeides, violet-like; philos, loving.] Staining darkly in iodine solution; appl. certain cytoplasmic inclusions and

vacuoles; iodinophilous.

iodopsin (i'odopsin) n. [Gk. ioeides, violet; opsis, sight.] Visual violet of retinal cones, a photo-sensitive protein-vitamin A compound.

iodothyrin (i'ōdöthi'rĭn) n. [Gk. ioeides, violet; thyreos, shield.] An iodine compound in the colloid

material of thyroid gland.

iodothyroglobulin (ī'ödöthī'röglöb'ūlīn) n. [Gk. ioeides, violet; thyreos, shield; L. globus, globe.] Compound of iodothyrin and nucleoprotein extractable hormone of the thyroid gland.

iridial angle,—filtration angle of eye: an angular recess between cornea and anterior surface of iris.

iridocytes (Ir'idösīts) n. plu. [Gk. iris, rainbow; kytos, hollow.] Guanin granules, bodies or plates, of which the reflecting tissue of skin of fishes is composed; iridescent cells in integument of Sepia.

iridomotor (Ĭr'idömō'tŏr) a. [L. iris, rainbow; movere, to move.] Connected with movements of iris.

iridophores,—iridocytes, q.v.

iris (I'ris) n. [L. iris, rainbow.] A thin, circular, contractile and vascular disc of eye between cornea and lens, and surrounding the pupil.

iris cells,—pigment cells surrounding cone and retinula of an omma-

tidium.

- irradiation (ĭr'rādĭā'shūn) n. [L. in, into; radius, ray.] Treatment with rays, as ultra-violet rays, X-rays, etc.; the spreading of an effect of a stimulus; spreading of an excitatory process; apparent enlargement of objects, due to difference in illumination.
- irritability (ir'ītābil'ītī) n. [L. irritare, to provoke.] Power of receiving external impressions, and reacting to them, inherent in living matter.

irritant (īr'ītănt) n. [L. irritare, to provoke.] An external stimulus which provokes a response.

irrorate (irr'ōrāt) a. [L. irrorare, to bedew.] Covered as if by minute droplets; dotted with minute colour markings, as wings of certain butterflies.

isadelphous (īsăděl'fūs) a. [Gk. isos, equal; adelphos, brother.]
With equal number of stamens in

two phalanges.

isauxesis (īsôksē'sis) n. [Gk. isos, equal; auxesis, growth.] Growth of a part at the same rate as that of the whole; ontogenetic heterauxesis.

ischiadic (is'kiăd'îk), ischial (is'kiăl)

a. [Gk. ischion, hip.] Pert. or
in region of hip; appl. artery, vein,
process of ischiopubic plate.

ischiocapsular (ĭs'kĭökāp'sūlăr) a. [Gk. ischion, hip; L. capsula, little chest.] Appl. a ligament joining

capsular ligament and hip.

ischiocavernosus (is'kiökăv'ērnō' sŭs) a. [Gk. ischion, hip; L. carus, hollow.] Appl. muscle between hip and corpora cavernosa; erector of penis, or of clitoris.

- ischioflexorius (is'kiöflekso'rius) n. [Gk. ischion, hip; L. flexus, bent.] Posterior thigh muscle in salamander, corresponding to semimembranosus.
- ischiopodite (ĭs'kĭŏpödīt) n. [Gk. ischion, hip; pous, foot.] Proximal joint of walking legs of certain Crustacea, or of maxillipedes.
- ischiopubic (is'kiöpū'bik) a. ischion, hip; L. pubes, adult.] Appl. a gap or fenestra between ischium and pubis; appl. a median cartilaginous plate with median and lateral processes, in Dipnoi.

ischiopubis (ĭs'kiöpū'bĭs) n. [Gk. ischion, hip; L. pubis, adult.] The ischium of pterodactyls, pubis being excluded from acetabulum; a fused ischium and pubis.

ischiorectal (ĭs'kĭörĕk'tăl) a.

[Gk. ischion, hip; L. rectus, straight.] Pert. ischium and rectum; appl. fossa and muscles.

ischium (is'kium) n. [Gk. ischion, hip.] The ventral and posterior bone of each half of pelvic girdle of vertebrates except fishes; an ischiopodite.

isidia (īsĭd'ĭă) n. plu. [Gk. isis, plant; idion, dim. | Coral-like soredia on surface of some lichens. Sing. isidium.

isidiferous (ī'sidif'erus) a. [Gk. isis, plant; L. ferre, to bear.] Bearing isidia; isidophorous.

isidioid (īsĭd'ĭoid) a. [Gk. isis, plant; idion, dim.; eidos, like.] Like an isidium.

isidium,-sing. of isidia.

island of Reil [J. C. Reil, German anatomist]. Insula, q.v.

islets of Langerhans [P. Langerhans, German anatomist]. Spherical or oval bodies scattered throughout the pancreas, concerned in metabolism of sugar in body; endocrine portion of pancreas.

isoagglutinin (ī'söagloot'inin) [Gk. isos, equal; L. agglutinare, to glue to.] Fertilizin or agglutinin of eggs which reacts on sperm of same species; cf. heteroagglutinin. isoagglutinogen (ī'söăglootin'öjěn),

isohaemagglutinogen (ī'söhēm'āglootĭn'ojēn) n. [Gk. isos, equal; haima, blood; genos, birth; L. agglutinare, to glue to.] Substance producing agglutination of erythrocytes within the same blood group.

isobilateral (ī'sobīlăt'ērāl) a. [Gk. isos, equal; L. bis, twice; latus, side.] Appl. a form of bilateral symmetry where a structure is divisible in two planes at right angles.

isobryonic (ī'söbriŏn'ik) a. isos, equal; bryein, to proliferate.] Developing equally, as lobes of dicotyledonous embryo; isobryous.

isocarpous (ī'sökâr'pŭs) a. isos, equal; karpos, fruit.] Having carpels and perianth divisions equal in number.

isocercal (ī'sösĕr'kăl) a. [Gk. isos, equal; kerkos, tail.] With vertebral column ending in median line of caudal fin.

isochela (ī'sökē'lă) n. [Gk. isos, equal; chele, claw.] A chela with two parts equally developed; a two-pronged or anchor-shaped spicule in certain sponges.

isochromosome (ī'sökrō'mösōm) n. [Gk. isos, equal; chroma, colour; soma, body.] Chromosome with identical arms united in a median centromere; metacentric derived from telocentric chromosome.

isochromous (ī'sökrō'mŭs) a. [Gk. equal; chroma, colour.] Equally tinted; uniformly coloured; isochromatic, isochroous.

isocortex (ī'sökôr'těks) n. [Gk. isos, equal; L. cortex, bark.] The part of cerebral cortex made up of six layers of nerve-cells. Opp. allocortex.

isocytic (î'sösĭt'ĭk) a. [Gk. isos, equal; kytos, hollow.] With all cells equal. isodactylous (ī'södăk'tĭlŭs) a. [Gk. isos, equal; daktylos, finger.] Having all digits of equal size.

isodemic (īsödem'ik) a. [Gk. isos, equal; demos, people.] With or pert. populations composed of an equal number of individuals; appl. lines on a map which pass through points representing equal population density.

isodiametric (ī'södīamet'rik) a. [Gk. isos, equal; dia, through; metron, measure.] Having equal diameters; appl. cells or other structures; appl. rounded or polyhedral cells.

isodont (ī'södŏnt) a. [Gk. isos, equal; odous, tooth.] Having teeth

all equal.

isodynamic (î'södÎnăm'îk) a. [Gk. isos, equal; dynamis, power.] Of equal strength; providing the same amount of energy; appl. foods.

isogamete (ī'sögămēt') n. [Gk. isos, equal; gametes, spouse.] One of a pair of undifferentiated gametes.

isogamous (īsŏg'ămŭs) a. [Gk. isos, equal; gamos, marriage.] Having

the gametes alike.

isogamy (īsŏg'ămĭ) n. [Gk. isos, equal; gamos, marriage.] Union of similar gametes, or of similar unicells.

isogenes (ī'söjēnz) n. plu. [Gk. isos, equal; genos, descent.] Lines on a map which connect points where same gene frequency is found.

isogenetic (ī'söjenet'ik) a. [Gk. isos, equal; genesis, descent.] Arising from the same or a similar origin; of the same genotype; isogenous.

isogenic (īsöjě'nĭk) a. [Gk. isos, equal; genos, race.] Homozygous. isogenomatic (ī'söjēnömăt'īk) [Gk. isos, equal; genos, race.] Containing similar sets of chromo-

somes; appl. nuclei; isogenomic. isognathous (īsŏg'nāthūs) a. [Gk. isos, equal; gnathos, jaw.] Having

both jaws alike.

isolate (i'sölāt) n. [It. isola, from; L. insula, island.] A breeding group restricted by isolation.

isolateral (ī'sölat'ērăl) a. [Gk. isos, equal; L. latus, side.] Having equal sides; appl. leaves with palisade tissue on both sides.

isolation (īsölā'shun) n. [L. insula, Separation from others; prevention of mating between breeding groups owing to spatial, topographical, ecological, phenological, physiological, genetic, or other barriers.

isolecithal (ī'sölēs'īthāl) a. [Gk. isos, equal; lekithos, yolk.] Appl. ova with yolk granules distributed nearly equally throughout substance.

isomar,—isophane, q.v.

isomastigote (ī'sömăs'tīgōt) a. [Gk. isos, equal; mastix, whip.] Having flagella of equal length.

isomere (ī'sömēr) n. [Gk. isos, equal; meros, part. A homologous struc-

ture or part.

isomerous (īsom'erus) a. [Gk. isos, equal; meros, part.] Having equal numbers of different parts; appl. flowers with same number of parts in each whorl; exhibiting isomery.

isometry (īsom'etri) n. [Gk. isos, equal; *metron*, measure.] Growth of a part at the same rate as the

standard or the whole.

isomorphic (ī'sömôr'fĭk) a. [Gk. isos, equal; morphe, shape.] Superficially alike; isomorphous; appl. alternation of diploid and haploid phases in morphologically similar generations.

isomorphism (ī'sömôr'fĭzm) n. [Gk. isos, equal; morphe, shape.] Apparent similarity of individuals of

different race or species.

[Gk. isos. isomyaric (ī'sömīăr'ik) a. equal; mys, muscle.] With adductor muscles equal in size; isomyarian.

isopedin (īsŏp'edin) n. [Gk. isopedos, level.] Inner layer of laminated bony material in cosmoid and ganoid fish scales.

isopetalous (ī'söpēt'ālūs) a. isos, equal; petalon, petal.] Having

similar petals.

isophane (ī'söfān) n. [Gk. isos, equal; phainein, to show.] A line connecting all places within a region at which a biological phenomenon, e.g. flowering of a plant, occurs at the same time; isomar, phenocontour.

isophene (ī'söfēn) n. [Gk. isos, equal; phainein, to show.] contour line delimiting area corresponding to a given frequency of a variant form; phenocontour.

isophenous (i'söfē'nŭs) a. [Gk. isos, equal; phainein, to show.]
Being of the same phenotype.

isophytoid (i'söfi'toid) a. [Gk. isos, equal; phyton, plant; eidos, form.] An 'individual' of a compound plant not differentiated from the rest.

isoplankt (i'soplangkt) n. [Gk. isos, equal; plangktos, wandering.] Line representing, on a map, distribution of equal amounts of plankton, or of particular plankton species.

isoploid (i'söploid) a. [Gk. isos, equal; aploos, onefold.] With an even number of chromosome sets in somatic cells. n. An isoploid individual.

isopodous (īsŏp'ödŭs) a. [Gk. isos, equal; pous, foot.] Having the legs alike and equal.

isopogonous (i'söpō'gōnūs) a. [Gk. isos, equal; pogon, beard.] Of feathers, having the two webs equal and similar.

isopolyploid (i'söpŏl'iploid) a., n. [Gk. 1sos, equal; polys, many; aploos, onefold; eidos, form.] Polyploid with an even number of chromosome sets, as tetraploid, hexaploid, octoploid, etc.

isopygous (i'sŏp'igous) a. [Gk. isos, equal; pyge, rump.] With pygidium and cephalon of equal size; appl. trilobites.

isospore (i'söspōr) n. [Gk. isos, equal; sporos, seed.] An agamete produced by schizogony; opp. anisospore.

isosporous (īsŏs'pörūs, īsöspō'rūs) a.
[Gk. isos, equal; sporos, seed.]
Having spores of one kind only;
homosporous.

isostemonous (īsöstēm'önūs) a. [Gk. isos, equal; L. stemon, warp.]
 Having stamens equal in number to that of sepals or of petals.

isotelic (i'sotel'ik) a. [Gk. isos, equal; telos, end.] Exhibiting, or tending to produce, the same effect; homoplastic; appl. food factors that can replace each other; pert. isotely.

isotels (i' htělz) n. plu. [Gk. isos,

equal; telos, end.] Substances having the same physiological, e.g. nutritional, effect.

isotely,—homoplasty, q.v.

isotomy (īsōt'ömī) n. [Gk. isos, equal; temnein, to cut.] Bifurcation repeated in a regular manner, as in crinoid brachia.

isotonic (īsötŏn'īk) a. [Gk. isos, equal; tonos, strain.] Of equal tension; having equal osmotic pressure.

isotonicity (i'sotonis'iti) n. [Gk. isos, equal; tonos, tone.] Normal tension under pressure or stimulus.

isotopic (i'sotop'ik) a. [Gk. isos, equal; topos, place.] Pert. isotopes, chemical elements having the same atomic number and identical chemical properties, but differing in atomic weight.

isotropic (l'sötröp'lk) a. [Gk. isos, equal; tropikos, turning.] Singly refracting in polarised light, appl. the light stripes of voluntary muscle fibres; opp. anisotropic; symmetrical around longitudinal axis; not influenced in any one direction more than another, appl. growth rate; without pre-determined axes, as eggs; isotropous.

isotropy (īsŏt'röpĭ) n. [Gk. isos, equal; trepein, to turn.] Absence of predetermined axes in eggs.

isotype (i'sotīp) n. [Gk. isos, equal; typos, pattern.] A specimen collected from the same plant as the holotype and at the same time.

isozoic (ī'sozō'īk) a. [Gk. isos, equal; zoon, animal.] Inhabited by similar forms of animal life.

isozooid (ī'sözō'oid) n. [Gk. isos, equal; zoon, animal; eidos, like.] A zooid similar to parent stock.

isthmiate (isth'miāt) a. [Gk. isthmos, neck.] Connected by an isthmuslike part.

isthmus (Isth'mus) n. [Gk. isthmos, neck.] A narrow structure connecting two larger parts, as those of aorta, acoustic meatus, limbic lobe, prostate, thyroid, etc., or between semi-cells; junction between perikaryon and axon-base.

iter (ĭt'ër) n. [L. iter, way.] A passage or canal, as those of middle ear, brain, etc.; an aqueduct.

ivory (ī'vorī) n. [L. ebur, ivory. through F. ivoire.] Dentine of teeth, usually that of elephant's tusks and similar structures, formed from odontoblasts.

Jacobson's cartilage [L. L. Jacobson, Danish anatomist]. Vomeronasal cartilage supporting Jacobson's organ.

Jacobson's nerve, — tympanic branch of the glossopharyngeal

nerve.

Jacobson's organ,—a diverticulum of olfactory organ in many vertebrates, often developing into an epithelium-lined sac opening into mouth; vomeronasal organ.

jactitation (jäktītā'shun) n. [L. jactare, to toss.] Process of scattering seeds by censer mechanism,

q.v.

jaculator (jăk'ūlātŏr) n. [L. jaculator, shooter.] A placental process, usually hooked, of certain fruits.

jaculatory (jäk'ūlătörī) a. [L. jaculatorius, throwing.] Darting out; capable of being emitted.

jaculatory duct,—portion of vas deferens which is capable of being protruded, in many animals.

jaculiferous (jăk'ūlif'ērŭs) a. [L. jaculum, a dart; ferre, to carry.] Bearing dart-like spines.

jarovization (yâr'ov\(^1z\)ā'sh\(^1u\)) n. [Russ. yarovizatsya, from yarovoi, vernal.] Vernalisation, q.v.

jaw (jô) n. [Akin to chaw, chew.] A structure, of vertebrates, supported by bone or cartilage, naked or sheathed in horn, or bearing teeth or horny plates, forming part of mouth, and helping to open or shut it; a similarly placed structure in invertebrates.

jaw foot,—maxillipede of Arthropoda. J-disc,—isotropic or I-disc, q.v.

jecorin (jek'orin) n. [L. jecur, liver.] A lecithin-like substance or phosphatide present in liver and other organs of the body.

jejunum (jëjoon'ŭm) n. [L. jejunus, empty.] Part of small intestine between duodenum and ileum.

jelly of Wharton [T. Wharton, English anatomist]. The gelatinous connective tissue surrounding the vessels of umbilical cord.

Johnston's organ [C. Johnston, British entomologist]. A statical or chordotonal organ in second segment of insect antenna.

joint (joint) n. [O.F. joindre, from L. jungere, to join.] Place of union or separation of two parts, as between bones; articulation; a node; portion between two nodes or joints.

Jordan's organ [K. Jordan, 200-

logist]. The chactosemata.

juba (joob'a) n. [L. juba, mane.]
A mane; a loose panicle.

jubate (joob'āt) a. [L. jubatus, maned.] With mane-like growth.

jugal (joog'āl) n. [L. jugum, yoke.]
The malar bone, between maxilla and squamosal. a. Pert. a jugum. jugate (joog'āt) a. [L. jugum, yoke.]
Having pairs of leaflets; furnished

with a jugum.

jugular (jūg'ūlār, joog'ūlār) a. [L. jugulum, collar-bone.] Pert. neck or throat; appl. veins, foramen, fossa, etc.; appl. nerve, the hyoidean or posterior post-trematic nerve; appl. ventral fish-fins beneath and in front of pectoral fins.

jugulum (joog'ūlūm) n. [L. jugulum, collar-bone.] The foreneck region of a bird's breast; in insects, the

jugum of wing.

jugum (joog'ũm) n. [L. jugum, yoke.] A pair of opposite leaflets; ridge on mericarp of umbelliferous plants; small lobe on posterior border of fore-wing of certain moths; ridge or depression connecting two structures; union of small sphenoidal wings in first year after birth. Jurassic (joorăs'īk) a. [Jura moun-Pert. or appl. Mesozoic period between Triassic and Cretaceous.

juvenal (joo'vënăl) a. [L. juvenalis, youthful.] Youthful; appl. plumage replacing nestling-down of first

plumage.

juvenile hormone, - secreted by corpus allatum and inhibiting development of adult characters, in certain insects.

juxta (jŭk'stă) n. [L. juxta, close to.] A ring-walled structure supporting

sheath of aedeagus.

juxtaglomerular (jŭk'stăglöměr'ūlăr) a. [L. juxta, close to; glomerare, to form into a ball.] Appl. cells surrounding arteriole of glomerulus of kidney.

juxtanuclear (jūk'stănū'kleăr) a. [L. juxta, close to; nucleus, kernel.] Appl. bodies: basophil deposits in cytoplasm of vitamin D-deficient

parathyroid cells.

K

kaino-,--see caeno-.

kako-,--see caco-.

kalidium (kălĭd'ĭŭm) n. [Dim. of Gk. kalia, hut.] A form of sporocarp, or cystocarp.

kalymma (kăl'imă) n. [Gk. kalymma, covering.] Vacuolated part of outer layer of certain radiolarians.

kalymmocytes (kăl'imösīts) n. plu. [Gk. kalymma, covering; kytos, cell.] In ascidians, certain folliclecells which migrate into the egg after maturation.

karvaster (kărĭăs'tër) n. [Gk. karyon, nut, nucleus; aster, star.] A starshaped group of chromosomes.

karyenchyma (karieng'kima) n. [Gk. karyon, nucleus; engchyma, infusion.] Nuclear sap; achromatin; karyochylema.

karyogamy (kăriŏg'ămi) n. karyon, nucleus; gamos, marriage.] Union and interchange of nuclear material.

karyokinesis (kăr'iökinē'sis) n. [Gk. karyon, nucleus; kinesis, movement. Indirect cell-division; mit-

karyology (kăr'iŏl'ögi) n. ſGk. karyon, nucleus; logos, discourse.] Nuclear cytology.

karyolymph (kar'iölimf') n. karyon, nucleus; L. lympha, water.] Nuclear sap; karyenchyma.

karyolysis (kār'iŏl'isis) n. [Gk. karyon, nucleus; lyein, to loosen.] Supposed dissolution of the nucleus in mitosis; liquefaction of nuclear membrane.

karyomere (kăr'iömēr) n. [Gk. karyon, nucleus; meros, part.] In mitosis, a small vesicle into which a chromosome is converted in one type of nuclear construction; karvomerite.

karyomicrosome (kăr'iömī'krösōm) [Gk. karyon, nucleus; mikros, small; soma, body.] A nuclear granule.

karyomite (kăr'iömīt') n. [Gk. karyon, nucleus; mitos, thread.] A chromosome.

karyomitome (kăr'ĭöm**i**tōm') n. [Gk. karyon, nucleus; mitoma, network.] The nuclear thread-work.

karyomitosis (kăr'iömitō'sis) n. [Gk. karyon, nucleus; mitos, thread.] Indirect nuclear division; mitosis.

karyon (kār'iŏn) n. [Gk. karyon, nucleus.] The cell-nucleus.

karyophans (kăr'iöfănz') n. plu. [Gk. karyon, nucleus; phainein, to appear. Microsomes or nucleus-like granules surrounded by an ovoid matrix, which form the spironeme and axoneme in stalk of infusoria.

karyophore (kăr'iöfōr) n. [Gk.karyon, nucleus; pherein, to bear.] System of ectoplasmic fibrils or membranes for mooring the nucleus, in certain

karyoplasm (kăr'iöplăzm) n. [Gk. karyon, nucleus; plasma, mould.] Nucleoplasm, the nuclear substance; cf. cytoplasm.

karyorhexis (kăr'iörěk'sis) n. [Gk. karyon, nucleus; rhexis, breaking.] Fragmentation of the nucleus.

248

- karyosome (kăr'iösōm) n. [Gk. karyon, nucleus; soma, body.] A nucleolus of the 'net-knot' type; a chromosome; a special aggregation of chromatin in resting nucleus; the cell-nucleus itself; cf. plasmosome.
- karyosphere (kăr'iösfēr) n. [Gk. karyon nucleus; sphaira, globe.]
 The large nucleolus from which arise all or most of the chromosomes of Protista.
- karyota (kărĭō'tă) n. plu. [Gk. karyon, nucleus.] Nucleated cells.
- karyotheca (kăr'iöthē'kă) n. [Gk. karyon, nucleus; theke, covering.] The nuclear membrane.
- **karyotin** (kăr'iötĭn) n. [Gk. karyon, nucleus.] Chromatin; nuclear substance.
- karyotype (kăr'iotīp) n. [Gk. karyon, nucleus; typos, pattern.] Group of individuals with the same chromosome number and similar linear arrangement of genes in homologous chromosomes; chromosome complement of such a group.

kata-,-also cata-.

- katabolism (kătăb'ölizm) n. [Gk. kata, down; bole, throw.] The destructive chemical processes in living organisms, opp. anabolism.
- katabolite (kătăb'ölît) n. [Gk. kata, down; bole, throw.] Any product of katabolism, e.g. urea.
- katagenesis (kăt'ăjën'ësis) n. [Gk. kata, down; genesis, descent.] Retrogressive evolution.
- katakinetic (kăt'ākinĕt'ĭk) a. [Gk. kata, down; kinein, to move.] Appl. process leading to discharge of energy; cf. anakinetic.
- katakinetomeres (kăt'ākinĕt'ömērz)
 n. plu. [Gk. kata, down; kinein, to move; meros, part.] Unreactive, stable, atoms or molecules.
- kataphase (kăt'ăfāz) n. [Gk. kata, down; phasis, appearance.] The stages of mitosis from formation of chromosomes to division of cell; cf. anaphase.
- kataphoresis (kăt'ăfŏrē'sis) n. [Gk. kata, down; pherein, to carry.]
 Transfer of fluids through a mem-

- brane from anode to kathode; electrical osmosis.
- kataphoric (kătăfôr'ik) a. [Gk. kata, down; pherein, to carry.] Appl. passive action, the result of lethargy.
- kataplexy (kăt'āplēksī) n. [Gk. kata, down; plessein, to strike.] Condition of an animal feigning death; maintenance of a postural reflex induced by restraint or shock; cataplexis; cf. catalepsis.

katastate (kăt'ăstăt, kătăs'tāt) n. [Gk. kata, down; stasis, state.] Any product of katabolic activity of protoplasm; katabolite.

katatrepsis (kăt'ătrēp'sis) n. [Gk. kata, down; trepein, to turn.] Stage of decreasing movement in blastokinesis.

katharobic (kăth'ărŏbĭk) a. [Gk. katharos, pure; bios, life.] Living in clean waters, appl. Protista; opp. saprobic.

kathodic (kăthŏd'ik) a. [Gk. kathodos, descent.] Not arising in conformity with genetic spiral; appl. leaves.

kation (kăt'iŏn, kăti'ŏn) n. [Gk. kata, down; ienai, to go.] A positively charged ion which moves towards kathode or negative pole; opp. anion.

Keber's organ [G. A. F. Keber, German zoologist]. Pericardial glands in lamellibranchs.

keel (kel) n. [A.S. ceol, ship.] The carina or breast-bone of flying birds; boat-shaped structure formed by two anterior petals of Leguminosae; ridge on blade or on other parts of grasses.

kenenchyma (kěněng'kimä) n. [Gk. kenos, empty; engchyma, infusion.] A tissue devoid of its living contents, as cork.

keph-,-see ceph-.

keraphyllous (kërăfil'ŭs) a. [Gk. keras, horn; phyllon, leaf.] Appl. layer of a hoof between horny and sensitive parts.

keratin (kĕr'átĭn) n. [Gk. keras, horn.] A scleroprotein forming the basis of epidermal structures such as horns, nails, hairs.

- keratinisation (ker'atīnīzā'shun) n. [Gk. keras, horn.] State of becoming horny; appl. cells of epiderm developing in a horny material.
- keratogenous (keratoj'enus) a. [Gk. keras, horn; -genes, producing.]
 Horn-producing.
- keratohyalin (ker'átöhī'álīn) n. [Gk. keras, horn; hyalos, glass.] Substance contained in stratum lucidum of skin.
- keratoid (kĕr'ătoid) a. [Gk. keras, horn; eidos, form.] Horny; resembling horn.
- keratose (kĕr'ātōs) a. [Gk. keras, horn.] Having horny fibres in skeleton, as certain sponges.
- kernel (kër'nël) n. [A.S. cyrnel, small grain.] The inner part of a seed containing the embryo.
- **ketogenic hormone,**—a prepituitary principle which influences fat metabolism.
- key-fruit,—winged achenes hanging in clusters, as of Acer and Fraxinus.
- key gene,—oligogene, q.v.
- kidney (kid'në) n. [A.S. cwith, womb; neere, kidney.] Nephros; paired organ which elaborates and excretes urine.
- kinaesthesis (kĭn'ēsthē'sis) n. [Gk. kinein, to move; aisthesis, perception.] Perception of movement due to stimulation of muscles, tendons, and joints; proprioception.
- kinaesthetic (kin'ésthět'ik) a. [Gk. kinein, to move; aisthesis, perception.] Pert. sense of movement or muscular effort; appl. sense, area.
- kinase (kinās') n. [Gk. kinein, to move.] A substance which transforms zymogens to enzymes.
- kinesiodic (kin'ēsiŏd'ik) a. [Gk. kinesis, movement; hodos, way.] Pert. motor nerve paths; cf. kinesodic.
- kinesis (kinē'sīs) n. [Gk. kinesis, movement.] Random movement; locomotor reactions depending on intensity of stimulus; variation in linear or angular velocity.
- kinesodic (kin'ēsŏd'ík) a. [Gk. kinesis, movement; hodos, way.]
 Conveying motor impulses.
- kinetic (kinět'ik) a. [Gk. kinein, to

- move.] Active; appl. function of movement, opp. static; energy employed in producing or changing motion; appl. division centre in cell-division.
- kinetoblast (kine'töbläst) n. [Gk. kinein, to move; blastos, bud.] Outer ciliated investment of aquatic larvae with special locomotor properties.
- kinetochore (kinē'tökōr) n. [Gk. kinein, to move; choros, place.] Spindle-attachment or -insertion region; centromere.
- kinetogenesis (kǐnē'töjēn'ēsīs) n.
 [Gk. kinein, to move; genesis,
 descent.] The evolution theory that
 animal structures have been produced by animal movements.
- kinetomeres (kinē'tomērz) n. plu. [Gk. kinein, to move; meros, part.] Molecules or atoms, reactive or stable, ana- and kata-kinetomeres.
- kinetonucleus (kǐnē'tönū'klēŭs) n. [Gk. kinein, to move; L. nucleus, kernel.] The secondary nucleus, kinetoplast, or parabasal body, in forms such as trypanosomes, in close connection with flagellum and undulating membrane; cf. trophonucleus.
- kinetoplasm (kine'töplazm) n. [Gk. kinein, to move; plasma, something formed.] An iron-containing nucleo-protein forming a source of energy to Nissl granules.
- kinetoplast (kine'toplast) n. [Gk. kinein, to move; plastos, formed.] Composite body formed by union of parabasal body with blepharoplast in some Mastigophora.
- kinetosome (kinē'tosōm) n. [Gk. kinein, to move; soma, body.] Oneofa groupof granules occupying the polar plate region in moss sporogenesis.
- kinoplasm (kin'oplăzm) n. [Gk. kinein, to move; plasma, mould.]
 The substance of attraction-sphere, astral rays, and spindle-fibres; archiplasm; ergastoplasm.
- kinoplasmosomes (kin'öpläz'mösōmz) n. plu. [Gk. kinein, to move; plasma, form; soma, body.] Phragmoplast fibres seen at periphery of cell plate.

klado-, -clado-, q.v.

klasma-plates (klaz'mă-plāts) n. plu.
[Gk. klasma, fragment; L. platus, flat.] Small parts of compound ambulacral plates separated by growth pressure, in echinoids.

kleisto-,-also cleisto-, q.v.

kleistogamous (klīstŏg'āmŭs) a. [Gk. kleistos, closed; gamos, marriage.] Fertilised in closed flowers.

kleronomous (klērŏn'ömŭs) a. [Gk. kleronomos, heir.] Inherited; appl.

paths in nervous system.

klinokinesis (kli'nökinë'sis) n. [Gk. klinein, to slope; kinesis, movement.] Change in rate of change of direction, or angular velocity, due to intensity of stimulation; cf. kinesis.

klon,—clone, q.v.

knee (nē) n. [A.S. cneow, knee.]
Genu; joint between femur and tibia; root-process of certain swamp-inhabiting trees.

knephoplankton (něf'öplängk'tön) n. [Gk. knephas, twilight; plangktos, wandering.] Plankton living at depths between thirty and five hundred metres; cf. phaoplankton, skotoplankton.

knot (nöt) n. [A.S. cnotta, knot.] In wood, base of branch surrounded by new layers of wood and hardened by pressure; in nuclear meshwork, small particles of chromatin where meshes cross.

Kölliker's canal [R. A. von Kölliker, Swiss zoologist]. A canal leading from otocyst towards exterior, as in

certain Cephalopoda.

kolyone (koli'on) n. [Gk. kolyein, to hinder.] Substance elaborated in, and conveyed from, a tissue or organ, which lessens or inhibits function of other tissues.

komma (kŏm'ă) n. [Gk. komma, clause.] Sarcomere; inocomma.

koniocortex (kön'iökôr'těks) n. [Gk. konis, dust; L. cortex, bark.] Granular part of cortex, characteristic of sensory areas of brain.

Krause's end-bulbs, — see end-bulbs.

Krause's membrane [W. J. F. Krause, German anatomist]. The

single or double row of dots in the light transverse band of striated muscle; telophragma, Z-disc.

Kupffer cells [K. W. von Kupffer, German anatomist]. Stellate macro-

phages of liver sinuses.

kyogenic (kiöjěn'ík) a. [Gk. kyesis, pregnancy; genos, descent.] Appl. prepituitary hormone stimulating secretion of progestin by corpora lutea.

kyto-,—cyto-, q.v.

L

labella (lăběl'ă) n. [L. labellum, small lip.] Paraglossa of insects; plu. of labellum.

labellate (lăběl'āt) a. [L. labellum, small lip.] Furnished with labella or small lips.

labelloid (lăběl'oid) a. [L. labellum, small lip; Gk. eidos, form.] Like a labellum.

labellum (lăběl'um) n. [L. labellum, small lip.] The lower petal, morphologically posterior, of an orchid; two fused lateral staminodes, as in flower of Zingiberaceae; small lobe beneath labrum, or labial palp, in insects; proboscis lobe.

labia (lā'bia) n. plu. [L. labium, lip.]

Lips; lip-like structures.

labia cerebri,—margins of cerebral hemispheres overlapping corpus callosum.

labia majora,—outer lips of vulva.

labia minora,—inner lips of vulva.
labial (lā'biāl) a. [L. labium, lip.]
Pert. or resembling a lip.

labial palp,—lobe-like structure near mouth of molluscs; jointed appendage on labium of insects.

labiate (lā'bīāt) a. [L. labium, lip.] Lip-like; possessing lips or thickened margins; having limb of calyx or corolla so divided that one portion overlaps the other.

labiatiflorous (la'biātiflō'rūs) a. [L. labium, lip; flos, flower.] Having the corolla divided into two lip-like

portions.

- labidophorous (lāb'idŏf'örŭs) a. [Gk. labis, forceps; pherein, to carry.] Possessing pincer-like organs.
- labiella (la biel'a) n. [L. labium, lip.] A mouth-part of Myriopoda.
- labile (lā'bīl, lāb'īl) a. [L. labilis, apt to slip.] Readily undergoing change; unstable; appl. genes that are constantly mutating.
- labiodental (lā'bĭödĕn'tăl) a. [L. labium, lip; dens, tooth.] Pert. lip and teeth; appl. an embryonic lamina; appl. labial surface of tooth.
- labiosternite (lā'biöstĕr'nīt) n. [L. labium, lip; sternum, breast-bone.]
 A median area between palpigers of insect head.
- labiostipes (lā'biöstī'pēz) n. [L. labium, lip; stipes, stalk.] A portion of basal part of insect labium.
- labipalp (lā'bĭpălp) n. [I.. labium, lip; palpare, to feel.] Labipalpus, labial palp of insects.
- labium (lā'bĭum) n. [L. labium, lip.] A lip, or lip-shaped structure; the fused second maxillae of insects. Plu. labia.
- labral (lā'brăl) a. [L. labrum, lip.] Pert. a labrum.
- labrocyte (lăb'rösīt) n. [Gk. labros, greedy; kytos, hollow.] A mast cell.
- labrum (lă'brum) n. [L. labrum, lip.] Anterior lip of certain arthropods; hypostoma of trilobites; outer margin of mouth of gastropod shell; ring of fibrocartilage, ambon, q.v.
- labyrinth (lăb'īrīnth) n. [L. labyrinthus, labyrinth.] The complex internal ear, bony or membranous; lateral mass of air-cells of ethmoidal bone; portions of kidney cortex with uriniferous tubules; tracheal tympanum.
- labyrinthodont (lab'irin'thödönt) a.
 [Gk. labyrinthos, labyrinth; odous, tooth.] Having teeth with great complexity of dentine arrangement.
- lac (lăk) n. [Persian, lak, lacquer.] A resinous secretion of lac glands of certain Coccidae, composition depending on the food plant.

- laccate (lăk'āt) a. [It. lacca, varnish.] Appearing as if varnished.
- lacerated (lăs'ērātēd) a. [L. lacerare, to tear.] Having margin or apex deeply cut into irregular lobes.
- lacertiform (lăsĕr'tifôrm) a. [L. lacerta, lizard; forma, shape.]
 Having the shape of a lizard.
- lacertus (lăsĕr'tūs) n. |L. lacertus, arm-muscle.] Lacertus fibrosus, aponeurosis of tendon of biceps muscle of the arm; bicipital fascia.
- lachrymal,—lacrimal, q.v.
- lacinia (lăsīn'iā) n. [L. lacinia, flap.]
 Segment of an incised leaf; slender projection from margin of a thallus; extension of posterior margin of proglottis over anterior part of following proglottis; inner division of endopodite or stipes of maxilla of insects; fimbria, q.v.
- laciniate (lăsin'iāt) a. [L. lacinia, flap.] Irregularly incised, as petals; fringed; appl. a ligament of the ankle, the internal annular ligament.
- laciniform (lăsĭn'ĭfôrm) a. [L. lacinia, flap; forma, shape.] Shaped like lacinia; fringe-like.
- lacinula (lăsin'ūlă) n. [L. lacinia, dim., flap.] Small lacinia; inflexed sharp point of petal.
- lacinulate (lăsin'ūlāt) a. [L. lacinia, dim., flap.] Having lacinulae.
- lacrimal (lāk'rīmāl) a. [L. lacrima, tear.] Secreting or pert. tears; pert. or situated near lacrimal organ; appl. artery, bone, duct, glands, nerve, papillae, sac; also lachrymal.
- lacrimonasal (lăk'rĭmönā'zăl) a. [L. lacrima, tear; nasus, nose.] Pert. lacrimal and nasal bones or duct.
- lacrimose (lăk'rimōs) a. [L. lacrima, tear.] Bearing tear-shaped appendages, as gills of certain fungi.
- lactalbumin (lăk'tălbū'mĭn) n. [L. lac, milk; albumen, egg-white.] An albumin found in milk.
- lactase (lăk'tās) n. [L. lac, milk.] An intestinal enzyme converting lactose into glucose and galactose; also found in certain plants.

lactation (lăktā'shūn) n. [L. lac, milk.] Secretion of milk in mammary glands; period during which milk is secreted.

lacteals (lăk'tëălz) n. plu. [L. lac, milk.] Chyliferous or lymphatic vessels of small intestine; ducts which carry latex.

lactescent (lăktěs'ënt) a. [L. lactescere, to turn to milk.] Producing milk; yielding latex.

lactic (lăk'tik) a. [L. lac, milk.]

Pert. milk; appl. bacilli, acid.

lactiferous (lāktĭf'ērūs) a. [L. lac, milk; ferre, to carry.] Forming or carrying milk; carrying latex.

lactochrome (lăk'tökrōm) n. [L. lac, milk; Gk. chroma, colour.] A nitrogenous colouring matter in milk; lactoflavin, q.v.

lactoflavin (lăk'töflā'vĭn) n. [L. lac,
milk; flavus, yellow.] Vitamin
B₂; riboflavin, q.v.; C₁₇H₂₀O₆N₄.

lactogenic (lāk'töjēn'ik) a. [L. lac, milk; Gk.-genes, producing.] Pert., or stimulating, secretion of milk; appl. a prepituitary hormone inducing secretion of milk in mammals and of crop milk in the pigeon; appl. interval between parturition and ovulation, or between parturition and menstruation.

lactoglobulin (lăk'töglŏb'ūlĭn) n. [L. lac, milk; globulus, dim. of globus, globe.] The specific protein of milk, insoluble in water.

lactoproteid (lăk'töpro'tëid) n. [L. lac, milk; Gk. protos, first; eidos, form.] Any milk proteid.

lactose (lăk'tōs) n. [L. lac, milk.] Milk-sugar, C₁₂H₂₂O₁₁.

lacuna (lăkū'nā) n. [L. lacuna, cavity.] A space between cells; sinus; urethral follicle; cavity in bone; small cavity or depression on surface in lichens; a leaf gap. Plu. lacunae.

lacunar (lăkū'năr) a. [L. lacuna, cavity.] Having, resembling, or pert. lacunae.

lacunate (lăkū'nāt) a. [L. lacuna, a hollow.] Lacunar; pert. lacunae; appl. collenchyma, with cell-walls thickened where bordering intercellular spaces.

lacunose (laku'nos) a. [L. lacuna, cavity.] Having many cavities; pitted.

lacunosorugose (lăkū'nōsöroo'gōs)
a. [L. lacuna, cavity; rugosus, wrinkled.] Having deep furrows or pits, as some seeds and fruits.

lacus lacrimalis,—the triangular space between eyelids which contains lacrimal caruncle and receives tears from orifices of the lacrimal ducts.

lacustrine (lăkŭs'trĭn) a. [L. lacus, lake.] Pert. or living in or beside lakes.

laeotropic (lē'ötröp'ik), laeotropous (lēŏt'röpŭs) a. [Gk. laios, left; trope, turning.] Inclined, turned, or coiled to the left.

laevulose (lē'vūlōs) n. [L. laevus, left.] Fruit - sugar; fructose, C₆H₁₂O₆.

lagena (lage'nă, lăje'nă) n. [L. lagena, flask.] Terminal portion of the cochlear duct or scala media.

lageniform (lăgē'nĭfôrm, lăjē'nĭfôrm) a. [L. lagena, flask; forma, shape.] Shaped like a flask.

lagopodous (lăgŏp'ödŭs) a. [Gk. lagos, hare; pous, foot.] Possessing hairy or feathery feet.

Lamarckian (lămârk'ĭăn) a. [J.-B. de Lamarck, French biologist]. Of or pert. theories put forward by Lamarck.

Lamarckism (lămârk'izm) n. The evolution theory of Lamarck, embodying the principle that acquired characteristics are transmissible.

lambda (lăm'dă) n. [Gk. Λ, lambda.] The junction of lambdoid and sagittal sutures.

lambdoid (lăm'doid] a. [Gk. A, lambda; eidos, form.] A-shaped; appl. the cranial suture joining occipital and parietal bones.

lamella (lămel'ă) n. [L. lamella, small plate.] Any thin plate- or scale-like structure; the gill of an agaric. lamellar (lăměl'ăr), lamellate (lăm'ělāt) a. [L. lamella, small plate.] Composed of, or possessing

thin plates.

lamellibranchiate (lăměl'ibrăng'kiāt)
a. [L. lamella, small plate;
branchiae, gills.] Having platelike gills on each side; with bilaterally compressed symmetrical
body, like a bivalve.

lamellicorn (lăměl'īkôrn) a. [L. lamella, small plate; cornu, horn.]
Having antennal joints expanded

into flattened plates.

lamelliferous (lăm'ēlĭf'érŭs) a. [L. lamella, small plate; ferre, to carry.] Having small plates or scales.

lamelliform (lăměl'ifôrm) a. [L. lamella, small plate; forma, shape.] Plate-like.

lamellirostral (lăměl'írŏs'trăl) a. [L. lamella, small plate; rostrum, beak.] Having inner edges of bill bearing lamella-like ridges.

lamina (lăm'înă) n. [L. lamina, plate.] A thin layer, or scale; blade of leaf; one of thin plate-like expansions of sensitive tissue which fit into grooves on inside of horse-hoof.

lamina choriocapillaris,—capillary plexus constituting inner layer of

choroid.

lamina cribrosa,—cribriform plate, g.v.; membraneous portion of sclera at site of attachment of optic nerve and with perforations for axons of ganglion cells of retina.

lamina perpendicularis,—median process of mesethmoid or ethmoid forming proximal or bony part of

nasal septum.

lamina terminalis,—thin layer of grey matter forming anterior boundary of third ventricle of brain.

- lamina vasculosa,—outer layer of choroid beneath suprachoroid membrane
- laminar (lăm'ĭnăr), laminiform (lăm'ĭnĭfôrm) a. [L. lamina, plate; forma, shape.] Consisting of plates or thin layers; laminous.
- laminarian (lăminā'rian) a. [Lamin-

aria, a genus of brown seaweeds.]
Appl. zone between low water to about fifteen fathoms.

lamination (lăm'ĭnā'shūn) n. [L. lamina, plate.] The formation of thin plates or layers; arrangement in layers, as nerve cells of cerebral cortex.

laminiplantar (lăm'inīplăn'tăr) a. [L. lamina, plate; planta, sole of foot.] Having scales of metatarsus meeting

behind in a smooth ridge.

lanate (lā'nāt) a. [L. lana, wool.] Woolly; covered with short hairlike processes giving woolly appearance to surface.

lance-linear (lăns'-lĭn'eăr) a. [L. lancea, lance; linea, line.] Between lanceolate and linear in form.

lance-oblong (lăns'-ŏb'lŏng) a. [L. lancea, lance; oblongus, oblong.]
Oblong with tapering ends.

lanceolate (lăn'sëölāt) a. [L. lanceola, little lance.] Slightly broad, or tapering, at base and tapering to point; lance-shaped.

lance-oval (lăns'-ō'văl), lance-ovate (lăns'-ō'vāt) a. [L. lancea, lance; ovalis, oval.] Having a shape intermediate between lanceolate and oval

lancet (lăn'sët) n. [F. lancette, from L. lancea, lance.] Stylet of sting, as in bees.

lancet - plates, — plates supporting water-vascular vessels of Blastoidea.
 Langerhans,—see follicles, islets.

Langhans' cells [T. Langhans, German histologist]. Giant cells of inner layer of trophoblast or layer of Langhans.

languet, languette (lănggět') n.
[F. languette, small tongue.] A
process on branchial sac of ascidians.
laniary (lān'išri) a. [I. laniare to

laniary (lān'ĭārĭ) a. [L. laniare, to tear to pieces.] Term appl. to canine tooth.

lantern,-see Aristotle's lantern.

lanthanin (lănthā'nĭn) n. [Gk. lanthanein, to conceal.] Oxychromatin; linin, q.v.

lanuginous (lănū'jinŭs) a. [L.
lanugo, down.] Covered with down;
cf. lanate.

- lanugo (lănū'gō) n. [L. lanugo, down.] The downy covering on a foetus, begins to be shed before birth. lapidicolous (lăp'idik'ölŭs) a. [L.
- lapis, stone; colere, to cultivate.]
 Appl. animals that live under stones.
 lapillus (lapil'us) n. [L. lapillus,
- lapillus (lăpil'ŭs) n. [L. lapillus, pebble.] A small otolith in utriculus of teleosts.
- lappaceous (lăpā'shŭs) a. [L. lappa, bur.] Prickly.
- lappet (lăp'ēt) n. [M.E. lappe, loose hanging part.] One of paired lobes extending downwards from distal end of stomodaeum in jelly-fish; lobe of a sea-anemone gullet; wattle of a bird.
- larmier (lâr'myër) n. [F. larme, tear.] Tear pit; saccus lacrimalis.
- larva (larva) n. [L. larva, ghost.]
 An embryo which becomes selfsustaining and independent before
 it has assumed the characteristic
 features of its parents.
- larval (lâr'văl) a. [L. larva, ghost.] Pert. a larva; in the larval stage.
- larviform (lâr'vifôrm) a. [L. larva, ghost; forma, shape.] Shaped like a larva.
- larviparous (lârvip'ărŭs) a. [L. larva, ghost; parere, to produce.] Producing live larvae.
- larvivorous (lârviv'örŭs) a. [L. larva, ghost; vorare, to devour.] Larvaeating.
- larvule (lâr'vūl) n. [L. larvula, small larva.] Young larva.
- laryngeal (lărĭn'jeăl) a. [Gk. laryngx, upper part of windpipe.] Pert. or near the larynx; appl. artery, vein, nerve, etc.
- laryngopharynx (lăring'göfăr'ingks)
 n. [Gk. laryngx, larynx; pharyngx,
 gullet.] Part of pharynx between
 soft palate and oesophagus.
- laryngotracheal (lărĭng'götrā'këăl)
 a. [Gk. laryngx, windpipe; L.
 trachea, trachea.] Pert. larynx and
 trachea; appl. embryonic groove
 and tube.
- larynx (lar'ingks) n. [Gk. laryngx, larynx.] The organ of voice in most vertebrates, except birds.
- asso-cells,-filamented hemispheri-

- cal adhesive cells, investing tentacles of Ctenophora.
- lata-type,—a mutant with one or more supernumerary chromosomes as compared with its parent (from Oenothera lata).
- latebra (lăt'ëbră) n. [L. latebra, hiding-place.] The bulb or flaskshaped mass of white yolk in eggs.
- latebricole (lăt'ĕbrĭkōl) a. [L. latebra, hiding-place; colere, to inhabit.] Inhabiting holes.
- latent (la tent) a. [L. latens, hidden.] Lying dormant but capable of development under favourable circumstances; appl. buds, resting stages, characters.
- latent bodies,—the resting stage of certain Haemoflagellata.
- latent period,—the time interval between completion of presentation of a stimulus and the beginning of a reaction; reaction time.
- lateral (lăt'erāl) a. [L. latus, side.]
 Pert. or situated at a side; or at a
 side of an axis.
- lateral chain theory,—see side-chain.
 lateral line,—longitudinal line at each side of body of fishes marking position of sensory cells.
- lateral mesenteries, the mesenteries of Zoantharia, excluding directive or dorsal and ventral pairs.
- lateralia (lăterā'lĭă) n. plu. [L. latus, side.] The lateral plates of Cirripedia.
- laterigrade (lat'erigrad) a. [L. latus, side; gradus, step.] Walking sideways, as a crab.
- laterinerved (lăt'ërinervd') a. [L. latus, side; nervus, sinew.] With lateral veins.
- laterite (lăt'ērīt) n. [L. later, brick.]

 Appl. tropical red soils containing alumina and iron oxides and little silica owing to leaching under hot, moist conditions.
- laterocranium (lăt'ërökră'niŭm) n. [L. latus, side; cranium, skull.] Area of insect head comprising genae and postgenae.
- latero-sensory (lat'erösen'söri) a. [L. latus, side; sensus, sense.] Appl. system of lateral sense-organs in fishes, or lateral line system.

- laterosternites(lăt'eröster'nīts)n. plu.
 [L. latus, side; sternum, breastbone.] Sclerites at side of eusternum, as in Dermaptera and Isoptera.
- laterotergites (lăt'eröter'jīts) n. plu.
 [L. latus, side; tergum, back.]
 Small sclerites adjoining tergum
 of abdominal segments in some
 crustaceans and insects.
- latex (lā'tēks) n. [L. latex, a liquid.]
 A milky, or clear, sometimes coloured, juice or emulsion of diverse composition found in some plants, as in spurges, rubber trees, certain agarics, etc.
- laticifer (latis'ifer) n. [L. latex, a liquid; ferre, to carry.] Any latex-containing cell, series of cells, or duct
- laticiferous (lăt'īsīf'ērŭs) a. [L. latex, a liquid; ferre, to carry.] Conveying latex; appl. cells, tissue, vessels.
- latiplantar (lăt'îplăn'tăr) a. [L. latus, side; planta, sole of foot.] Having hinder tarsal surface rounded.
- latirostral (lăt'ĭrŏs'trăl) a. [L. latus, broad; rostrum, beak.] Broadbeaked.
- latiseptate (lăt'isĕp'tāt) a [L latus, broad; septum, septum.] Having a broad septum in the silicula.
- latitudinal furrow,—one running round a segmenting egg above and parallel to the equatorial.
- Laurer-Stieda canal,—a canal leading from junction of oviduct and vitelline duct to opening on dorsal surface in trematodes.
- laurinoxylon (lôr'inözī'lŏn) n. [L.
 laurus, laurel; Gk. xylon, wood.]
 Any fossil wood.
- law of acceleration,—the generalisation that organs of greater importance develop more quickly.
- lax (lăks) a. [L. laxus, loose.] Loose, as appl. panicle.
- layer of Langhans [T. Langhans, German histologist]. Cytotrophoblast.
- leader (lē'dër) n. [A.S. laedan, to lead.] Highest shoot or part of trunk of a tree.
- leaf (lef) n. [A.S. leaf, leaf.] An

- expanded outgrowth of a stem, usually green.
- leaf-buttress,—lateral prominence on shoot axis, due to underlying leaf primordium, representing leaf-base.
- leaf-cushions,—prominent persistent leaf-bases, furnishing diagnostic characters in certain extinct plants.
- leaf-gap,—mesh of stelar network, corresponding to site of leaf attachment in ferns; gap in vascular cylinder of stem, a parenchymatous region associated with leaf-traces; lacuna.
- leaflet,—a small leaf; individual unit of a compound leaf.

leaf-stalk,-petiole.

- leaf-trace,—vascular bundles extending from stem bundles to leafbase. *Cf.* girdle bundles.
- leberidocytes (lěbērī'dösīts) n. plu.
 [Gk. leberis, exuvia; kytos, hollow.]
 Cells containing glycogen, and
 developing from and regressing to
 leucocytes, found in blood of Arachnida at moulting.
- lechriodont (lěk'riödŏnt') a. [Gk. lechrios, crosswise; odous, tooth.] With vomerine and pterygoid teeth in a row nearly transverse.
- lecithalbumin (lěs'īthălbū'mĭn) n.
 [Gk. lekithos, egg-yolk; L.
 albumen, white of egg.] A substance, consisting of albumin and
 lecithin, of various body organs.
- lecithelles (lĕs'īthĕlz) n. plu. [Gk. lekithos, egg-yolk.] Yolk granules in hypoblastic or other lecithoblasts. lecithin (lĕs'īthīn) n. [Gk. lekithos, egg-yolk.] A phosphorised fat or phospholipide of cell-protoplasm.
- lecithoblast (lĕs'īthöblāst) n. [Gk. lekithos, egg-yolk; blastos, bud.] In developing eggs, the yolk-containing blastomeres.
- lecithocoel (lēs'īthösēl) n. [Gk. lekithos, egg-yolk; koilos, hollow.] Segmentation cavity of holoblastic eggs.
- lectoallotype (lek'töäl'ötīp) n. [Gk. lektos, chosen; allos, other; typos, pattern.] A specimen of the opposite sex to that of the lectotype and subsequently chosen from the original material.

lectotype (lěk'tötip) n. [Gk. lektos, chosen; typos, pattern.] A specimen chosen from syntypes to designate type of species.

legume (lĕg'ūm) n. [L. legumen, pulse.] Dehiscent one-celled, twovalved carpel, as pod of pea or

bean.

legumin (lěgū'min) n. [L. legumen, pulse.] A globulin found in seeds of Leguminosae.

leiotrichous (liŏt'rīkŭs) a. [Gk. leios, smooth; thrix, hair.] Having straight hair; leiothric.

leiotropic,—laeotropic, q.v.

lemma (lĕm'ā) n. [Gk. lemma, husk.] A valve or flowering glume; lower or outer palea, bract with axillary flower.

lemniscus (lěmnis'kŭs) n. [Gk. lemniskos, fillet.] One of paired club-shaped organs at base of acanthocephalan proboscis; a fillet of fibres on each side of cerebral peduncles.

lenitic (lēnīt'īk) a. [L. lenis, smooth.]

Lentic, q.v.

lens (lenz) n. [L. lens, lentil.] A transparent part of eye, which focuses rays of light on retina; crystalline lens; modified portion of cornea in front of each element of a compound eye.

lentic (lĕn'tĭk) a. [L. lentus, slow.]
Appl. or pert. standing water;
living in swamp, pond, or lake;

lenitic. Opp. lotic.

lenticel (len'tisel) n. [L. lens, lentil.]

Ventilating pore in angiosperm stems or roots; canal in cork;

a lenticular gland.

- lenticular (lentik'ūlăr) a. [L. lenticula, dim. of lens, lentil.] Shaped like a double convex lens; lentiform; appl. glands, lymphoid structures between pyloric glands. n. Tip of incus articulating with stapes, often ossified as a separate unit.
- lenticulate (lěntik'ūlāt) a. [L. lens, lentil.] Meeting in a sharp point; depressed, circular, and frequently ribbed.
- lentiform (lěn'tifôrm) a. [L. lens,

lentil; forma, shape.] Lentilshaped; lenticular; appl. nucleus, the extraventricular portion of corpus striatum.

lentigerous (lentij'erus) a. [L. lens, lentil; gerere, to bear.] Furnished

with a lens.

lentiginose (lěntij'īnōs), lentiginous (lěntij'īnŭs) a. [L. lentigo, freckle.] Freckled; speckled; bearing numerous small dots.

lento-capillary point,—point, just above wilting coefficient, at which flow of water towards root hairs is impeded on account of surface tension resistance.

leotropic,—laeotropic, q.v.

lepidodendroid (lĕp'idödĕn'droid) a. [Gk. lepis, scale; dendron, tree; eidos, form.] Pert. Lepidodendron; having scale-like leaf-scars.

lepidoid (lĕp'ĭdoid) a. [Gk. lepis, scale; eidos, form.] Resembling a scale or scales.

lepidomorial,—pert., or composed of, lepidomoria.

lepidomorium (lěp'ĭdömŏr'ĭŭm) n. [Gk. lepis, scale; morion, constituent part.] Small scale, or unit of composite scale, with bony base and conical or conoid crown of dentine, containing pulp cavity and sometimes covered with enamel. Plu. lepidomoria.

lepidophyte (lep'idöfit) n. [Gk. lepis, scale; phyton, plant.] A

fossil plant of fern family.

lepidopterous (lĕp'ĭdŏp'térŭs) a. [Gk. lepis, scale; pteron, wing.] Having wings covered with minute overlapping scales; pert. moths, butterflies.

lepidosteoid (lĕp'idŏs'tëoid) a. [Gk. lepis, scale; osteon, bone; eidos, form.] Appl. a ganoid scale lacking cosmine.

lepidote (lěp'ídōt) a. [Gk. lepidotos, scaly.] Covered with minute scales. lepidotic (lěp'ídō'tik) a. [Gk. lepi-

dotos, scaly.] Appl. an acid found in wings of some Lepidoptera.

lepidotrichia (lěp'Idötrik'iă) n. plu. [Gk. lepis, scale; thrix, hair.] The bony actinotrichia of teleosts.

lepocyte (lĕp'ösit) n. [Gk. lepis, husk; kytos, hollow.] A cell with a defining cell-wall, opp, gymnocyte.

lepospondylous (lēp'öspŏn'dīlūs) a. [Gk. lepis, husk; sphondylos, vertebra.] Having amphicoelous, or hour-glass shaped, vertebrae.

leptocentric (lep'tosen'trik) a. [Gk. leptos, slender; kentron, centre.] Appl. concentric bundle with central leptome.

leptocephaloid (lep'tökef'äloid, -sef-) a. [Gk. leptos, slender; kephale, head; eidos, form.] Resembling or having the shape of eel larvae.

leptocercal (lep'töser'käl) a. [Gk. leptos, slender; kerkos, tail.] With long slender tapering tail, as some fishes; leptocercous, appl. protozoa.

leptodactylous (lěp'tödāk'tĭlūs) a. [Gk. leptos, slender; daktylos, finger.] Having slender digits.

leptodermatous (lěp'töděr'mātŭs) a. [Gk. leptos, slender; derma, skin.] Thin-skinned; appl. various thecae; leptodermic, leptodermous.

leptoid (lěp'toid) n. [Gk. leptos, slender; etdos, form.] One of the thin-walled cortical cells forming strand projecting into the central cylinder of rhizome in mosses; a tubular cell in stem of certain pteridophytes.

leptome (lep'tōm) n. [Gk. leptos, slender.] Phloem-like part of vascular tissue of plant stems; leptomestome; bast; also leptom.

leptomeninges (lep'tömenin'jez) n. plu. [Gk. leptos, slender; meningx, membrane.] The pia mater and arachnoid membrane.

leptonema (lĕp'tönē'mā) n. [Gk. leptos, slender; nema, thread.]
Fine unpaired chromosome thread at leptotene.

leptophloem (lep'toflo'em) n. [Gk. leptos, slender; phloios, smooth bark.] Rudimentary bast tissue.

leptophyllous (lĕp'töfil'ūs) a. [Gk. leptos, slender; phyllon, lcaf.]
With slender leaves; having a small leaf area, under 25 square millimetres.

leptosome (lěp'tösom) a. [Gk. leptos,

slender; soma, body.] Tall and slender; opp. eurysome.

leptosporangiate (lěp'töspöran'jlāt) a. [Gk. leptos, slender; sporos, seed; anggeion, vessel.] With sporogenous tissue developing from outer cell of periclinal division. Opp. eusporangiate.

leptostroterate (lĕp'tostrō'terāt) a. [Gk. leptos, slender; strotos, covered.] With ambulacral plates narrow and crowded together, as in certain Stelleroidea.

leptotene (lěp'totēn) n. [Gk. leptos, slender; tainia, band.] Early stage of the prophase of meiosis where chromatin is in form of fine threads.

leptotrombicula (lĕp'totrombĭk'ūlā)

n. [Gk. leptos, slender; It. tromba,
trumpet.] The larval form of a
trombicula.

leptoxylem (lěp'tózī'lëm) n. [Gk. leptos, slender; xylon, wood.] Rudimentary wood tissue.

lepto-zygotene (lěp'tözī'gotēn) a. [Gk. leptos, slender; zygon, yoke; tainia, band.] Appl. transition stage between leptonema and zygonema.

leptus (lĕp'tŭs) n. [Gk. leptos, small.]
The six-legged larva of mites.

lethal (lē'thāl) a. [L. letum, death.] Causing death; of a parasite, fatal or deadly in relation to a particular host; appl. a hereditary factor which so influences development that the individual is rendered non-viable. n. A lethal factor.

lethality (lēthāl'īti) n. [I.. letum, death.] The ratio of fatal cases to total number of cases affected by a disease or other harmful agency.

leuceine (lū'sein) n. [Gk. leukos, white.] An amino-acid formed during decomposition of proteids.

leucine (lū'sĭn) n. [Gk. leukos, white.] Amino-caproic acid found as a constituent of pancreatic juice and of various tissues and organs, also of some plants; C_eH₁₃O₂N.

leucism (lū'sizm, loo-) n. [Gk. leukos, white.] The presence of white plumage or pelage in animals with pigmented eyes and skin.

leucite (lū'sīt) n. [Gk. leukos, white.]
A colourless plastid.

leuco-,-also leuko-.

leucoblast (lū'köblăst, loo-) n. [Gk. leukos, white; blastos, bud.] A colourless blood-corpuscle in development.

leucocarpous (lū'kokâr'pus, loo-) a. [Gk. leukos, white; karpos, fruit.]

With the fruit white.

leucocyan (lū'kösī'ān, loo-) n. [Gk. leukos, white; kyanos, dark blue.]
A pigment found in certain algae.

leucocyte (lū'kösīt, loo-) n. [Gk. leukos, white; kytos, hollow.] An amoebocyte; a colourless blood-corpuscle; leukocyte.

leucocytogenesis (lū'kösī'tojēn'ësīs, loo-) n. [Gk. leukos, white; kytos, hollow; genesis, descent.] Leucocyte formation; leucopoiesis.

leucophore, leukophore (lū'köfōr, loo-) n. [Gk. leukos, white; pherein, to bear.] A yellow-pigment-bearing cell; iridocyte.

leucoplastids (lū'köplăs'tidz, loo-) n. plu. [Gk. leukos, white; plastos, formed; idion. dim.] Colourless plastids from which amylo-, chloro-, and chromoplastids arise.

leucoplasts (lū'köplästs, loo-) n. plu. [Gk. leukos, white; plastos, formed.] Colourless granules of plant cytoplasm, opp. chromoplasts;

leucoplastids.

leucopoiesis (lū'köpoi'ēsīs, loo-) n.
[Gk. leukos, white; poiesis, making.]
The formation of white blood corpuscles.

leucopterin(e) (lū'kŏp'tĕrin, loo-) n.
[Gk. leukos, white; pteron, wing.]
A white wing pigment of certain
Lepidoptera; C₁₉H₁₉O₁₁N₁₅.

leuko-, -see leuco-.

levator (leva'tor) n. [L. levare, to raise.] A name given to muscles serving to raise an organ or part. Opp. depressor.

levulose,—laevulose, q.v.

Leydig's cells [F. von Leydig, German anatomist]. Cells in testicular interstitial tissue.

Leydig's duct,—the Wolffian duct.

Leydig's organs, -minute organs on antennae of arthropods, supposed to be organs of smell.

liana (lian'a), liane (lian') n. [F. liane, from L. ligare, to bind.] Any luxuriant woody climbing plant of tropical or semitropical forests.

Lias (li'ás) n. [layers.] Marine and estuarine deposits of Jurassic period, containing remains of cycads, insects, ammonites. saurians, and other fossils.

liber (lī'ber) n. [L. liber, inner bark.] Inner bark; bast.

libido (libī'do) n. [L. libido, desire.]
Excitation within body associated with instinct; sexual energy; psychic energy; horme, q.v.; élan vital.

libriform (li'brifôrm) a. [L. liber, inner bark; forma, shape.] Resembling bast; appl. fibres, woody, later becoming septate.

lichenism (lī'kenizm) n. [Gk. leichen, lichen.] Symbiotic relationship

between fungi and algae.

lichenoid (li'kėnoid) a. [Gk. leichen, hchen; cidos, form.] Resembling a lichen.

lichenology (lī'kënôl'öjĭ) n. [Gk. leichen, lichen; logos, discourse.] The study of lichens.

Lieberkühn's crypts [J. N. Lieberkühn, German anatomist]. Tubular glands of the small intestine.

Liebig's law | J. von Liebig, German chemist|. The food element least plentiful in proportion to the requirements of plants limits their growth; law of the minimum, q.v.

lien (lī'ën) n. [L. lien, spleen.] Spleen. lienal (līē'năl) a. [L. lien, spleen.] Pert. spleen; appl. artery, vein, nerve plexus.

lienculus (lien'kūlūs) n. [Dim. of L. lien, spleen.] An accessory spleen. lienogastric (lie'nögas'trīk) a. [L. lien, spleen; gaster, stomach.] Pert. spleen and stomach; appl. artery supplying spleen and parts of

supplying spleen and parts of stomach and pancreas; appl. vein of hepatic portal system. lienorenal,—see phrenicolienal.

life-cycle,--- the various phases through which an individual species passes to maturity.

ligament (lig'ămënt) n. [L. ligamentum, bandage.] Α strong fibrous band of tissue connecting two or more moveable bones.

ligamenta flava,—yellow elastic ligaments connecting laminae of adjoining vertebrae.

ligneous (lig'neus) a. [L. lignum, wood.] Woody; of nature of wood. lignescent (lignes'ent) a. [L. lignes-

cere, to become woody. Developing the characters of woody tissue.

lignicolous (lignik'olus) a. [L. lignum, wood; colere, to inhabit.] Growing on or in wood.

lignification (lig'nifikā'shun) n. [L. lignum, wood; facere, to form.] Wood-formation; thickening of plant cell-walls by deposition of lignin.

lignin (lǐg'nǐn) n. [L. lignum, wood.] A complex substance which, associated with cellulose, causes the thickening of plant cell-walls, and so forms wood.

lignivorous (ligniv'orus) a. [L. lignum, wood; vorare, to devour.] wood: appl. various Eating insects.

lignocellulose (lǐg'nösĕl'ūlōs) n. [L. lignum, wood; cellula, little cell.] Essential constituent of woody tissue, lignin and cellulose combined.

lignose (lig'nos) n. [L. lignum, wood.] A variety of cellulose.

ligula (lig'ūlă) n. [L. ligula, little tongue.] A band or taenia of white matter in dorsal wall of fourth ventricle; median structure between labial palps of insects; lobe of parapodium in certain annelids; ligule, q.v.; lingula. q.v.

ligulate (lig'ūlāt) a. [L. ligula, little tongue.] Having or pert. ligules; strap-shaped, as ray florets of

Compositae.

ligule (lig'ūl) n. [L. ligula, little tongue.] A membranous outgrowth at junction of blade and leaf-sheath or petiole; small scale on upper surface of leaf-base in Lepidodendreae, Selaginellaceae, a tongue-shaped and Isoètes; corolla, as of certain florets.

liguliflorous (lǐg'ūlǐflō'rūs) a. ligula, little tongue; flos, flower.] Having ligulate flowers only.

limacel (lim'asel) n. [F., from L. limax, slug.] Concealed vestigial shell of slugs.

limaciform (limăs'ifôrm), limacine [L. limax, slug; (līm'āsin) a. forma, shape.] Like a slug; slugshaped.

[A.S. lim, limb.] limb (lim) n. Branch; arm; leg; wing; expanded portion of unguiculate petal.

limbic (lim'bik) a. [L. limbus, border.] Bordering; appl. a cerebral lobe, including hippocampal and cingulate gyri.

limbous (lim'bus) a. [L. limbus, border.] Appl. overlapping sutures. limbus (lim'būs) n. [L. limbus, border.] Any border if distinctly marked off by colour or structure.

limen (lī'men) n. [L. limen, threshold.] Threshold, minimum stimulus, or quantitative difference in stimulation, that is perceptible; boundary, as between vestibule of nostril and nasal cavity, limen

limicolous (līmik'ölūs) a. [L. limus, mud; colere, to dwell.] Living in

liminal (lim'inal) a. [L. limen, threshold.] Pert. a threshold; appl. stimulus; appl. sensation.

limited,--appl. chromosomes germinal, not in somatic, nuclei.

limivorous (līmiv'örus) a. [L. limus, mud; vorare, to devour.] Mud-eating; appl. certain aquatic animals which swallow mud.

limnetic (limnět'ĭk) a. [Gk. limne, marsh.] Living in, or pert., marshes or lakes.

limnocryptophyte (lim'nokrip'töfit) n. [Gk. limne, marsh; kryptos, hidden; phyton, plant. A helophyte or marsh plant; limnophyte.

- limnology (limnot oji) n. [Gk. limne. marshy lake; logos, discourse.] Science dealing with biological and other phenomena pert. inland waters; the study of standing waters.
- limnophilous (limnof'ilūs) a. [Gk. limne, marsh; philein, to love.] Living in fresh-water marshes; also limnobiotic.

limnophyte (lim'nöfit) n. [Gk. limne, marshy lake; phyton, plant.] A

pond plant; a helophyte.

limnoplankton (lim'nöpläng'ktön) n. [Gk. limne, marshy lake; plang-ktos, wandering.] The floating animal and plant life in fresh-water lakes, ponds, and marshes. Opp. haloplankton.

limosphere (lī'mösfēr) n. [Gk. limen, receptacle; sphaira, globe.] A spherical body containing a vacuole, situated near blepharoplast in spermiogenesis of some mosses.

linea (lǐn'ēă) n. [L. linea, line.] A line-like structure or mark.

linear (lin'ëăr) a. [L. linea, line.]

Pert. or in a line; tape- or threadlike; asthenic, appl. constitutional
type.

linear-ensate,—between linear and ensiform in shape.

linear - lanceolate,—between linear and lanceolate in shape.

linear-oblong, between linear and oblong in shape.

linellae (līněl'ē) n. plu. [L. linella, fine thread.] A system of filaments in certain Sarcodina holding together the xenophya.

lineolate (lin'ëölāt) a. [L. linea, line.]
Marked by fine lines or striae.

lingua (ling'gwă) n. [L. lingua, tongue.] The floor of mouth in mites; hypopharynx of insects; a tongue, or tongue-like structure.

lingual (ling'gwăl) a. [L. lingua, tongue.] Pert. tongue; appl. artery, gyrus, nerve, vein, etc.; appl. radula of molluscs.

lingula (ling'gūlā) n. [L. lingula, little tongue.] A small tongue-like process of bone or other tissue, as of cerebellum or sphenoid; a genus of brachiopods; ligula, q.v.

lingulate, -ligulate, q.v.

linin (li'nin) n. [L. linum, flax.] The substance of achromatinic or oxyphilic reticulum of cell-nucleus.

lininoplast (lin'inöplast) n [L. linum, flax; Gk. plastos, moulded.] Plasmosome, q.v.

linkage (ling'këj) n. [A.S. hlince, link.] Tendency of certain hereditary factors to remain associated through several generations; gametic coupling.

Linnaean (linē'an) a. [C. Linné or Linnaeus, Swedish naturalist]. Pert. or designating the system of classification established by Linnaeus.

linneon (lǐnē'ŏn) n. [C. Linné, Swedish naturalist]. Linnaean or taxonomic species.

lipase (lip'ās) n. [Gk. lipos, fat.] A lipolytic or fat-splitting enzyme, found in blood and in various plant and animal organs, and in various seeds

lip-cell,—a sporangium cell at the point of dehiscence.

lipides (lip'idz) n. plu. [Gk. lipos, fat.] Heterogeneous compounds soluble in fats and their solvents, including fats, waxes, chromolipides, sterols, glycolipides, phospholipides; lipids; cf. lipoids.

lipines (lip'inz) n. plu. [Gk. lipos, fat.] Compound lipides, including phospholipides and cerebrosides; lipins.

lipochondria (līp'ökŏn'driă) n. plu. [Gk. lipos, fat; chondros, grain.] Lipoid granules in the Golgi zone; Golgi presubstance.

lipochrin (līpō'krīn) a. [Gk. lipos, fat; ochros, sallow.] Appl. yellow lipoid droplet, fading by light, in unpigmented base of retinal cell

lipochromes (lǐp'ökrōmz) n. plu. [Gk. lipos, fat; chroma, colour.] A more or less indefinite group of plant and animal pigments, as carotins, luteins, chlorophane, rhodophane, xanthophane, zoonerythrin, etc.

lipoclastic (lĭp'öklăs'tik) a. [Gk. lipos, fat; klastos, broken.] Fat-splitting; lipolytic; appl. enzymes. lipogastry (li'pögăs'tri) n. [Gk. leipesthai, to be lacking; gaster, stomach.] Temporary obliteration of gastral cavity, as in some sponges.

lipogenous (lǐpŏj'ēnūs) a. [Gk. lipos, fat; genos, birth.] Fat-producing.

- lipohumour (lip'ohū'mor) n. [Gk. lipos, fat; L. humor, moisture.] A fat-soluble substance produced by nerves and acting on chromatophores.
- lipoid (lip'oid) a. [Gk. lipos, fat; eidos, form.] Resembling a fatty substance. n. A lipide; one of various fat-soluble substances occurring in plants and animals, as a sterol, carotene, terpene.

lipolytic (lip'olit'ik) a. [Gk. lipos, fat; lyein, to dissolve.] Capable of dissolving fat; fat-reducing.

- lipomerism (lipom'erizm) n. [Gk. leipesthai, to be lacking; meros, part.] Suppression of segmentation, or coalescence of segments, as in crustaceans.
- lipopalingenesis (lī'popāl'injēn'ēsis)

 n. [Gk. leipesthai, to be lacking;
 palin, anew; genesis, descent.]

 The omission of some stage or stages in phylogeny.

lipophore (lip'ofor) n. [Gk. lipos, fat; -phoros, -bearing.] A wandering cell originating in neural crest and containing a lipochrome.

- lipostomy (līpŏs'tömi) n. [Gk. leipesthai, to be lacking; stoma, mouth.] Temporary obliteration of mouth or osculum.
- lipotropic (lipotrop'ik) a. [Gk. lipos, fat; trope, turn.] Influencing fat metabolism; accelerating removal of fat.
- lipoxenous (lipözěn'ús, lipŏk'sėnŭs)

 a. [Gk. lipein, to abandon; xenos, host.] Leaving the host before completion of development.
- lirella (lĭrĕl'a) n. [L. lira, furrow.]
 A linear apothecium of lichens.
- lissencephalous (lĭs'ĕnkĕf'ālŭs, -sĕf-)
 a. [Gk. lissos, smooth; engkephalos,

brain.] Having few or no convolutions of the brain; lissencephalic.

- lissoflagellate (lĭs'öflāj'ēlāt) a. [Gk. lissos, smooth; L. flagellum, whip.] Having no collar surrounding base of flagellum.
- lithite (lith'it) n. [Gk. lithos, stone.] A calcareous secretion found in connection with ear, or with otocysts, lithocysts, and tentaculocysts, sensory organs of many invertebrates.
- lithocarp (lith'okârp) n. [Gk. Inthos, stone; karpos, fruit.] Fossil fruit or carpolith.
- lithocysts (lith'ösists) n. plu. [Gk. lithos, stone; kystis, bladder.] Minute sacs or grooves, containing lithites, found in various invertebrates; enlarged cells of plant epidermis, in which cystoliths are formed.
- lithodesma (lǐth'odēs'mā) n. [Gk. lithos, stone; desma, bond.] A small plate, shelly in nature, found in certain bivalves.
- lithodomous (lithŏd'omŭs) a. [Gk. lithos, stone; demein, to build.]
 Living in rock-holes or clefts.
- lithogenous (lithŏj'enŭs) a. [Gk. lithos, stone; -genes, producing.] Rock-forming, or rock-building, as certain corals.
- lithophagous (lithöf'águs) a. [Gk. lithos, stone; phagein, to eat.] Stone-eating, as birds; rock-burrowing, as some molluses and sea-urchins.
- lithophilous (lithòf'ilūs) a. [Gk. /ithos, stone; philein, to love.] Growing on stones or rocks; saxicoline.
- lithophyll (lith'ofil) n. [Gk. lithos, stone; phyllon, leaf.] A fossil leaf, or leaf-impression.
- lithophyte (lith'ofit) n. [Gk. lithos, stone; phyton, plant.] Plant growing on rocky ground
- lithosere (līth'ösēr) n. [Gk. lithos, stone; L. serere, to put in a row.] A plant succession originating on rock surfaces.
- lithotomous (lithot'ömus) a. [Gk. lithos, stone; temnein, to cut.] Stone-boring, as certain molluscs.

- litoral (lit'orăl) a. [L. litus, seashore.] Growing or living at or near the sea-shore, appl. zone between high and low water marks; also appl. cells, fixed macrophages, lining sinuses of reticular tissues and the wall of lymph channels; littoral.
- Littré's glands [A. Littré, French surgeon]. Urethral mucous glands. liver (liv'ër) n. [A.S. lifer, liver.] The bile-secreting gland of vertebrates; digestive gland of some invertebrates.
- liver-pancreas,—an organ in molluses and crustaceans, combining functions of liver and pancreas.
- lobar (lō'băr) a. [L.L. lobus, lobe.] Of or pert. a lobe.
- lobate (lō'bāt) a. [Gk. lobos, lobe.] Divided into lobes; lobose.
- lobe (lōb) n. [L.L. lobus, from Gk. lobos, lobe.] Any rounded projection of an organ; lobus; a flap-like structure on toes of certain birds.
- lobed (lobd) a. [Gk. lobos, lobe.] Having margin cut up into rounded divisions by incisions which reach less than half-way to mid-rib.
- lobopodia (lŏb'öpō'dĭā) n. plu. [Gk. lobos, lobe; pous, foot.] Blunt pseudopodia of Protozoa.
- **lobose** (lŏb'ōs) a. [Gk. lobos, lobe.] Divided into lobes; lobate.
- lobular (lŏb'ūlăr) a. [Gk. lobos, lobe.] Like or pert. small lobes.
- lobulate (lŏb'ūlāt) a. [Gk. lobos, lobe.] Divided into small lobes
- lobule (löb'ūl) n. [Dim. of L.L. lobus, lobe.] A small lobe or subdivision of a lobe; lobulus.
- lobus (lö'bus) n. [L.L. lobus, lobe.] Lobe; portion of an organ, as of glands and brain, delimited by fissures or septa.
- local sign,—characteristic quality of a tactile or other sensation associated with point of stimulation.
- localisation (lō'kālīzā'shūn) n. [L. localis, local.] Determination of a position; restriction to a limited area; restriction of pairing and chiasma formation at pachytene to one part of the chromosome.

- localisation of function,—reference to different parts of brain as communicating centres of various senses.
- localisation of sensation,—identification on surface of body of exact spot affected.
- locellus (lösěl'ŭs) n. [L. locellus from locus, place.] A small compartment of an ovary.
- lociation (lŏsiā'shūn) n. [L. locus, place.] Local differences in abundance or proportion of dominant species; local faciation.
- locomotor rods,—hooked or knobbed rods for crawling, on ventral surface of certain Nematoda.
- locular (lŏk'ūlăr) a. [L. loculus, little place.] Containing, or composed of loculi; loculate.
- locule, loculus.
- loculicidal (lŏk'ūlĭsī'dăl) a. [L. loculus, compartment; caedere, to cut.] Dehiscent dorsally down middle of
- carpels.

 loculus (lŏk'ūlŭs) n. [L. loculus, compartment.] A small chamber or cavity; cavity of an ovary or of an anther; cavity between septa in certain Coelenterata; chamber of foraminiferal shell.
- locus (lō'kūs) n. [L. locus, place.] Position of gene in the chromosome; location of a stimulus.
- locusta (lokŭs'tă) n. [I.. locusta, locust.] Spikelet of grasses; a locust.
- lodicule (lŏd'ikūl) n. [L. lodicula, coverlet.] A scale at base of ovary in grasses, supposed to represent part of a perianth.
- lodix (lō'diks) n. [L. lodix, blanket.] A ventral sclerite of seventh abdominal segment, covering genital plate, in Lepidoptera.
- logotype (lŏg'otīp) n. [Gk. logos, word; typos, pattern.] A genotype by subsequent designation, not originally described as such.
- loma (lô/ma) n. [Gk. loma, hem.] A thin membranous flap forming a fringe round an opening; fringe of toe in birds.

- lomastome (lō'măstōm) a. [Gk. loma, hem; stoma, mouth.] Having margin of lip recurved or reflected. loment,—lomentum.
- lomentaceous (lō'mentā'shūs) a. [L. lomentum, bean meal.] Pert., resembling, or having lomenta.
- lomentum (löměn'tům) n. [L. lomentum, bean meal.] A legume or pod constricted between seeds; loment.
- long-day,-appl. plants in which the flowering period is hastened by a relatively long photoperiod, ordinarily more than 12 hours.
- longicorn (lŏn'jĭkôrn) a. [L. longus, long; cornu, horn.] Having long antennae; appl. certain beetles.
- longipennate (lŏn'jipĕn'āt) a. [L. longus, long; penna, wing.] Having long wings, or long feathers.
- longirostral (lön'jĭrŏs'trăl) a. [L. longus, long; rostrum, beak.] With a long beak; longirostrate.
- loph (lof) n. [Gk. lophos, crest.] Crest which may connect cones in teeth and so form a ridge.
- lophiostomate (lŏf'iŏs'tomāt) a. [Gk. lophion, small crest; stoma, mouth.] With crested conceptacle-opening.
- lophobranchiate (löf'obrăng'kiāt) a. [Gk. lophos, crest; brangchia, gills.] With tufted gills.
- lophocaltrops (löf'ökäl'tröps) n. [Gk. lophos, crest; A.S. coltraeppe, kind of thistle.] A sponge spicule with rays crested or branched.
- lophocercal (lŏf'ösër'kăl) a [Gk. lophos, crest; kerkos, tail.] Having a rayless caudal fin like a ridge round end of vertebral column.
- lophodont (löf'ödönt) a. [Gk. lophos, crest; odous, tooth.] Having transverse ridges on the cheek-teeth grinding surface.
- lophophore (löf'öför) n. [Gk. lophos, crest; pherein, to carry.] A horseshoe-shaped tentacle-supporting organ in Polyzoa and Brachiopoda.
- lophoselenodont (löf'ösĕlē'nödönt)
 a. [Gk. lophos, crest; selene, moon; odous, tooth.] Having cheek-teeth ridged with crescentic cuspid ridges on grinding surface.

- lophosteon (lofôs'teŏn) n. [Gk. lophos, crest; osteon, bone.] The keel-ridge of a sternum.
- lophotriaene (löf'ötrī'ēn) n. [Gk.
 lophos, crest; triaina, trident.]
 Lophocaltrops, q.v.
- lophotrichous (löföt'rikus) a. [Gk. lophos, tuft; thrix, hair.] Having long whip-like flagella; with a tuft of flagella at one pole; appl. bacteria; lophotrichate, lophotrichic.
- loral (lô'răl) a. [L. lorum, thong.]

 Pert. or situated at the lore.
- lorate (lō'rāt) a. [L. lorum, thong.] Strap-shaped.
- lore (lor) n. [L. lorum, thong.]
 Space between bill and eyes in birds.
- lorica (lōrī'kă, lōr'ikă) n. [L. lorica, corselet.] A protective external case found in rotifers, infusorians, and diatoms.
- loricate (lō'rikāt) a. [L. lorica, corselet.] Covered with protective shell or scales.
- lorication moment,—the occasion of deposition of silica or calcium carbonate for an entire skeleton at one time; dictyotic moment.
- lorulum (lo'rūlum) n. [L. dim. of lorum, thong.] The small strapshaped and branched thallus of certain lichens.
- lorum (lō'rūm) n. [L. lorum, thong.] The piece of under jaw on which submentum lies in certain insects; dorsal plate protecting pedicle in spiders.
- lotic (lō'tik) a. [L. lotum, flowed against.] Appl. or pert. running water; living in brook or river; opp. lentic.
- Louis, angle of [A. Louis, French surgeon]. Angulus Ludovici or sternal angle.
- loxodont (lök'södönt) a. [Gk. loxos, oblique; odous, tooth.] Having molar teeth with shallow grooves between the ridges.
- luciferase (loos'iferas) n. [L. lux, light; ferre, to carry.] An oxidising enzyme which acts on luciferin, causing luminescence; photogenin.

luciferin (loos'iferin) n. [L. lux, light; ferre, to carry.] Intracellular or extracellular substance oxidised by luciferase, causing luminescence; photophelein.

lumbar (lum'bar) a. [L. lumbus, loin.] Pert. or near the region of the loins; appl. artery, vein, ver-

tebrae, plexus, gland.

lumbocostal (lum'bokos'tal) a. [L. lumbus, loin; costa, rib.] loins and ribs; appl. arch, liga-

lumbosacral (lum'bosa'kral) a. [L. lumbus, loin; sacrum, sacred.] Pert. loins and sacrum; appl. nerve and trunk, plexus.

lumbriciform (lumbris'ifôrm), lum-

(lŭm'brĭkoid) bricoid lumbricus, earth-worm; forma, shape; Gk. eidos, form.] Like a worm.

lumen (lū'men, loo-) n. [L. lumen, light. The cavity of a tubular part or organ; central cavity of a plant

luminescent organs, -- specialised organs for the production of light, found in various plant and animal organisms.

lunar (lū'năr, loo-) a. [L. luna, moon.] Appl. carpal bone, os lunare or lunatum, also called semilunar and intermedium;

lunate (lū'nāt, loo-) a. [L. luna, moon.] Somewhat crescent-shaped, semilunar.

lunatum,—semilunar bone.

lunette (lūnět') n. [F. lunettes, spectacles.] Transparent lower eyelid of snakes.

lung (lung) n. [A.S. lunge, lung.] The paired or single respiratory organ of air-breathing higher animal forms.

lung-book,—the respiratory organ of Scorpionidae, formed like a purse with numerous compartments.

lunula (lū'nūlă, loo-) n. [L. lunula, small moon.] Lunule.

lunular (lū'nūlăr, loo-) a. [L. lunula, small moon.] With crescentshaped marking; lunulate.

lunule (lū'nūl, loo-) n. [L. lunula, small moon.] A crescent-shaped structure or marking; lunula; small crescentic sclerite, the frontal lunule, above antennal bases in certain Diptera; white opaque portion of nail near root.

lunulet (lū'nūlět, loo-) n. [L. lunula, small moon.] A small lunule.

lupulin (lū'pūlĭn, loo'pūlĭn) n. [L. lupus, hop.] The resinous glandular scales of hops; an organic compound, bitter and acrid, obtained from these; C₂₆H₃₈O₄.

luteal (lū'teăl, loo-) a. [L. luteus, orange-yellow.] Pert. or like cells of corpus luteum; appl. lutein and paralutein cells; appl. hormone: progesterone.

lutein (lū'tein, loo-) n. [L. luteus, orange-yellow.] The yellow lipochrome pigment of egg-yolk and corpus luteum; C40H56O2.

lutein cells,—modified granulosa cells during formation of corpus luteum; follicular lutein cells, opp. lutein cells of theca interna.

luteination (lū'teinā'shun) n. luteus, orange-yellow.] luteum formation; luteinisation.

luteinisation (lū'teinīzā'shun, loo-) n. [L. luteus, orange-yellow.] The formation of corpus luteum.

luteinising hormone,—a pituitary hormone which stimulates thecalutein cell formation and interstitial cells of testis; LH, prolan B.

proluteosterone,—progesterone, gestin.

luteotrophic (lū'tëötrŏf'ĭk) a. luteus, orange-yellow; Gk. trophe, Appl.nourishment.] hormone which assists in maintaining corpus luteum, and may also be lactogenic.

lychnidiate (liknid'iāt) a. [Gk. lychnidion, small lamp.] Luminous.

lycopene (lī'köpēn) n. [L.L. lycopersicum, tomato, from Gk. lykopersikon.] The red carotenoid pigment of fruits of tomato, rose, etc.; lycopin; $C_{40}H_{56}$.

lygophil (lī'göfil) a. [Gk. lyge, shadow; philos, friend.] Prefer-

ring shade or darkness

lymph (limf) n. [L. lympha, water.]
An alkaline colourless fluid contained in lymphatic vessels.

lymph heart,—contractile expansion of a lymph vessel where it opens into a vein, in many vertebrates.

lymphatic (limfāt'ik) a. [L. lympha, water.] Pert. or conveying lymph. lymphocyte (lim'fösīt) n. [L. lympha, water; Gk. kytos, hollow.] A

small mononuclear colourless cor-

puscle of blood and lymph.

lymphogenic (lǐm'fojěn'ík) a. [L. lympha, water; Gk. -genes, producing.] Produced in lymph-glands. lymphogenous (lǐmfōj'énūs) a. [L. lympha, water; Gk.-genes, producing.] Lymph-forming.

lymphoid (lim'foid) a. [L. lympha, water; Gk. eidos, form.] Appl. retiform tissue with meshes largely occupied by lymph corpuscles;

adenoid.

lymphoidocyte (lǐm'foidosīt) n. [L. lympha, water; Gk. eidos, form; kytos, hollow.] Haemocytoblast.

lymphomyelocyte (lim'fomī'elosīt) n.
[L. lympha, water; Gk. myelos, marrow; kytos, hollow.] Myeloblast.
lyochromes (lī'ökrōmz) n. plu. [Gk.

lyein, to loose; chroma, colour.]
Water-soluble yellow cell pigments, or flavins, including vitamin B₂.

lyocytosis (lī'osītō'sīs) n. [Gk. lyein, to loose; kytos, hollow.] Histolysis by extra-cellular digestion, as in insect metamorphosis.

Lyonnet's glands,—paired accessory silk glands in lepidopterous larvae;

Filippi's glands.

lyophil (lī'öfil) a. [Gk. lyein, to loose; philos, loving.] Appl. solutions which, after evaporation to dryness, go readily into solution again on addition of fluid; cf. lyophobe.

lyophobe (li'öföb) a. [Gk. lyein, to loose; phobos, fear.] Appl. solutions which, after evaporation to dryness, remain as a solid; cf.

lyophil.

lyotropic (liötröp'ik) a. [Gk. lyein, to loose; trope, turn.] Appl. solutions which are dependent on changes in the solvent itself. lyra (li'ră) n. [Gk. lyra, lyre.] Triangular lamina or psalterium joining lateral parts of fornix, marked with fibres as a lyre; a lyrate pattern as on some bones; a series of chitinous rods forming part of the stridulating organ in certain spiders.

lyrate (li'rāt) a. [Gk. lyra, lyre.]
Lyre-shaped; appl. certain leaves.
lyriform (li'rĭfôrm) a. [L. lyra, lyre;
forma, shape.] Lyre-shaped; appl.
a sensory organ, the lyra, in

spiders.

265

lysactinic (līsāktīn'īk) a. [Gk. lysis, loosing; aktis, ray.] Of Stelleroidea, having podia limited to lower half of body instead of continued to apical plates; cf. desmactinic.

lysigenous (līsij'ēnūs) a. [Gk. lysis, loosing; -genes, producing.] Appl. formation of tissue cavities caused by degeneration and breaking down of cell-walls in centre of mass.

lysin (lī'sīn) n. [Gk. lysis, loosing.] Any substance capable of causing dissolution or lysis of cells or bacteria.

lysine (lī'sēn) n. [Gk. lysis, loosing.]
A diamino-acid, constituent of some plant proteins, a dietary factor, and cleavage product of certain animal proteins; C₆H₁₄O₂N₂.

lysogenesis (lī'sojen'esis) n. [Gk. /ysis, loosing; genesis, descent.]

The action of lysins. lysogenous, -lysigenous.

lytta (lit'ā) n. [Gk. lytta, madness.]
A vermiform structure of muscle, fatty and connective tissue, or cartilage, under the tongue of mammals; lyssa or rabies; cantharis, a blister-beetle.

M

macerate (măs'erāt) v. [L. macerare, to soften.] To wear away or to isolate parts of a tissue or organ; to soften and wear away by digestion or other means.

- machopolyp (măk'öpŏl'ĭp) n. [Gk. mache, fight; polys, many; pous, foot.] A nematophore, of certain Hydromedusae, provided with cnidoblasts or adhesive globules.
- macrandrous (măkrăn'drus) a. [Gk. makros, large; aner, man.] Having large male plants or elements.
- macraner (mākrā'nër) n. [Gk. makros, large; aner, male.] Male ant of unusually large size.
- macrergate (măkrčr'gāt) n. [Gk. makros, large; ergates, worker.] Worker ant of unusually large size. macro-,—a/so see mega-.
- macrobiotic (măk'robīot'ik) a. [Gk. makros, long; bios, life.] Long-lived; life-prolonging.
- macroblast (mak röblast) n. [Gk. makros, large; blastos, bud.] A large cell or corpuscle; a young normoblast.
- macrocarpous (măk'rökâr'pŭs) a. [Gk. makros, large; karpos, fruit.] Producing large fruit.
- macrocephalous (măk'rökēf'ālŭs, -sēf-) a. [Gk. makros, large; kephale, head.] Having the cotyledons thickened; big-headed
- macrocnemic (māk'röknē'mīk) a. [Gk. makros, large; kneme, tibia.] Appl. Zoanthidae having the sixth protocneme or primary pair of mesenteries perfect.
- macroconidium (măk'rökŏnĭd'iŭm)
 n. [Gk. makros, large; konis, dust; idion, dim.] A large asexual spore or conidium.
- macroconjugant (măk'rökŏn'joogănt) n. [Gk. makros, large; L. conjugare, to unite.] The larger individual of a conjugating pair.
- macrocyst (măk'rösist) n. [Gk. makros, large; kystis, bladder.] A large reproductive cell of certain fungi; a large cyst or case, as for spores.
- macrocytase (măk'rösī'tās) n. [Gk. makros, large; kytos, hollow.] The enzyme of macrophages or endothelial cells.
- macrodactylous (måk'rödåk'tǐlůs) a. [Gk. makros, long; daktylos, finger.] With long digits.

- macrodont (măk'rödŏnt) a. [Gk. makros, large; odous, tooth.] With large teeth.
- macroevolution (māk'röēvölū'shūn)

 n. [Gk. makros, large; L. evolvere,
 to unroll.] Evolutionary processes
 extending through geological eras;
 large-scale evolution of new genera
 and species owing to mutations
 resulting in marked changes in
 chromosome pattern and reaction
 system. Opp. microevolution.
- macrogamete (măk'rögamēt') n.
 [Gk. makros, large; gametes,
 spouse.] The larger of two conjugants, usually considered as equivalent to ovum or female conjugant.
- macrogametocyte (māk'rögāmē'tösīt) n. [Gk. makros. large; gametes, spouse; kytos, hollow.] The mother-cell of a macrogamete, considered female; term used mainly in connection with Protista.
- macrogamy (măkrŏg'ămĭ) n. [Gk. makros, large; gamos, marriage.] Syngamy between full-grown individuals of a species, as in Actinophrys; hologamy.
- macroglia (măkrōglī'ā) n. plu. [Gk. makros, large; glia, glue.] Astrocytes or true neuroglia; astroglia and oligodendroglia.
- macroglossate (măk'röglŏs'āt) a. [Gk. makros, large; glossa, tongue.] Furnished with a large tongue.
- macrognathic (māk'rönāth'ik) a. [Gk. makros, large; gnathos, jaw.] Having specially developed jaws.
- macrogonidium (māk'rögŏnīd'īum)

 n. [Gk. makros, large; gone,
 generation; idion, dim.] A large
 gonidium.
- macrogyne (măk'röjīnē) n. [Gk. makros, large; gyne, woman.] Female ant of unusually large size.
- macroleucocyte (măk'rölū'kösīt, -loo-) n. [Gk. makros, large; leukos, white; kytos, hollow.] A chromophil leucocyte, developed from a proleucocyte.
- macromere (mak'romer) n. [Gk. makros, large; meros, part.] In cleavage of telolecithal eggs, a larger cell of lower hemisphere.

macromerozoite (măk'röměrozô'īt) n. [Gk. makros, large; meros, part; zoon, animal.] One of many divisions produced by macroschizont stage of Sporozoa.

macromesentery (măk'röměs'entëri, -mez-) n. [Gk. makros, large; mesos, middle; enteron, gut. | One of the larger complete mesenteries of Anthozoa.

macromitosome (măk'romī'tösōm) n. [Gk. makros, large; mitos, thread; soma, body. The paranucleus, as

in Lepidoptera.

macromutation (mak'romuta'shun) n. |Gk. makros, large; L. mutare, to change. | Simultaneous mutation of a number of different charac-

macromyelon (māk'romī'elŏn) n. |Gk. makros. long; mvelos. The medulla oblonmarrow. gata.

macronotal (măk'ronō'tăl) a. [Gk. makros, large; noton, back. | With large thorax, as a queen ant.

macront (măk'rŏnt) n. [Gk. makros, The larger large; on, being. of two sets of cells formed after schizogony in Neosporidia, the macront giving rise to macrogametes.

macronucleocyte (măk'rönū'kleosīt) n. [Gk. makros, large; L. nucleus, kernel; Gk. kytos, hollow.] leucocyte having a relatively large nucleus; chromophil leucocyte of insects.

macronucleus (māk'rönū'kleus) n. [Gk. makros, large; L. nucleus, kernel.] The larger of two nuclei in a cell, usually supposed to be of a vegetative or somatic nature; meganucleus.

macrophage (măk'röfāj) n. [Gk. makros, large; phagein, to eat.] A large phagocytic cell, fixed or wandering; a large mononuclear leucocyte; a histiocyte, clasmatocyte, pericyte, etc.

(măkrŏf'ăgŭs) macrophagous [Gk. makros, large; phagein, to Feeding on relatively large masses of food; opp. microphagous. macrophyllous (māk'rofil'ŭs) a. [Gk. makros, large; phyllon, leaf.l Having large leaves or leaflets.

macropodous (măkrŏp'odus) a. [Gk. makros, long; pous, foot.] Having a long stalk, as a leaf or leaflet; having hypocotyl large in proportion to rest of embryo; long-footed.

macropterous (măkrop'terus) a. [Gk. makros, large; pteron, wing.] With unusually large fins or wings; fully

winged, opp. brachypterous.

macropyrenic (măk'ropīrē'nĭk) a. IGk. makros, large; pyren, fruit stone. | With nuclei markedly larger than average for the species or other group. n. A macropyrenic individual.

macroschizogony (măk'roskizog'-22. [Gk. makros, schizein, to cleave; gone, generation.] Method of multiplication of macroschizonts; schizogony giving rise to large merozoites.

macroschizont (măk'röskīz'ŏnt) n. [Gk. makros, large; schizein, to cleave; on, being.] Stage in lifecycle of certain Haemosporidia developed from sporozoite, and giving rise to macromerozoites.

macrosclereids (măk'rösklē'reidz) n. plu. [Gk. makros, large; skleros, hard; eidos, form.] Relatively large columnar sclereids, as in coat of certain seeds.

macroscopic (măk'röskŏp'ik) a. [Gk. makros, large; skopein, to view.] Visible by the naked eye.

macrosepalous (măk'rösĕp'ălŭs) a. [Gk. makros, large; F. sépale, sepal.] With specially large sepals.

macroseptum (măk'rösĕp'tŭm) n. [Gk. makros, large; L. septum, inclosure.] A primary or perfect septum of Anthozoa.

macrosiphon (măkrösī'fŏn) n. [Gk. makros, large; siphon, tube.] Large internal siphon of certain cephalopods.

macrosmatic (măk'rŏsmăt'ik) a. [Gk. makros, large; osme, smell.] With well-developed sense of smell.

macrosomatous (măk'rösō'mătŭs) a. [Gk. makros, large; soma, body.] Possessing abnormally large body.

- macrosome (māk'rosom) n. [Gk. makros, large; soma, body.] A large alveolar sphere or granule in protoplasm.
- macrosplanchnic (măk'rösplăngk'nĭk) a. [Gk. makros, large; splangchnon, entrail.] Large-bodied and short-legged.

macrosporangiophore (măk'rösporăn'jiöför) n. [Gk. makros, large; sporos, seed; anggeion, vessel; pherein, to bear.] A structure bearing a macrosporangium.

macrosporangium (mak'rosporan'jium) n. [Gk. makros, large; sporos, seed; anggeion, vessel.] A sporangium developing macrospores or megaspores.

macrospore (măk'röspōr) n. [Gk. makros, large; sporos, seed.] A large anisospore or gamete of Sarcodina; a larger spore of heterosporous plants; embryo-sac; megaspore.

macrosporophore (măk'röspō'röfōr)

n. [Gk. makros, large; sporos, seed; pherein, to bear.] A leafy lobe developing macrosporangia.

macrosporophyll (măk'rospō'rofil) n. [Gk. makros, large; sporos, seed; phyllon, leaf.] Macrosporophore; carpel.

macrosporozoite (māk'rospōr'ozō'īt)

n. [Gk. makros, large; sporos, seed;
zoon, animal.] A larger endogamous sporozoite of Sporozoa.

macrostomatous (māk'röstŏm'átūs)
a. [Gk. makros, large; stoma,
mouth.] With very large
mouth.

macrostylospore (măk'röstī'löspōr)

n. [Gk. makros, large; stylos,
pillar; sporos, seed.] A large
spore-like stalked body.

macrostylous (mäk'röstī'lūs) a. [Gk. makros, long; stylos, pillar.] With long styles.

macrotherm (māk'röthērm) n. [Gk. makros, large; therme, heat.] A tropical plant; macrothermophyte, megatherm.

macrotous (măkrō'tus) a. [Gk. makros, large; ous, ear.] With large ears. macrotrichia (măk'rötrĭk'ĭă) n. plu. [Gk. makros, large; thrix, hair.] The larger setae on body or wings of insects.

- macrotype (māk'rötīp) n. [Gk. makros, large; typos, a type.] A modified arrangement of mesenteries containing more macromesenteries than normal microtype, in Anthozoa.
- macrozoogonidium (māk'rözō'ögōnid'ĭūm) n. [Gk. makros, large;
 zoon, animal; gone, generation;
 idion, dim.] A large zoogonidium.
 macrozoospore (māk'rözō'öspōr) n.

[Gk. makros, large; zoon, animal; sporos, seed.] Large motile spore. macruric (makroor'ik) a. [Gk.

makros, long; oura, tail.] Long-tailed; macrural, macrurous.

macula (măk'ūlă) n. [L. macula, spot.] A spot or patch of colour; a small pit or depression; a tubercle.

macula cribrosa,—area on wall of vestibule of car, perforated for passage of auditory nerve filaments. macula germinitiva,—the germinal spot, nucleolus of an ovum.

macula lutea,—yellow spot of retina, an oval yellowish area in centre of posterior part of retina at point of most perfect vision.

macular (măk'ūlăr) a. [L. macula, spot.] Pert. a macula; pert. macula lutea.

maculate (măk'ūlāt), maculiferous (măk'ūlīf'ērūs), maculose (măk'ūlōs) a. [L. macula, spot.] Spotted.

maculation (māk'ūlā'shūn) n. [L. maculare, to spot.] The arrangement of spots on a plant or animal.

- madrepore (măd'rēpōr) n. [F. madrepore—from L. mater, mother; Gk. poros, friable stone.] A branching stony coral; plate at external opening of stone canal in echinoderms.
- madreporic (măd'rēpŏr'ik) a. [F. madrėpore, madrepore.] Pert. a madrepore or madreporite; appl. body, plate, tubercle, canal.

madreporic canal —stone canal, q.v.

- madreporite (mădrep'örīt) n. [F. madrepore, madrepore.] A flat circular or pentagonal grooved, perforated plate at end of an interambulacral area, or between two such areas in Echinoidea, or between rays in Asteroidea; a modified genital plate.
- Magendie's foramen [F. Magendie, French physiologist]. Median aperture in roof of fourth ventricle. connecting the latter with subarachnoid cavities.
- maggot (mag'ot) n. [M.E. magot, grub.] The worm-like insect larva, without appendages or distinct head, as that of the blow-fly.

maiosis, -- meiosis, q.v.

- mala (mā'lā) n. [L. mala, cheek.]
 Part of maxilla of some insects. of
 mandible of certain myriopods;
 part of exterior of lower jaw of
 birds; cheek; malar bone.
- malacoid (măl'ăkoid) a. [Gk. malakos, soft; eidos, form.] Soft in texture.
- malacology (măl'ăkŏl'öjī) n. [Gk. malakos, soft; logos, discourse.] The study of molluscs.
- malacophilous (māl'ākōf'īlūs) a.
 [Gk. malakos, soft; philein, to love.]
 Pollinated by agency of gastropods.
- malacopterous (māl'ākŏp'terūs) a. [Gk. malakos, soft; pteron. wing] Soft-finned.
- malacostracous (măl'ākŏs'trākūs) a. [Gk. malakos, soft; ostrakon, shell.] Soft-shelled.
- malar (mā'lār) a. [L. mala, cheekbone.] Pert. or in region of cheek.

 n. The jugal or zygomatic bone.
- malaxation (mălăksā'shūn) n. [Gk. malassein, to soften.] Compression of mandibles, or chewing, as by wasps.
- male (māl) a. [L. mas, male.] Pert. masculine organism; appl. organs of reproduction, as testes, or stamens; symbol 3.

male pronucleus,—nucleus of spermatozoon.

malleate (măl'eāt) a. [L. malleus, bammer.] Hammer-shaped; appl. a type of trophi of rotifer gizzard.

- malleoincudal (mål'eoĭnkū'dål) a. [L. malleus, hammer; incus, anvil.] Pert. malleus and incus of car.
- malleolar (mălē'olār) n. [L. dim. of malleus. hammer.] The vestigial fibula of ruminants. a. Pert. or in region of malleolus; appl. arteries. folds, sulcus.
- malleolus (mălē'olūs) n. [L. dam. of malleus, hammer.] Medial and lateral malleolus, lower extremity prolongations of tibia and fibula respectively; one of the club- or racket-shaped appendages on basal segments of hind legs of Solpugidæ.
- malleoramate (măl'eörā'māt) a. [L. malleus, hammer; ramus, branch.] Appl. type of trophi with looped manubrium and toothed incus in rotifer gizzard.
- malleus (māl'eūs) n. [L. malleus, hammer.] A part of rotifer mastax or gizzard; one of the chain of auditory ossicles of mammals; one of the Weberian ossicles of fishes.
- mallochorion (măl'okōr'iŏn) n. [Gk. mallos, wool; chorion, skin.] The primitive mammalian chorion.
- malloplacenta (māl'ōplāsēn'ta) n. [Gk. mallos. wool; L. placenta, flat cake] Non-deciduate placenta with villi evenly distributed, as in cetaceans and some ungulates.

Malpighian (mālpīg'īān) n. [M. Malpighi, Italian anatomist]. Discovered by or named after Malpighi.

- Malpighian body, or corpuscle,—in spleen, a nodular mass of lymphoid tissue ensheathing the smaller arteries; in kidney, a glomerulus of convoluted capillary blood-vessels enclosed in a dilatation of uriniferous tubule.
- Malpighian layer,—basal layer of epidermis next to true skin; rete Malpighii.
- Malpighian tubules,—thread-like excretory tubes leading into posterior part of gut of insects.
- maltase (môl'tās) n. [A.S. mealt, malt.] An enzyme which converts malt-sugar into grape-sugar.

- maltose (môl'tōs) n. [A.S. mealt, malt.] Malt-sugar, formed from starch by ptyalin and amylase; C₁₂H₁₂O₁₁.
- mamelon (mām'ēlŏn) n. [F. mamelon, from L. mamilla, nipple.]
 Small pimple-like structure in centre of tubercle of echinoid interambulacral plate; papilla forming nucellus in cycads.
- mamilla (mămĭl'ā) n. [L. mamilla, nipple.] A nipple; a nipple-shaped structure; mammilla.
- mamillary bodies,—corpora mamillaria or albicantia, q.v.
- mamillary process or tubercle,—superior tubercle connected with transverse process of lower thoracic vertebrae.
- mamillate (măm'ilāt) a. [L. mamilla, nipple.] Studded with small protuberances.
- mamma (măm'ă) n. [L. mamma, breast.] Milk-secreting organ of female mammals.
- mammal (mām'āl) n. [L. mamma, breast.] An animal of a class of vertebrates of which the females suckle the young.
- mammalogy (mămăl'öji) n. [L. mamma, breast; Gk. logos, discourse.] The study of mammals.
- mammary (măm'ări) a. [L. mamma, breast.] Pert. the breast; appl. arteries, veins, glands, tubules, etc.
- mammiferous (māmif'ërūs) a. [L. mamma, breast; ferre, to bear.] Developing mammae; milk-secreting; mammalian.

mammilla,—mamilla, q.v.

- mammogenic (măm'ŏjēn'īk) a. [L. mamma, breast; Gk. gennaein, to produce.] Appl. pituitary hormone complex which promotes growth of the lobe-alveolar and duct systems of the mammary gland.
- manchette (mănshět') n. [F. manchette, cuff.] Membrane enveloping the cytoplasm surrounding the axial filament of a spermatid.
- mandible (măn'dibl) n. [L. mandibulum, jaw.] The lower jaw of vertebrates, either a single bone or composed of several; a paired

- mouth appendage of arthropods; mandibulum.
- mandibular (măndĭb'ūlăr) a. [L. mandibulum, jaw.] Pert. the lower jaw; appl. arch, canal, foramen, fossa, nerve, notch.
- mandibulate (măndib'ūlāt) a. [L. mandibulum, jaw.] Having a lower jaw; having functional jaws; having mandibles.
- mandibuliform (măndib'ūlifôrm) a. [L. mandibulum, jaw; forma, shape.] Resembling, or used as a mandible; appl. certain insect maxillae.
- mandibulohyoid (mandib'ūlöhī'oid)

 a. [L. mandibulum, jaw; Gk.
 hyoeides, T-shaped.] In region of
 mandible and hyoid.
- mandibulomaxillary (măndib'ūlömăksil'ări) a. [L. mandibulum, jaw; maxilla, jaw.] Pert. maxillae and mandibles of arthropods.
- manducation (măn'dūkā'shūn) n. [L. manducare, to chew.] Chewing; mastication.
- manicate (măn'īkāt) a. [L. manicatus, sleeved.] Covered with entangled hairs or matted scales.
- manna (măn'ā) n. [Gk. manna, manna.] Hardened exudation of bark of certain trees; honey-dew secreted by certain Coccidae.
- mannose (măn'ōs) n. [Gk. manna, manna.] A sugar of various plants; $C_aH_{10}O_a$.
- manoxylic (mănozi'lik) a. [Gk. manos, slack; xylon, wood.] Having soft loose wood, as Cycadales. Opp. pycnoxylic.
- mantle (man'tl) n. [L. mantellum, cloak.] Outer soft fold of integument next shell of molluscs; pallium; sheath of spongoblast cells; body-wall of ascidians; scapulars and wing coverts of birds; ocrea, q.v.
- mantle cavity,—a space between the mantle and body proper.
- mantle cell,—a cell of tapetum or investing tissue of a sporangium.
- mantle fibres,—the spindle-fibres of a fully formed spindle.

mantle layer,—a layer of embryonic medulla spinalis representing the future gray columns.

mantle lobes,—dorsal and ventral flaps of mantle in bivalves.

manual (măn'ūăl) n. [I.. manus, hand.] A wing-quill borne on manus of birds; remex primarius, primary feather.

manubrial (mănū'briăl) a. [L. manubrium, handle.] Pert. a manu-

brium; handle-shaped.

- manubrium (mănū'brĭūm) n. [L. manubrium, handle.] A cell projecting inwards from shield of an antheridial globule of thallophytes; a hypostome or conical elevation at distal end of a hydrozoan polyp; clapper-like portion hanging down from under surface of medusae; handle-like part of malleus of ear; handle of malleus of mastax; presternum or anterior part of sternum.
- manus (mān'ūs) n. [L. manus, hand.] Hand, or part of fore-limb corresponding to it, as found in vertebrates from Amphibia onwards.
- manyplies,—omasum or psalterium, third chamber of stomach of ruminants—so-called from its folded structure.

marcescent (marses'ent) a. [L. marcescere, to wither.] Withering but not falling off; appl. a calyx or corolla persisting after fertilisation.

marginal (mâr'jīnāl) a. [L. margo, edge.] Pert. at or near the margin, edge, or border; appl. a form of nervation; appl. a convolution of frontal lobe; appl. a type of placenta; appl. plates round margin of chelonian carapace.

marginalia (mâr'jĭnā'lĭa) n. plu. [L. margo, edge.] Prostalia or defensive spicules on body surface round

osculum.

marginate (mâr'jĭnāt) a. [L. margo, edge.] Having a distinct margin

in structure or colouring.

marginicidal (mâr'jīnĭsī'dăl) a. [L. margo, edge; caedere, to cut.] Dehiscing by line of union of carpels.

- marginiform (mar'jinifôrm) a. [I.. margo, edge: forma, shape.] Like a margin or border in appearance or structure.
- marginirostral (mâr'jīnĭrŏs'trăl) a. [L. margo, edge; rostrum, beak.] Forming the edges of a bird's bill.
- marita (mărī'tă, marē'tă) n. [L. maritus, conjugal.] Sexually mature stage in helminth life history.
- marital (măr'ītăl) a. [L. maritus, conjugal.] Pert. marita; producing fertilised eggs, appl. trematodes.
- marker,—an identifying factor; a gene of known location and effect which makes possible the determination of the distribution of other, less conspicuously effective, genes.

marmorate (mar'mörāt) a. [L. marmor, marble.] Of marbled appearance.

- marrow (măr'ō) n. [A.S. mearg, pith.] Connective tissue filling up cylindrical cavities in bodies of long bones, and spaces of cancellous tissue, differing in composition in different bones; medulla ossium; pith of certain plants; vegetable marrow.
- marsupial (mârsū'piāl) a. [L. marsupium, pouch.] Pert. a marsupium; pouch-bearing, as a kangaroo; appl. bones of pelvic girdle in certain mammals.
- marsupium (mârsū'pĭūm) n. [L. marsupium, pouch.] Any pouch-like structure in which the young of an animal complete their development, such as abdominal pouch of marsupials; gill cavities of bivalves; recess formed by diverging spines and a supporting membrane in stelleroids; structure protecting the acrocyst in Sertularia; a nursing-sac surrounding certain archegonia.
- mask (mäsk) n. [F. masque, mask.] A hinged prehensile structure, corresponding to adult labium, peculiar to dragon-fly nymph.
- masked (mas'kd) a. [F. masque, mask.] Personate, appl. corolla; concealed, appl. fat of cell which is not evident microscopically.

- massa intermedia,—grey matter connecting thalami across third ventricle; middle commissure.
- masseter (măsē'ter) n. [Gk. masseter, one that chews.] Muscle which raises lower jaw and assists in chewing.
- masseteric (măs'ētēr'ik) a. [Gk. masseter, one that chews.] Pert. or near masseter muscle of cheek; appl. artery, vein, nerve.
- massive (mas'īv) a. [L. massa, mass.] Bulky; heavy; compacted; appl. nuclei deficient in nuclear sap.
- massula (măs'ūlă) n. [L. massula, small mass.] A mass of microspores in a sporangium of certain pteridophytes; a massed group of microspores in orchids.
- mast cells,—spheroid or ovoid cells of very granular protoplasm, numerous in connective tissue where fat is being laid down; Mastzellen of Ehrlich.
- mastax (măs'tăks) n. [Gk. mastax, jaws.] The gizzard or pharyngeal mill of rotifers.
- mastication (măs'tikā'shŭn) n. [L. masticare, to chew.] Process of chewing food with teeth till reduced to small pieces or to a pulp.
- masticatory stomach,—the gastric mill or stomodaeal apparatus of crustaceans, for grinding and straining food material.
- mastidion (măstid'iŏn) n. [Gk. mastos, breast; idion, dim.] Nipple-like protuberance on paturon, in some spiders.
- mastigium (măstij'ium) n. [Gk. mastigion, little whip.] Defensive posterior lash of certain larvae.
- mastigobranchia (mās'tǐgöbrāng'kǐā) n. [Gk. mastix, whip; brangchia, gills.] Epipodite of adult Decapoda, a bilobed membranous lamina extending upwards between gills.
- mastigosome (măstī'gösōm') n. [Gk. mastix, whip; soma, body.] A blepharoplast.
- mastoid (măs'toid) a. [Gk. mastos, breast; eidos, form.] Nipple-shaped; appl. a process of temporal

- bone, cells, foramen, fossa, notch.
- mastoideosquamous (măstoid'ëöskwā'mŭs) a. [Gk. mastos, breast; eidos, like; L. squama, scale.] Pert. mastoid and squamous parts of temporal bone.
- mastoidohumeralis (măstoid'öhūmerā'līs) a. [Gk. mastos, breast; eidos, like; L. humerus, humerus.] Λ muscle of certain quadrupeds connecting mastoid and humerus.
- masto-occipital (măs'tö-ŏksīp'ităl) a. [Gk. mastos, breast; L. occiput, occiput.] Pert. occipital bone and mastoid process of temporal.
- mastoparietal (măs'töpărī'ētăl) a. [Gk. mastos, breast; L. paries, wall.] Pert. parietal bone and mastoid process of temporal.
- mastotympanic (măs'tötimpăn'îk) a. [Gk. mastos, breast; tympanon, drum.] Appl. part of tympanic cavity's boundary in certain reptiles.
- mating types,—groups, the individuals of which do not conjugate with individuals of other groups, as of ciliates.
- matriclinous (măt'rĭklī'nūs) a. [L. mater, mother; Gk. klinein, to bend.] With hereditary characteristics more maternal than paternal; matroclinic; matroclinal.
- matrix (mắt'riks) n. [L. mater, mother.] Ground substance of connective tissue; part beneath body and root of nail; uterus; body upon which lichen or fungus grows; envelope of chromatid.
- matroclinal,—matriclinous, q.v.
- mattula (māt'ūlā) n. [L. matta, mat.] Fibrous network covering petiole bases of palms.
- maturation (măt'ūrā'shŭn) n. [L. maturus, ripe.] Ripening; completion of germ-cell development, consisting of reduction of chromatin; meiosis; reduction of chromosomes from somatic or diploid to genetic or haploid number.
- Mauthner's cells [L. Mauthner, Austrian physician]. A layer be-

- tween medullary sheath and neurolemma of nerve fibre.
- maxilla (māksīl'ā) n. [L. maxilla, jaw.] The upper jaw; part of upper jaw behind premaxilla; an appendage of most arthropods, posterior to mandible, modified in various ways in adaptation to function and requirements.
- maxillary (măksil'ări) a. [L. maxilla, jaw.] Pert. or in region of maxilla or upper jaw; appl. artery, nerve, process, sinus, tuberosity, vein, etc.
- maxillary glands, paired renal organs opening at base of maxilla in Crustacea.
- maxilliferous (măk'silif'erūs) a. [L. maxilla, jaw; ferre, to carry.]
 Bearing maxillae.
- maxilliform (măksĭl'ĭfôrm) a. [L. maxilla, jaw; forma, shape.] Like a maxilla.
- maxillipede (mäksil'ĭpēd) n. [L. maxilla, jaw; pes, foot.] An appendage, in one, two, or three pairs, posterior to maxillae in arthropods; also maxilliped.
- maxillodental (măksil'oděn'tăl) a. [L. maxilla, jaw; dens, tooth.] Pert. jaws and teeth.
- maxillojugal (măksil'ojoo'găl) a. [L. maxilla, jaw; jugum, yoke.] Pert. jaw and jugal bone.
- maxillolabial (māksīl'olā'bīāl) a. [L. maxilla, jaw; labium, lip.] Pert. maxilla and labium; appl. dart in ticks.
- maxillomandibular (măksil'ömăndĭb'ūlăr) a. [L. maxilla, jaw; mandibulum, jaw.] Appl. arch forming jaws of primitive fishes; pert. maxilla and mandible.
- maxillopalatal (măksĭl'öpăl'ătăl) a. [L. maxilla, jaw; palatus, palate.] Pert. jaw and palatal bones; appl. a maxillary process of birds.
- maxillopharyngeal (măksĭl'öfărĭn'jeăl) a. [L. maxilla, jaw; Gk. pharyngx, gullet.] Pert. lower jaw and pharynx.
- maxillopremaxillary (măksil'öprēmāksil'ări) a. [L. maxilla, jaw;

- pre, before.] Pert. whole of upper jaw; appl. jaw when maxilla and premaxilla are fused.
- maxilloturbinal (māksil'ötūr'bīnāl)
 a. [L. maxilla, jaw; turbo, whorl.]
 Pert. maxilla and turbinals. n. A
 bone arising from lateral wall of
 nasal cavity, which supports sensory
 epithelium.
- maxillula (māksīl'ūlā) n. [L. dim. of maxilla, jaw.] A first maxilla in Crustacea when there are more pairs than one; an appendage between mandible and first maxilla in primitive insects.
- mazic (mā'zĭk) a. [Gk. maza, cake.] Placental; pert. placenta.
- M-chromosome, a microchromosome; or, a mediocentric chromosome.
- **M-disc,** -a line in middle of Hensen's line, q.v.
- meatus (mēā'tŭs) n. [L. meatus, passage.] A passage or channel, as acoustic, nasal, etc.
- mechanism (měk'ănĭzm) n. [Gk. mechane, machine.] The view that all vital phenomena are due to physical and chemical laws.
- mechanocyte (měk'ănösīt) n. [Gk. mechane, contrivance; kytos, hollow]. A cell derived from bone, cartilage, connective tissue, tendon, or muscle; a supporting cell; a fibrocyte.
- mechanoreceptor (měk'ánörēsěp'tŏr) n. [Gk. mechane, contrivance;
 L. recipere, to receive.] A specialised structure sensitive to contact,
 pressure, or gravity.
- Meckel's cartilage or rod [J. F. Meckel, junior, German anatomist]. The lower jaw of lower vertebrates, and in higher vertebrates, the axis round which membrane bones of jaw are arranged and formed.
- Meckel's ganglion [J. F. Meckel, senior, German anatomist]. The sphenopalatine ganglion.
- meconidium (mē'kōnid'iŭm) n. [Gk. mekon, poppy; idion, dim.] Sessile or pedicellate extracapsular medusa usually lying on top of gonangium of certain hydroids.

- meconium (mēkō'nĭūm) n. [Gk. mekon, poppy.] Waste products of a pupa or other embryonic form; contents of intestine of new-born mammal.
- media (mē'diā) n. [L. medius, middle.] A middle structure, such as a layer of tissue, a central nervure; plu. of medium, substances of various kinds in which cultures are grown; transmitting substances. mediad (mē'diād) adv. [L. medius, middle; ad, to.] Towards but not
- quite in the middle line or axis.

 medial (mē'dĭāl) a. [L. medius,
 middle.] Situated in the middle.
- median (mē'dĭān) a. [L. medius, middle.] Lying or running in axial plane; intermediate; middle. n. The middle variate when variates are arranged in order of magnitude.
- mediastinal (me'diasti'nal) a. [L. mediastinus, servant.] Pert. or in region of mediastinum; appl. cavity, arteries, glands, pleura.
- mediastinum (mē'diāstī'nūm) n. [L. mediastinus, servant.] Space between right and left pleura in and near median sagittal thoracic plane; incomplete vertical septum of testis, Highmore's body.
- mediator (me'diator) n. [L. medius, middle.] A nerve cell maintaining relation between receptor and effector.
- mediocentric (mē'dĭösĕn'trĭk) a. [L. medius, middle; centrum. centre.] Having a medial, or mediad, centromere; appl. chromosome.
- mediodorsal (mē'dĭödôr'săl) a. [L. medius, middle; dorsum, back.]
 In the dorsal middle line.
- mediopalatine (mē'diöpāl'ātīn) a. [L. medius. middle; palatus, palate.] Between palatal bones; appl. a cranial bone of some birds.
- mediopectoral (mē'diöpēk'törāl) a.
 [L. medius, middle; pectus, breast.]
 Appl. middle part of sternum.
- mediostapedial (mē'dĭöstāpē'dĭāl) n.
 [L. medius, middle; stapes, stirrup.]
 Pert. that portion of columella auris external to stapes.

- mediotarsal (mē'diötâr'săl) a. [I... medius, middle; tarsus, ankle.]
 Between tarsal bones.
- medioventral (mē'diövěn'trăl) a. [L. medius, middle; venter, belly.] In the middle ventral line.
- mediproboscis (mē'dīpröbŏs'īs) n. [L. medius, middle; Gk. proboskis, trunk.] Middle portion of insect proboscis, part of ligula.
- medithorax (mē'dīthō'rāks) n. [L. medius, middle; Gk. thorax, breast.] The middle part of thorax; the mesothorax of insects.
- medium (mē'dium) n. [L. medium, middle.] Any of the structures through which a force acts, as refracting media of eyebulb; substance in which cultures are reared or tissues propagated.
- medulla (mëdul'ā) n. [L. medulla, marrow, pith.] Marrow of bones; central part of an organ or tissue; pith or central portion of stem.
- medulla oblongata, posterior portion of brain continuous with medulla spinalis or spinal cord.
- medullary (mědůl'ári) a. [L. medulla, pith.] Pert. or in region of medulla; appl. axis, artery, lamina, membrane, bone, spaces, canal, etc.
- medullary canal,—hollow cylindrical portion of a long bone containing marrow; the neurocoel; neural tube
- medullary groove,—a groove on surface of medullary plate, bounded by folds which grow and coalesce, converting groove into a canal, the neurocoel.
- medullary keel,—a downward growth towards archenteron, the rudiment of central nervous system in development of certain primitive vertebrates.
- medullary layer,—a thick subcortical layer of the thallus of some lichens.
- medullary phloem, internal phloem in a bicollateral bundle, as in Cucurbitaceae.
- medullary plate,—plate-like formation of ectoderm cells bordering blastopore of early embryo; neural

plate, earliest rudiment of nervous system.

medullary rays—a number of strands of connective tissue extending between pith and pericycle.

medullary sheath,—a ring of protoxylem round pith of certain stems; a layer of white substance, composed of myelin, surrounding axis cylinder of medullated nervefibre.

medullary velum,—valve of Vieus-

sens, q.v.

medullated (měď úlātěd) a. [L. medulla, pith.] Provided with pith, or with a medullary sheath.

medullated nerve-fibres,—fibres of brain and spinal cord, consisting of axis-cylinder or neuraxis of primitive fibrillae, surrounded by medullary sheath, in turn covered by delicate neurilemma.

medullispinal (mëdül'ispī'năl) a. [L. medulla, pith; spina, spine.] Of the spinal cord.

medusa (mëdū'să) n. [Gk. Medousa, one who rules.] A jelly-fish.

medusiform (mëdū'sifôrm) a. [Gk. Medousa, Medusa; L. forma, shape.] Like a medusa or jelly-fish.

medusoid (mëdū'soid) n. [Gk. Medousa, Medusa; eidos, like.]
A medusa-like free-swimming gonophore of Hydrozoa. a. Like a jelly-fish or medusa.

medusome (mëdū'sōm) n. [Gk. Medousa, Medusa; soma, body.] Medusoid stage in life-history of Obelia.

mega-,--also see macro-.

megacephalic (měg'ákěfál'ík, -sěf-) a. [Gk. megas, large; kephale, head.] With abnormally large head; having a cranial capacity of over 1450 c.c.; cf. mesocephalic, microcephalic.

megagamete (měg'ăgămēt') n. [Gk. megas, large; gametes, spouse.]
A rounded cell regarded as an ovum or its equivalent, developed from a megagametocyte after a process akin to maturation; macrogamete.

megagametocyte (měg'ăgămē'tösīt) n. [Gk. megas, large; gametes, spouse; kytos, hollow.] A cell developed from a merozoite, and itself giving rise to a megagamete.

megagametogenesis (még'ágámétójén'ésis) n. [Gk. megas, great; gametes, spouse; genesis, descent.]
Development of megagametes or ova.

megakaryocyte (měg'ākār'iösīt) n. [Gk. megas, large; karyon, nut; kytos, hollow.] An amoeboid giant cell of bone-marrow, with one large annular lobulated nucleus, containing a number of nucleoli.

megalaesthetes (měg'álēsthēt'ēz) n. plu. [Gk. megalon, great; aisthetes, perceiver.] Sensory organs, sometimes in form of eyes, in Placo-

phora.

megalecithal,—see megalolecithal. megaloblast (meg'ălöblast) n. [Gk. megalos, greatly; blastos, bud.] A primitive large erythroblast.

megalogonidum (měg'álogŏnīd'iŭm)

n. [Gk. megalos, greatly; gonos, offspring; idion, dim.] A large gonidium.

megalolecithal (meg'aloles'ithal) a. [Gk. megalos, greatly; lekithos, yolk.] Containing much yolk, as telolecithal eggs; megalecithal.

megalopic (měg'álŏp'îk) a. [Gk. megalos, greatly; ops, eye.] Belonging to the megalops stage

ing to the megalops stage.

megalopore (meg'alöpor) n. [Gk. megalon, great; poros, channel.]

Pore in dorsal plates of Chiton, for placing a megalaesthete in direct communication with exterior.

megalops (měg'álŏps) n. [Gk. megalos, greatly; ops, eye.] A larval stage of certain Crustacea, as crabs, conspicuous by large stalked

eyes; megalopa.

megalospheric (měg'álösfěr'ík) a. [Gk. megalos, greatly; sphaira, globe.] Of polythalamous foraminifer shells, having a megalosphere or large initial chamber; megaspheric.

megamere (mēg'āmēr) n. [Gk. megas, large; meros, part.] One of the large cells formed after primary divisions of a developing ovum.

- megameric (měg'áměr'ík) a. [Gk. megas, large; meros, part.] With relatively large parts; appl. chromosomes with large heterochromatic regions; pert. megameres.
- meganephridia (meg'anefrid'ia) n.
 plu. [Gk. megas, large; nephros,
 kidney; idion, dim.] Large
 nephridia, occurring as one pair
 per segment; holonephridia.
- meganucleus (mēg'ānū'klēūs) n. [Gk. megas, large; L. nucleus, kernel.] The larger or vegetative nucleus of infusoria; macronucleus; trophonucleus.
- megaphanerophyte (měg'áfán'eröfit) n. [Gk. megas, large; phaneros, manifest; phyton, plant.] Tree exceeding 30 metres in height.
- megaphyllous (měg'áfil'ús) a. [Gk. megas, large; phyllon, leaf.] Having relatively large leaves.
- megasclere (měg'ásklēr) n. [Gk. megas, large; skleros, hard.] Skeletal spicule of general supporting framework of sponges. Opp. microsclere. megasome,—macrosome, q.v.
- megasorus (měg'ásō'rūs) n. [Gk. mega, large; soros, heap.] A sorus containing megasporangia, opp. microsorus.
- megasporangium (měg'áspörán'jium) n. [Gk. megas, large; sporos, seed; anggeion, vessel.] A macrospore-producing sporangium; ovule.
- megaspore (měg'áspōr) n. [Gk. megas, great; sporos, seed. A larger-sized spore of dimorphic forms in reproduction by spore-formation; larger spore of heterosporous plants, regarded as female; gynospore; embryo-sac cell of seed plant macrospore.
- megasporocyte (měg'áspō'rösît) n. [Gk. megas, large; sporos, seed; kytos, hollow.] The embryo-sac mother-cell, diploid cell in ovary that undergoes meiosis, producing four haploid megaspores.
- megasporophyll (meg'aspō'röfil) n. [Gk. megas, great; sporos, seed; phyllon, leaf.] A spore-bearing

- leaf developing megasporangia; carpel.
- megatherm (měg'áthěrm) n. [Gk. megas, great; therme, heat.] A tropical plant; a plant requiring moist heat.
- megazooid (měg'ázō'oid) n. [Gk. megas, great; zoon, animal; eidos, form.] The larger zooid resulting from binary or other fission.
- megazoospore (měg'ázō'öspōr) n. [Gk. megas, great; zoon, animal; sporos, seed.] A large zoospore, as in reproduction of certain Radiolaria; a zoogonidium of certain Algae.
- megistotherm (měj'Istöthěrm, měg-) n. [Gk. megistos, greatest; therme, heat.] A plant that thrives at a more or less uniformly high temperature.
- Mehlis' glands,—acinous glands surrounding the ootype; shell gland of trematodes.
- Meibomian glands [H. Meibom, German anatomist]. The tarsal glands, modified sebaceous glands of the eyelids, the ducts opening on the free margins.
- meiocyte (mī'ösīt) n. [Gk. meion, smaller; kytos, hollow.] A reproductive cell prior to meiosis; auxocyte, q.v.
- meiogenic (mi'ojen'ik) a. [Gk. meion, smaller; gene, descent.] Promoting nuclear division.
- meiolecithal (mī'olĕs'īthăl) a. [Gk. meion, less; lekithos, yolk.] Having little yolk, as homolecithal and isolecithal eggs.
- meiomery (miom'eri) n. [Gk. meion, smaller; meros, part.] Condition of having fewer than the normal number of parts.
- meiophylly (mī'öfīl'ī) n. [Gk. meion, smaller; phyllon, leaf.] Suppression of one or more leaves in a whorl.
- meiosis (miō'sis) n. [Gk. meion, smaller.] Process of reduction division of germ-cell chromosomes from diploid to haploid number at maturation; also maiosis.
- meiostemonous (mī'östěm'önus) a. [Gk. meion, smaller; stemon, spun thread.] Having fewer stamens than petals or sepals.

- meiotaxy (mi'ötäk'si) n. [Gk. meion, smaller; taxis, arrangement.] Suppression of whorl or set of organs.
- meiotic (miŏt'ik) a. [Gk. meion, smaller.] Appl. reduction division, pert. meiosis.
- Meissner's corpuscles [G. Meissner, German histologist]. Tactule corpuscles, associated with sense of pain, in skin of digits, lips, nipple, and certain other areas.
- Meissner's plexus, a gangliated plexus of nerve fibres in submucous coat of small intestine.
- melanin (měl'ánīn) n. [Gk. melas, black.] Black or dark-brown pigment; cf. eumelanin, phaeomelanin, dopa, haemozoin; C₇₇H₉₈O₃₃N₁₄S.
- melaniridosome (mel'anir'idosom) n. [Gk. melas, black; iris, rainbow; soma, body.] A pigment body consisting of a melanophore and associated iridocytes in corium of fishes.
- melanism (měl'ánĭzm) n. [(k. melas, black.] Excessive development of black pigment.
- melanoblast (měl'ánoblást) n. [Gk. melas, black; blastos, bud.] A cell of rete mucosum giving rise to melanin formation in the Malpighian layer of epidermis.

melanocyte (měl'ánösīt) n. [Gk. melas, black; kytos, hollow.] A black pigmented lymphocyte.

- melanophore (měl'ánofôr) n. [Gk. melas, black; pherein, to bear.] A black pigment cell.
- melanosoma (měl'ánosô'mă) n. [Gk. melas, black; soma, body.] Dark, pigment mass associated with ocellus, as in certain Dinoflagellata.
- melanospermous (měl'ánöspěr'můs) a. [Gk. melas, black; sperma, seed.] Appl. seaweeds with darkcoloured spores.
- melanotic (měl'ánŏt'īk) a. [Gk. melas, black.] Having black pigment unusually developed.
- meliphagous (mělif'ágůs) a. [Gk. meli, honey; phagein, to eat.] Feeding on honey; mellivorous.

melliferous (mělif'ërus) a. [L. mel,

- honey; ferre, to carry.] Honey-producing.
- mellisugent (měl'isū'jent) a. [L. mel, honey; sugere, to suck.] Honeysucking.
- mellivorous (mělĭv'örūs) a. [L. mel, honey; vorare, to devour.] Honeyeating; meliphagous.
- member (mem'ber) n. [I.. membrum, member.] A limb or organ of the body; a well-defined part or organ of a plant.
- membrana (měmbrá'nă) n. [L. membrana, membrane.] A thin film, skin, or layer of tissue covering a part of animal or plant; a thin covering of cells or of unicellular organisms; a membrane.
- membranaceous (měm'brănā'shūs)

 a. [L. membrana, membrane.] Of
 the consistency, or having the
 structure, of a membrane.
- membrane bone,—a bone developing directly in membrane without passing through a cartilage stage.
- membranella (měm'brăněl'ă) n. [L. membrana, membrane.] An undulating membrane formed by fusion of rows of cilia, in some protozoa; ciliated band, in tornaria.
- membraniferous (měm'brănĭf'ërŭs)

 a. [L. membrana, membrane; ferre,
 to carry.] Enveloped in or bearing
 a membrane.
- membranoid (měm'brănoid) a. [L. membrana, membrane; Gk. eidos, form.] Resembling a membrane.
- membranous (mem'branus) a. [L. membrana, membrane.] Resembling or consisting of membrane; pliable and semitransparent.
- membranous cranium,—a mesenchymal investment enclosing brain.
 membranous labyrinth, internal ear, separated from bony cavities by perilymph, and itself containing endolymph.
- membranous vertebral column, continuous sheath of mesoderm enveloping notochord and neural tube.
- membranula (membran'ūlā) n. [L. dim. of membrana, membrana.] A concrescence of cilia, as in certain infusoria.

- membranule (měm'brănūl) n. [I.. dim. of membrana, membrana.] A small opaque space close to body of insect, in anal area of wing of some dragonflies.
- menadione,—vitamin K, present in green vegetables and other foods, essential to formation of prothrombin.
- menarche (měnâr'kē) n. [Gk. men, month; arche, beginning.] First menstruation; age at first menstruation.
- **Mendelian,**—*pert.* character which behaves according to results of Mendel's law, manifesting allelomorphic inheritance.
- Mendelism,—a law or rule governing inheritance of characters in plants and animals, discovered by Gregor Mendel. This principle deals with inheritance of 'unit characters,' presence or absence of one or other of a pair of contrasting characters, dominant and recessive. It also shows that offspring of organisms with a pair of contrasting characters will exhibit these in a definite ratio, and it is extended to deal with groups of characters.
- meningeal (mēnĭn'jëăl) a. [Gk. meningx, membrane.] Pert. or in region of meninges; appl. arteries, veins, nerves, etc.
- meninges (měnin'jēz) n. plu. [Gk. meningx, membrane.] The three membranes enclosing brain and spinal cord, from without inwards: dura mater, arachnoid, and pia mater.
- meningocyte (měnīng'gösīt) n. [Gk. meningx, membrane; kytos, hollow.] A phagocytic cell of the subarachnoid space.
- meningosis (měn'inggō'sis) n. [Gk. meningx, membrane.] Attachment by means of membranes.
- meningospinal (měnǐng'göspi'năl) a. [Gk. meningx, membrane; L. spina, spine.] Pert. spinal cord membranes. meninx,—sing. of meninges.
- meniscus (mēnīs'kūs) n. [Gk. meniskos, small moon.] Interarticular fibro-cartilage found in joints exposed to violent concussion; inter-

- vertebral disc; a tactile disc, being terminal expansion of axis cylinder in tactile corpuscles.
- menopause (mē'nöpôz) n. [Gk. men, month; pausi, ending.] Climacterical cessation of menstruation. Cf. climacteric.
- menotaxis (měn'ötăk'sĭs) n. [Gk. menein, to remain; taxis, arrangement.] Compensatory movements to maintain a given direction of body axis in relation to sensory stimuli; maintenance of visual axis during locomotion.
- mensa (měn'să) n. [L. mensa, table.] Chewing surface of tooth.
- menstruation (měn'strooā'shūn) n.
 [L. mensis, month; struere, to flow.] Periodic discharge from uterus of various vertebrates, chiefly higher mammals.
- mental (měn'tăl) a. [L. mentum, chin.] Pert. or in region of chin; appl. foramen, nerve, spines, tubercle, muscle; appl. scale or plate of fish and of reptile; pert. mentum of insects. [L. mens, mind.] Pert. the mind.
- mentigerous (mentij'erus) a. [L. mentum, chin; gerere, to carry.]
 Supporting or bearing the mentum.
- mentomeckelian (mën'tomëkē'liăn)
 a. [L. mentum, chin; J. F. Meckel, junior, German anatomist]. Appl. a cartilage bone, present in a few lower vertebrates, at either side of mandibular symphysis.
- mentum (měn'tům) n. [L. mentum, chin.] The chin; medial part of gnathochilarium in Diplopoda; region of labium between prementum and submentum in insects; projection between head and foot of some gastropods.
- mericarp (mer'ikârp) n. [Gk. meros, part; karpos, fruit.] A one-seeded indehiscent part of a schizocarp, as of a cremocarp.
- mericlinal (měrikli'năl) a. [Gk. meros, part; klinein, to bend.] Partly periclinal, appl. chimaera with inner tissue of one species only partly surrounded by outer tissue of the other.

- meridional canal,—in ctenophores, a canal into which adradial canals open.
- meridional furrow,—a longitudinal furrow extending from pole to pole of a segmenting egg.
- merisis (měr'isis) n. [Gk. meris, division.] Increase in size owing to cell division; cf. auxesis.

merism,—metamerism, q.v.

- merismatic (mer'ismat'ik) a. [Gk. merismos, partition.] Dividing or separating into cells or segments; meristematic, q.v.
- merismoid (meris/moid) a. [Gk. merismos, partition; eidos, like.] With branched pileus.
- merispore (měr'ĭspōr) n. [Gk. meris, part; sporos, seed.] A segment or spore of a multicellular sporebody.
- meristele (mer'istele) n. [Gk. meris, part; stele, pillar.] A separate part of a monostelic stem passing outwards from stele to leaves; the branch of a stele supplying a leaf.
- meristem (mer'istem) n. [Gk. meristos, divided.] Tissue formed of cells all capable of diversification, as found at growing points; merismatic or meristematic tissue.
- meristematic (měr'istėmăt'ik) a. [Gk. meristos, divided.] Pert. or consisting of meristem; appl. tissue, cells of growing point; merismatic.
- meristematic ring,—tube of meristematic tissue between cortex and pith, subtending the apical meristem and giving rise to vascular tissues.
- meristic (meristic) a. [Gk. meristos, divided.] Segmented; divided off into parts; differing in number of parts.
- meristic variation, changes in number of parts or segments, and in geometrical relations of the parts; cf. substantive variation.
- meristogenetic (měrīs'tojënět'īk) a. [Gk. meristos, divided; genesis, descent.] Developing from meristem.
- merithallus (mě'rĭthăl'ús) n. [Gk. meris, part; thallos, young shoot.]
 A stem unit; an internode.

- Merkel's corpuscle [F. S. Merkel, German anatomist]. A tactile receptor, in skin and in submucosa of mouth.
- mermaid's purse,—horny, floating or fixed, egg-envelope of elasmobranchs.
- mermithaner (měr'mǐthā'ner) n.
 [Gk. mermis, cord; aner, male.]
 Male ant parasitised by Mermis.
- mermithergate (měr'mĭthěr'gāt) n. [Gk. mermis, cord; ergates, worker.] An enlarged worker ant parasitised by Mermis.
- mermithogyne (měrmí'thöjín'ē) n. [Gk. mermis, cord; gyne, female.] Female ant parasitised by Mermis.
- meroandry (měröän'drĭ) n. [Gk. meros, part; aner, male.] The condition of having a reduced number of testes, as a single pair in certain Oligochaeta; cf. holandry.
- meroblast (měr'oblast) n. [Ġk. meros, part; blastos, bud.] Intermediate stage between schizont and merozoite in some Sporozoa; a meroblastic ovum.
- meroblastic (měr'oblās'tīk) a. [Gk. meros, part; blastos, bud.] Appl. ova which undergo only partial segmentation or cleavage in development; developing from part of the oosphere only.
- merocerite (mērŏs'erīt) n. [Gk. meros, thigh; keras, horn.] The fourth segment of crustacean antennae.
- merocrine (měr'okrin) a. [Gk. meros, part; krinein, to separate.] Appl. glands in which secreting cells are able to function repeatedly, as sudoriferous and lactiferous glands. Opp. holocrine.
- merocytes (měr'ösīts) n. plu. [Gk. meros, part; kytos, hollow.] Nuclei formed by repeated division of supernumerary sperm-nuclei, as in egg of selachians, reptiles, and birds; schizonts, q.v.
- merogametes (měr⁷ögămēts⁷) n. plu. [Gk. meros, part; gametes, spouse.] Protozoan individuals specialised for syngamy; microgametes. Opp. hologametes.

merogamy, --microgamy, q.v.

merogastrula (měr'ögăs'troolă) n. [Gk. meros, part; dim. of gaster, stomach.] The gastrula formed from a meroblastic ovum.

merogenesis (měr'öjěn'ësis) n. [Gk. meros, part; genesis, descent.] Formation of parts; segmentation. merogeny,—merogony, q.v.

merognathite (mērŏgnā'thīt) n. [Gk. gnathos, jaw.] meros, thigh; of crustacean Fourth segment

mouth-part.

merogony (měrog'oni) n. [Gk. meros, part; gone, generation.] Development of normal young of small size, from part of an egg, in which there was no female pronucleus.

meroistic (měrōĭs'tĭk) a. [Gk. meros, part; oon, egg.] Appl. ovariole containing nutritive or nurse cells; cf. acrortophic, polytrophic.

merokinesis (měr'ökinē'sis) n. [Gk. meros, part; kinesis, movement.] Formation and division of a threadlike chromosome in the karyomeres.

merome (měrōm) n. [Gk. meros, part]. A body segment; somite, metamere.

meromorphosis (měr'omôr'fōsis) n. [Gk. meros, part; morphosis, shaping.] Regeneration of a part with the new part less than that lost.

meron (mē'rŏn) n. [Gk. meros, upper thigh.] Posterior portion of coxa of insects; sclerite between middle and hind coxae, or immediately above hind coxa, in Diptera; meseusternum.

meronephridia,-micronephridia.

meront (měront') n. [Gk. meros, part; on, being.] A uninucleate schizont-stage in Neosporidia, succeeding the planont-stage.

meroplankton (měr'öplang'kton) n. [Gk. meros, part; plangktos, wandering.] Plankton living only parttime near the surface; temporary plankton, consisting mainly of eggs and larvae; seasonal plankton.

meropodite (mē'röpödīt'), meros (mē'ros) n. [Gk. meros, upper thigh; pous, foot.] Fourth segment of thoracic appendage in crustaceans; femur in spiders.

merosomatous (měr'ösōm'ătŭs) a. [Gk. meros, part; soma, body.] Appl. ascidiozooids divided into two regions, thorax and abdo-

merosome (měr'ösōm) n. [Gk. meros, part; soma, body.] A body segment, somite, or metamere.

merosthenic (mē'rŏsthěn'ĭk) a. [Gk. meros, upper thigh; sthenos, strength.] With unusually developed hind-limbs.

merotomy (měrŏt'ömĭ) n. [Gk. meros, part; temnein, to cut.] Segmentation or division into parts.

merotype (měr'otīp) n. [Gk. meros, part; typos, pattern.] Part of the same perennial plant or vegetatively propagated animal from which a holotype was taken.

merozoite (měr'ozo'īt) n. [Gk. meros, part; zoon, animal.] Divisionproduct of a schizont in Sporozoa.

merozoon (měr'öz \bar{o} 'ŏn) n. meros, part; zoon, animal.] fragment of a unicellular animal containing part of the macronucleus, obtained by artificial division.

merrythought,-furcula of birds, formed by coalesced clavicles.

merus,—meropodite, q.v.

Méry's glands [J. Méry, French anatomist]. Bulbo-urethral glands.

mesad,- mediad, mesiad.

mesadenia (měsădē'niă) n. plu. [Gk middle; aden. mesos. gland.] genital Mesodermal accessory glands in insects; cf. ectadenia.

mesal,—medial, mesial.

mesamoeboid (měs'ămē'boid) a. [Gk. mesos, middle; amoibe, change; eidos, form.] Appl. nucleated cells of blood islands from which blood corpuscles are derived.

mesanepimeron (měsăn'ěpimē'ron, mēz-) n. [Gk. mesos, middle; ana, up; meros, upper thigh.] Sclerite above epimeron and below wing base, in Diptera.

mesarch (měs'ark) a. [Gk. mesos, middle; arche, beginning.] Appl.

xylem having metaxylem developing in all directions from the protoxylem, characteristic of ferns; having the protoxylem surrounded by metaxylem; beginning in a mesic environment, appl. seres.

mesaticephalic (měs'ătikěfăl'īk,-sěf-) a. [Gk. mesatos, mid; kephale, head.] Having a cephalic index of 75 to 80; mesocephalic.

mesaxonic (měs'akson'ik) a. [Gk. mesos, middle; axon, axis.] With the line dividing the foot, passing

up the middle digit.

mesectoderm (měsěk'toděrm) [Gk. mesos, middle; ektos, outside; derma, skin.] Parenchyma formed of descendants of ectodermal cells which migrated inwards.

mesembryo (měsěm'briö) n. |Gk. mesos, middle; embryon, embryo.]

The blastula.

mesencephalon (měs'ěnkěf'alon, -sef-) n. [Gk. mesos, middle; en. in; kephale, head.] The mid-brain, comprising corpora quadrigemina (bigemina), cerebral peduncles, and aqueduct of Sylvius.

mesenchyma (měsěng'kimă), mesenchyme (měsěng'kim) n. [Gk. mesos, middle; engchein, to pour in.] A mass of tissue, intermediate between ectoderm and endoderm of a gastrula.

mesendoderm (měsěn'doděrm) n. [Gk. mesos, middle; endon, within; derma, skin.] Cells lying posteriorly to lip of blastopore, partly invaginated with endoderm in gastrulation, in development of some molluscs.

mesenterial (měs'entē'rĭăl, měz-) a. [Gk. mesos, middle; enteron, gut.] Pert. a mesentery; appl. filaments of Actinozoa.

mesenteric (měs'entěr'ik) a. [Gk. mesos, middle; enteron, gut.] Pert. a mesentery; appl. arteries, glands, nerves, veins, etc.

mesenteriole (měs'ëntē'riol) n. [L. dim. of mesenterium, mesentery.] A fold of peritoneum derived from mesentery, and retaining vermiform process or appendix in position.

mesenterium,-mesentery, q.v.

mesenteron (měsěn'tëron, mēz-) n. [Gk. mesos, middle; enteron, gut.] The main digestive cavity of Actinozoa and other Coelentera; portion of alimentary canal lined by endoderm; mid-gut.

mesentery (měs'enteri, měz-) n. [L. mesenterium, mesentery.] A peritoneal fold serving to hold viscera in position; a muscular partition extending inwards from body-wall in coelenterates.

mesepimeron (měs'epimē'rŏn, měz-) n. [Gk. mesos, middle; epi, upon; meros, upper thigh.] The epimeron of insect mesothorax; meskatepimeron in Diptera.

mesepisternum (měs'epister'nům) n. [Gk. mesos, middle; epi, upon; sternon, breast.] Meso-episternum, sclerite below anterior spiracle in

Diptera; mesanepisternum.

mesethmoid (měsěth'moid, mēz-) a. Gk. mesos, middle: ethmos, sieve: eidos, form.] Between the two ectethmoid bones; appl. ethmoid plate of cranium when it ossifies; median cranial bone of vertebrates.

mesiad (měz'ĭăd) adv. [Gk. mesos, middle; L. ad, to.] Towards or near

the middle plane.

mesial (mē'zĭāl), mesian (mē'zĭăn) a. [Gk. mesos, middle.] middle vertical or longitudinal plane.

mesic (měs'ik, mē'zik) a. [Gk. mesos, middle.] Conditioned by temperate moist climate, neither xeric nor hydric.

mesoarion,—mesovarium, q.v.

mesobenthos (mčs'oběn'thos, mēz-) n. [Gk. mesos, middle; benthos, depths.] Animal and plant life of sea-bottom when depth is between 100 and 500 fathoms.

mesoblast (měs'öblast, mēz-) n. [Gk. mesos, middle; blastos, bud.] The mesoderm or middle layer of an

embryo; mesoblastema

mesoblastic (měs'öblas'tik, mēz-) a. [Gk. mesos, middle; blastos, bud.] Pert. or developing from middle laver of an embryo.

- mesobranchial (měs'obrăng'kiăl, mēz-) a. [Gk. mesos. middle; brangchia, gills.] Pert. middle gillregion, as in Crustacea.
- mesobronchus (měs'öbröng'kůs, mēz-) n. [Gk. mesos, middle; brongchos. windpipe.] In birds, the main trunk of a bronchus giving rise to secondary bronchi.

mesocaecum (mesosecum mesocaecum) n. [Gk. mesos, middle; L. caecus, blind.] The mesentery connected with the caecum.

mesocardium (měs'okâr'dĭum, mēz-)
n. [Gk. mesos, middle; kardia,
heart.] An embryonic mesentery
binding heart to pericardial walls;
part of pericardium enclosing veins
(venous m.) or aorta (arterial m.);
mesocardial ligament or gubernaculum cordis.

mesocarp (měs'ökârp, mēz-) n. [Gk. mesos, middle; karpos, fruit.] The middle layer of the pericarp.

mesocentrous (měs'ösěn'trŭs,mēz-)a. [Gk. mesos, middle; kentron, centre.] Ossifying from a median centre.

mesocephalic (měs'ökéf'ālīk, mē'zösēf'ālīk) a. [Gk. mesos, middle; kephale, head.] Having a cranial capacity of between 1350 and 1450 c.c.; cf. megacephalic, microcephalic.

mesochilium (měs'ökī'lĭūm, mēz-) n.
[Gk. mesos, middle; cheilos, lip.]
The middle portion of labellum of orchids.

mesochiton (mčs'ókī'tŏn, mēz-) n. [Gk. mesos, middle; chiton, coat.] Middle layer of oogonial wall, between endochiton and exochiton, as in Fucales; mesochite.

mesocoel (měs'ösēl, mēz-) n. [Gk. mesos, middle; koilos, hollow.] Middle portion of coelomic cavity; the second of three main parts of coelom of molluscs; cavity of mesencephalon, aqueduct of Sylvius or iter.

mesocolic (měs'ökŏl'īk, mēz-) a. [Gk. mesos, middle; kolon, large intestine.] Pert. mesocolon; appl. lymph glands.

mesocolon (měs'ökō'lon, mēz-) n.

[Gk. mesos, middle; kolon, large intestine.] A mesentery or fold of peritoneum attaching colon to dorsal wall of abdomen.

mesocoracoid (měs'okŏr'ăkoid, mēz-)
a. [Gk. mesos, middle; korax,
crow; eidos, form.] Situated between hyper- and hypo-coracoid;
appl. middle part of coracoid arch
of certain fishes.

mesocotyl (měs'ökŏt'īl, mēz-) n. [Gk. mesos, middle; kotyle, cup.] Part of axis between scutellum and coleoptile.

mesocycle (měs'osīkl, mēz-) n. [Gk. mesos, middle; kyklos, circle.] A layer of tissue between xylem and phloem of a monostelic stem; part of conjunctive tissue of stele.

mesodaeum (mēs'ödē'ŭm, mēz-) n. [Gk. mesos, middle; odaios, pert. way.] Endodermal part of embryonic digestive tract, between stomodaeum and proctodaeum.

mesoderm (měs'öděrm, měz-) n. [Gk. mesos, middle; derma, skin.] The mesoblast or embryonic layer lying between ectoderm and endoderm.

mesodermal (měs'öděr'măl, mēz-) a.
[Gk. mesos, middle; derma, skin.]
Pert., derived, or developing from mesoderm.

mesodesm (měs'ödězm, mēz-) n. [Gk. mesos, middle; desma, bond.] Part of mesocycle.

mesodont (mēs'ödŏnt, mēz-) a. [Gk. mesos, middle; odous, tooth.] Appl. stag-beetles having a medium development of mandible projections.

mesogaster (měs'ögás'tër, mēz-) n. [Gk. mesos, middle; gaster, stomach.] The mesentery or fold of peritoneum supporting the stomach.

mesogastric (měs'ögăs'trĭk, mēz-) a. [Gk. mesos, middle; gaster, stomach.] Pert. a mesogaster or mesogastrium, or to middle gastric region.

mesogastrium (měs'ögăs'trīŭm, mēz-) n. [Gk. mesos, middle; gaster, stomach.] Mesentery connecting stomach with dorsal abdominal wall in embryo; middle abdominal region. mesoglia (měsöglī'ā, mēz-) n. [Gk. mesos, middle; gloia, glue.] Mesodermal phagocytic interstitial cells of nervous system; of Hortega: microglia; of Robertson: oligodendroglia.

mesogloea (měs'öglē'ă, mēz-) n. [Gk. mesos, middle; gloia, glue.] An intermediate non-cellular gelatinous layer in sponges and coelenterates

mesognathion (měs'onath'ion, mēz-) n. [Gk. mesos, middle; gnathos, The lateral segment of premaxilla, bearing lateral incisor.

mesohepar (měs'ohē'păr, mēz-) n. [Gk. mesos, middle; hepar, liver.] Mesentery supporting liver.

mesohydrophytic (měs'öhī'drofitik, mēz-) a. [Gk. mesos, middle; water; phyton, plant.] hydor, Growing in temperate regions but requiring much moisture.

mesolecithal (mes'oles'ithal, mez-) a. [Gk. mesos, middle; lekithos, yolk.] Having a moderate yolk content;

cf. centrolecithal.

mesology (měsŏl'öjĭ, mēz-) n. [Gk. mesos, middle; logos, discourse.] The study of relations between organism and environment; bionomics.

mesome (měs'ōm, mēz'ōm) n. [Gk. mesos, middle.] The axis regarded as a morphological unit of plants.

mesomere (měs'omēr, mēz-) n. [Gk. middle; meros, part.] mesos, Middle zone of coelomic pouches in embryo; mesoblastic somite or protovertebra; medial branch of phallic lobe in insects.

mesometrium (měs'ömē'trium, mēz-) [Gk. mesos, middle; metra, The mesentery of uterus

and connecting tubes.

mesomitosis (mes'omito'sis, mez-) n. [Gk. mesos, middle; mitos, thread.] Mitosis within nuclear membrane, without co-operation of cytoplasmic elements; cf. metamitosis.

mesomorphic (měs'omôr'fik, mēz-) [Gk. mesos, middle; morphe, form.] Having form, structure, or size normal or intermediate between extremes; mesoplastic.

mesomyodian (měs'omīo'dian, mēz-) [Gk. mesos, middle; mys, muscle; eidos, form.] Appl. birds with muscles of syrinx attached to middle of bronchial semi-rings.

meson (měs'ŏn, mē'zŏn) n. [Gk. mesos, middle.] The central plane,

or region of it.

mesonephric (měs'oněf'rik, mēz-) a. [Gk. mesos, middle; nephros, kidney.] *Pert.* mesonephros, or midkidney; appl. duct, tubules.

mesonephridium (měs'oněfrid'iŭm, [Gk. mesos, middle; mēz-) n. nephros, kidney; idion, dim.] A nephridium or excretory organ of certain invertebrates, derived from mesoblast.

mesonephros (měs'öněf'rŏs, mēz) n. [Gk. mesos, middle; nephros, kidney.] Intermediate part of excretory organ vertebrate in embryos; Wolffian body.

mesonotum (měs'önō'tům, mēz-) n. [Gk. mesos, middle; noton, back.] Dorsal part of insect mesothorax.

mesoparapteron (měs'oparap'teron, mēz-) n. [(ik. mesos, middle; para, beside; pteron, wing.] A small sclerite of mesothorax of some insects.

mesopetalum (měs'öpět'ălum, mēz-) n. | Gk. mesos, middle; petalon, leaf.] Labellum or lip of an orchid.

mesophanerophyte (měs'öfăn'ëröfīt, mēz-) n. [Gk. mesos, middle; phaneros, manifest; phyton, plant.] Tree from 8 to 30 metres in height.

mesophil (měs'öfil, měz-) a. [Gk. mesos, middle; philein, to love.] Thriving at moderate temperatures, at between 20° and 40° C. when appl. bacteria; mesophilic; mesic, q.v. n. Mesophile.

mesophloem (měs'öflő'ěm, mēz-) n. mesos, middle; phloios, [Gk. smooth bark.] Middle or green

bark; mesophloeum.

mesophragma (měs'öfrăg'mă, mēz-) n. [Gk. mesos, middle; phragma, fence.] A chitinous piece descending into interior of insect body with post-scutellum for base; M or Hensen's line, q.v.

- mesophryon (měsŏf'rĭŏn, mēz-) n. [Gk. mesos, middle; ophrys, eyebrow.] The elevated median headregion of trilobites.
- mesophyll (měs'öfil, mēz-) n. [Gk. mesos, middle; phyllon, leaf.] The internal parenchyma of a leaf.
- mesophyte (měs'öfīt, mēz-) n. [Gk. mesos, middle; phyton, plant.] A plant thriving in temperate climate with normal amount of moisture.
- mesoplankton (měs'öplăng'ktŏn, mēz-) n. [Gk. mesos, middle; plangktos, wandering.] Drifting animal and plant life from a hundred fathoms downwards; drifting organisms of medium size; cf. megaloplankton, microplankton, nanoplankton.
- mesoplast (měs'öplast, mēz-) n. [Gk. mesos, middle; plastos, moulded.]
 A cell nucleus.
- mesoplastic (měs'oplăs'tĭk, mēz-) a. [Gk. mesos, middle; plastos, moulded.] Having a normal or average form; mesomorphic; appl. constitutional type.
- mesoplastron (měs'öplăs'trŏn, mēz-)
 n. [Gk. mesos, middle; F. plastron,
 breast-plate.] Plate between hyoand hypo-plastron of certain
 turtles.
- mesopleurite (měs'öploo'rīt, mēz-) n. [Gk. mesos, middle; pleura, side.] Lateral mesothoracic sclerite. as in Diptera.
- mesopleuron (měs'öploo'rŏn, mēz-)

 n. [Gk. mesos, middle; pleura,
 side.] A lateral part of insect
 mesothorax; a mesopleurite
- mesopodial (měs'öpō'dĭăl, mēz-) a. [Gk. mesos, middle; pous, foot.] Pert. a mesopodium.
- mesopodium (měs'öpō'diŭm, mēz-)
 n. [Gk. mesos, middle; pous, foot.]
 Leaf-stalk or petiole region of leaf;
 middle part of molluscan foot; the
 metacarpus or metatarsus.
- mesopraescutum (měs'öprēskū'tǔm, mēz-) n. [Gk. mesos, middle; L. prae, before; scutum, shield.] Praescutum of mesothorax in insects.

- mesopterygium (měs'ötčrij'ĭŭm, mēz-) n. [Gk. mesos, middle, pterygion, little wing or fin.] The middle of three basal pectoral fin-cartilages in recent elasmobranchs.
- mesopterygoid (měs'ötěr'īgoid, mēz-) n. [Gk. mesos, middle; pteryx, wing; eidos, form.] The middle of three pterygoid bone elements of teleosts; the ectopterygoid.
- mesoptile (měs'ötīl, mēz-, -ptīl) n. [Gk. mesos, middle; ptilon, feather.] Prepenna following protoptile and succeeded by metaptile or by teleoptile.
- mesorchium (měsôr'kiŭm, mēz-) n. [Gk. mesos, middle; orchis, testicle.] Mesentery supporting testis.
- mesorectum (měs'örěk'tům, mēz-) n. [Gk. mesos, middle; I.. rectus, straight.] Mesentery supporting rectum.
- mesorhinal (měs'orī'nal, mēz-) a. [Gk. mesos, middle; rhis, nose.] Between nostrils.
- mesorhinium (měs'or!n'iŭm, mēz-) n.
 [Gk. mesos, middle; rhis, nose.]
 The internarial surface region of a bird's bill.
- mesosalpinx (měs'osăl'pĭngks, mēz-)
 n. [Gk. mesos, middle; salpingx,
 trumpet.] The portion of broad
 ligament enclosing uterine tube.
- mesoscapula (měs'öskăp'ūlă, mēz-) n. [Gk. mesos, middle; L. scapula, shoulder-blade.] Scapular spinc.
- mesoscutellum (měs'öskūtěl'ům, mēz-) n. [Gk. mesos, middle; L. scutellum, small shield.] Scutellum of insect mesothorax.
- mesoscutum (měs'öskū'tǔm, mēz-) n. [Gk. mesos, middle; L. scutum, shield.] Scutum of insect mesothorax.
- mesosoma (měs'ösō'mă, mēz-) n. [Gk. mesos, middle; soma, body.] Middle part of body; praeabdomen of Arthropoda.
- mesosome (měs'ösôm, měz-) n. [Gk. mesos, middle; soma, body.] A phallosome, q.v.

- mesosperm (měs'öspěrin, měz-) n. [Gk. mesos, middle; sperma, seed.] Integument investing nucellus of ovule.
- mesospore (měs'öspōr, mēz-), mesosporium (měs'öspō'rīŭm, mēz-) n. [Gk. mesos, middle; sporos, seed.] The intermediate of three spore coats; a unicellular teleutospore in certain rust fungi; a resting-spore stage.
- mesostate (měs'östāt, mēz-) n. [Gk. mesos, middle; stasis, standing.]
 Intermediate stage in metabolism.
- mesostereom (mēs'östēr'eŏm, mēz-) n. [Gk. mesos, middle; stereos, solid.] The middle layer of thecal plates of Cystidea.
- mesosternebra (měs'óstěr'nëbră, mēz-) n. [Gk. mesos, middle; sternon, breast-bone.] A part of developing mesosternum.
- mesosternum (měs'ostěr'nům, měz-)
 n. [Gk. mesos, middle; L. sternum.
 breast-bone.] Middle part of
 sternum of vertebrates; gladiolus;
 sternum of mesothorax of insects;
 mesosternal area, episternum of
 mesothorax, or meskatepisternum
 of Diptera.
- mesostethium (měs'östē'thiŭm, mēz-) n. [Gk. mesos, middle; stethos, chest.] A mesosternum.
- mesostylous (měs'óstī'lūs, mēz-) a. [Gk. mesos, middle; stylos, pıllar.] Having styles of intermediate length; appl. heterostylous flowers.
- mesotarsal (měs'ötâr'săl, mēz-) a. [Gk. mesos, middle; L. tarsus, ankle-joint.] Pert. mesotarsus.
- mesotarsus (mēs'otâr'sŭs, mēz-) n. [Gk. mesos, middle; L. tarsus, ankle-joint.] A middle-limb tarsus of insects.
- mesotergum (měsötěr'gům, měz-) n. [Gk. mesos, middle; L. tergum, back.] Median arched portion or axis of trilobite body.
- mesothecium (měs'öthē'siŭm, mēz-)
 n. [Gk. mesos, middle; theke, cup.]
 The middle investing layer of an anther-sac.
- mesothelium (měs'öthē'lĭŭm, mēz-)

- n. [Gk. mesos, middle; thele, nipple.] Mesoderm bounding primitive coelom and giving rise to muscular and connective tissue.
- mesotherm (měs'öthěrm, mēz-) n.
 [Gk. mesos, middle; therme, heat.]
 Plant thriving in moderate heat.
- mesothoracic (měs'öthōrăs'îk, mēz-)
 a. [Gk. mesos, middle; thorax, chest.] Pert. or in region of mesothorax; appl. a spiracle, of insects.
- mesothorax (měs'öthō'răks, mēz-) n. [Gk. mesos, middle; thorax, chest.] The middle segment of thoracic region of insects.
- mesotic (měső'tik, mēz-) a. [Gk. mesos, middle; ous, ear.] Appl. paired chondrocranial cartilages in birds, between parachordal and acrochordal; also basiotic.
- mesotriaene (měs'otrī'ēn, mēz-) n. [Gk. mesos, muddle; triaina, trident.] Aberrant type of triaene spicule.
- mesotrochal (měsŏť rokal, mēz-) a. [Gk. mesos, middle; trochos, wheel.] Appl. an annulate larva with circlet of cilia round middle of body.
- mesotrophic (měs'otrŏf'îk, mēz-) a. [Gk. mesos, middle; trophe, nour-ishment.] Mixotrophic, q.v.; providing a moderate amount of nutrition, appl. environment.
- mesotympanic (mesotympan'ik, mez-) n. [Gk. mesos, middle; tympanon, drum.] Symplectic; a bone in suspensory apparatus of lower jaw in fishes.
- mesovarium (měs'ōvā'rīŭm, mēz-) n. [Gk. mesos, middle; L. ovarium, ovary.] Mesentery of ovary; suspensory mesentery in fishes.
- mesoventral (měs'övěn'trāl, mēz-) a. [Gk. mesos, middle; L. venter, belly.] In middle ventral region.
- Mesozoic (měs'özō'îk, mēz-) a. [Gk. mesos, middle; zoe, life.] Appl. or pert. secondary geological era, the age of reptiles.
- mestome (mes'tom) n. [Gk. mestos, filled.] A vascular bundle, including hadrome and leptome.

- metabiosis (mět'ábīō'sīs) n. [Gk. meta, after; biosis, a living.] Condition in which one organism lives only after another has prepared its environment and has died.
- metabolic (mět'ábŏl'ík) a. [Gk. metabole, change.] Changeable; appl. chemical changes occurring in the living organism; influencing metabolism, appl. hormones; formed during metabolism; metamorphosing.

metabolin,—metabolite, q.v.

- metabolism (mětăb'ölīzm) n. [Gk. metabole, change.] The chemical change, constructive and destructive, occurring in living organisms.
- metabolite (mětăb'ölīt) n. [Gk. metabole, change.] Any product of metabolism.
- metaboly (mětăb'ölĭ) n. [Gk. metabole, change.] Change, particularly of shape, as in Eugleninae.
- metabranchial (mět'ábrăng'kĭăl) a. [Gk. meta, after; brangchia, gills.] Pert. or in region of posterior gill region.
- metacarpal (mět'ăkâr'păl) a. [Gk. meta, after; karpos, wrist.] Pert. metacarpus; appl. bones, articulations, etc. n. A primary wing-quill in the metacarpal region.
- metacarpophalangeal (mět'ākâr'pöfălăn'jēăl) a. [Gk. meta, after; karpos, wrist; phalangx, series.] Appl. articulations between metacarpals and phalanges.
- metacarpus (mět'ākār'pŭs) n. [Gk. meta, after; karpos, wrist.] The skeletal part of hand between wrist and fingers, consisting typically of five cylindrical bones.
- metacentric (mět'ásěn'trĭk) a. [Gk. meta, among; kentron, centre.] Having the centromere at or near the middle, appl. chromosomes; cf. acrocentric, telocentric. n. A metacentric or V-shaped chromosome; isochromosome.
- metacercaria (mět'ăsërkā'rīā) n. [Gk. meta, after; kerkos, tail.] A cercaria after encystment; adolescaria.

- metacestode (měť ásěs 'tōd) n. [Gk. meta, after; kestos, girdle; eidos, form.] Bladder-worm, encysted stage of a cestode; plerocestoid; plerocercoid.
- metachromas-y,-ie (mět'ákrō'măsĭ)

 n. [Gk. meta, change of; chroma, colour.] Condition of certain tissues and cell components which, treated with basic aniline stains, show other than the fundamental colour constituent.
- metachromatic (mět'ákrōmăt'îk) a. [Gk. meta., change of; chroma, colour.] Appl. substances characterised by metachromasy; appl. granules of reserve food substances which stain with basic dyes, in bacteria and algal cells.
- metachromatinic grains,—chromatoid bodies found in cells, very similar to chromatin in properties and characteristics.
- metachromy (mět'ákrō'mĭ) n. [Gk. meta, change of; chroma, colour.]
 Change in colour, as of flowers.
- metachrosis (měťákrō'sĭs) n. [Gk. meta, change of; chrosis, colouring.] Ability to change skin colour by expansion or contraction of pigment cells.
- metacneme (mět'áknēmē) n. [Gk. meta, after; kneme, tibia.] A secondary mesentery of Zoantharia.
- metacoel (mět'ásēl) n. [Gk. meta, after; koilos, hollow.] The posterior part of coelom of molluscs; anterior extension of fourth ventricle of brain.
- metacone (mět'ákōn) n. [Gk. meta, after; konos, cone.] Postero-external cusp of upper molar.
- metaconid (met'ākō'nid) n. [Gk. meta, after; konos, cone.] Posterointernal cusp of lower molar.
- metaconule (měťákô'nūl) n. [Gk. meta, after; konos, cone.] Posterior secondary cusp of upper molar.
- metacoracoid (měťákör'ákoid) n. [Gk. meta, after; korax, crow; eidos, form.] Posterior part of coracoid.

- metacromion (mět'ákrō'miŏn) n. [Gk. meta, after; akros, summit; omos, shoulder.] Posterior branch-process of acromion-process of scapular spine.
- metacyclic (mět'ásšk'lìk) a. [Gk. meta, after; kyklos, circle.] Appl. final infective forms, of certain parasitic protozoa, which pass on to next host.
- metadiscoidal (mět'ádiskoid'ál) a. [Gk. meta, after; diskos, disc; eidos, form.] Appl. placenta in which villi are at first scattered and later restricted to a disc, as in man and monkeys.
- metadromous (mětăd'römŭs) a. [Gk. meta, after; dromos, running.] With primary veins of segment arising from upper side of midrib
- meta-episternum, metepisternum, q.v.
- metaesthetism (mětēsthē'tizm) n. [Gk. meta, after; aisthetos, perceptible by senses.] Doctrine that "consciousness is a product of evolution of matter and force."
- metagastric (mět'āgăs'trĭk) a. [Gk. meta, after; gaster, stomach.] Pert. posterior gastric region.
- metagastrula (měťágăs'troolă) n. [Gk. meta, after; gaster, stomach.] A modified form of gastrula.
- metagenesis (mět'ájěn'esis) n. [Gk. meta, after; genesis, descent.] Alternation of sexual and asexual generations; cf. heterogenesis.
- metagyny (mëtaj'îni, mět'agi'ni) n. [Gk. meta, afterwards; gyne, female.] Protandry.
- metakinesis (mět'ákíně'sis) n. [Gk. meta, after; kinein, to move.] Middle stage of mitosis, during which chromosomes are grouped in equatorial plate; movement of chromosomes between prophase and metaphase.
- metallic (mětăl'îk) a. [Gk. metallon, mine.] Iridescent; appl. colours due to interference by fine striae or thin lamellae, as in insects.
- metaloph (měť alof) n. [Gk. meta, after; lophos, crest.] The posterior

- crest of a molar, uniting metacone, metaconule, and hypocone.
- metamere (mět'āmēr) n. [(k. meta, after; meros, part.] A body segment.
- metameric (mět'ăměr'îk) a. [Gk. meta, after; meros, part.] Pert. metamerism or segmentation.
- metamerised (mět amėrīzd) a. [Gk. meta, after; meros, part.] Segmented.
- metamerism (mět'āmërīzm) n. [Gk. meta, after; meros, part.] The condition of a body divided up into segments more or less alike; segmentation; zonal symmetry.
- metamitosis (mět'āmītō'sīs) n. [Gk. meta, after; mitos, thread.] Mitosis in which cytoplasmic and nuclear elements are both affected; cf. mesomitosis.
- metamorphosis (mět'ámôr'fōsīs) n. [Gk. meta. change of; morphe, form.] Change of form and structure undergone by an animal from embryo to adult stage, as in insects; transformation of one structure into another, as of stamens into petals; interference with normal symmetry in flowers; internal chemical change.
- metamps (mět'amps) n. plu. [Gk. meta, change of; morphe, form.] Different forms of same species, as in certain sponges.
- metamyelocyte (mětămí'élösīt) n. [Gk. neta, b-yond; myelos, marrow; kytos, hollow.] A myelocyte with horseshoe-shaped nucleus before transformation into a leukocyte.
- metanauplius (mět'ănôp'liŭs) n. [Gk. meta, after; L. nauplius, kind of shell-fish.] Larval stage of Crustacea, succeeding nauplius stage.
- metandry (mětăn dri) n. [Gk. meta, after; aner, male.] Meroandry with retention of posterior pair of testes only, opp. proandry; protogyny, opp. protandry.
- metanephric (měť aněť rík) a. [Gk. meta, after; nephros, kidney.] Pert. or in region of hind-kidney.

- metanephros (měť aněť roš) n. [Gk. meta, after; nephros, kidney.] The organ arising behind mesonephros and replacing it as functional kidney of fully-developed Amniota.
- metanotum (měť ano tum) n. [Gk. meta, after; noton, back.] Notum or tergum of insect metathorax.
- metanucleus (mět'ănū'klëŭs) n. [Gk. meta, after; L. nucleus, kernel.] Egg-nucleolus after extrusion from germinal vesicle.
- metapeptone (mět'ăpěp'tōn) n. [Gk. meta, after; peptos, digested.] A product of action of gastric juice on albumins.
- metaphase (mět'áfāz) n. [Gk. meta, after; phainein, to appear.] The stage in mitosis or meiosis in which chromosomes are split up in equatorial plate.
- metaphery (mětăf'ërĭ) n. [Gk. meta, beyond; pherein, to bear.] Displacement of organs.
- metaphloem (mět'āflō'ëm) n. [Gk. meta, after; phlotos, inner bark.]
 The phloem of secondary xylem.
- metaphragma (měť áfrág má) n. [Gk. meta, after; phragma, fence.] An internal metathoracic septum in insects.
- metaphysis (mětăf'isĭs) n. [Gk. meta, besides; physis, growth.] Paraphysis, q.v., of fungi.
- metaphyte (měť afit) n. [Gk. meta, after; phyton, plant.] A multi-cellular plant, opp. protophyte.
- metaplasia (mět'āplā'sĭā) n. [Gk. meta, change of; plasis, moulding.] Conversion of tissue from one form to another, as in ossification.
- metaplasis (mětăp'lăsīs) n. [Gk. meta, after; plasis, moulding.]
 The mature period in life of an individual.
- metaplasm (měťáplázm) n. [Gk. meta, after; plasma, mould.] Lifeless or ergastic ingredients of protoplasm, opp. to living material or organoids.
- metaplastic (mět'ăplăs'tĭk) a. [Gk. meta, after; plastos, moulded.] Pert. metaplasia; pert. metaplasm.

metaplastic or metaplasmic bodies,
—grains of protoplasm which are
stages or products of metabolism
and not true protoplasm.

metapleural (mčťáploo'rāl) a. [Gk. meta, after; pleura, side.] Posteriorly and laterally situated; pert. metapleure; pert. metapleuron.

- metapleure (měťáploor) n. [Gk. meta, after; pleura, side.] An abdominal or ventro-lateral fold of integument of certain primitive Chordata.
- metapleuron (měťáploor'ŏn) n. [Gk. meta, after; pleura, side.] The pleuron of insect metathorax.
- metapneustic (mět'ănū'stīk, -pnū'-)a. [Gk. meta, after; pneuma, breath.] Appl. insect larvae with only the terminal pair of spiracles.
- metapodeon (mět'ăpŏd'ëŏn), metapodeum (mět'ăpō'dëŭm) n. [Gk. meta, after; pous, foot.] That part of insect abdomen behind petiole or podeon.
- metapodial (mět'ápō'diăl) a. [Gk. meta, after; pous, foot.] Pert. a metapodeon or metapodium.
- metapodium (měťápo'diŭm) n. [Gk. meta, after; pous, foot.] Posterior portion of molluscan foot; portion of foot between tarsus and digits; in four-footed animals, metacarpus and metatarsus.
- metapodosoma (měť apodoso ma) n. [Gk. meta, after; pous, foot; soma, body.] Body region bearing third and fourth pair of legs in Acarina.
- metapolar cells,—second circlet of cells of polar cap of rhombogen of Rhombozoa.
- metapophysis (měť apof í isis) n. [Gk. meta, after; apo, from; phyein, to grow.] A prolongation of a vertebral articular process developed in certain vertebrates; mammillary process.
- metapore (mět'ápōr) n. [Gk. meta, after; poros, channel.] The medial aperture in roof of fourth ventricle of brain; Magendie's fora-

- metapostscutellum (měťápōst'skūtěl'ŭm) n. [Gk. meta, after; L. post, after; scutellum, small shield.] Postscutellum of insect metathorax.
- metapraescutum (mět'ăprēskū'tūm)
 n. [Gk. meta, after; L. prae, before;
 scutum, shield.] Praescutum of
 insect metathorax.
- metapterygium (měťátërij'íum) n. [Gk. meta, after; pterygion, little wing.] The posterior basal fincartilage, pectoral or pelvic, of recent elasmobranchs.
- metapterygoid (mět'átěr'ígoid) n. [Gk. meta, after; pteryx, wing; eidos, form.] Posterior of three pterygoid elements in certain lower vertebrates.
- metaptile (mět'átīl, -ptīl) n. [Gk. meta, after; ptilon, feather.] A plumose penna or feather; cf. mesoptile, teleoptile.
- metarachis (mět'ārā'kĭs) n. [Gk. meta, after; rhachts, spine.] Face of Pennatulacea which coincides with sulcar aspect of terminal zooid—so-called dorsal surface.
- metascutellum (mět'áskūtěl'ům) n. [Gk. meta, after; L. scutellum, small shield.] Scutellum of insect metathorax.
- metascutum (měť ásků tům) n. [Gk. meta, after; L. scutum, shield.] Scutum of insect metathorax.
- metasitism (mět'ăsītizm) n. [Gk. meta, after; sitos, food.] A cannibalistic mode of life.
- metasoma (mēt'āsō'mā) n. [Gk. meta, after; soma, body.] The six terminal segments of opisthosoma of Eurypterida; posterior body-region of Arachnoidea; postabdomen; abdomen, as of woodlice.
- metasomatic (měť asomať ik) a. [Gk. meta, after; soma, body.]

 Pert. or situated in metasoma.
- metasperm (mět'ăspěrm) n. [Gk. meta, after; sperma, seed.] A plant having seeds in a closed ovary; an angiosperm.
- metasporangium (mět'ăspörăn'jium)
 n. [Gk. meta, after; sporos, seed;
 anggeion, vessel.] A sporangium

- containing resting spores, as in Bacillaceae.
- metastasis (mětăs'tăsīs) n. [Gk. metastasis, removal.] Metabolism; transference of function from one organ to another; transport of bacteria by the circulatory system.
- metastatic life history,—that of certain Trematoda in which the young form, after entering intermediate host, metamorphoses into adult, after which intermediate host is swallowed by final host.
- metasternum (mět'ástěr'nům) n. [Gk. meta, after; L. sternum, breastbone.] The sternum of insect metathorax; posterior sternal part, or xiphisternum, of Anura; xiphoid or ensiform process, posterior part of sternum of higher vertebrates.
- metasthenic (měť asthěn'ík) a. [Gk. meta, after; sthenos, strength.] With well-developed posterior part of body.
- metastigmate (mět'ástig'māt) a. [Gk. meta, after; stigma, mark.] Having posterior tracheal openings or stigmata, as in mites.
- metastoma (měťástö'mā) n. [Gk. meta, after; stoma, mouth.] The two-lobed lower lip of Crustacea; 'hypopharynx' of Myriopoda; median plate belind mouth in Palaeostraca; metastome.
- metastructure (měťástrůk'tůr) n. [Gk. meta, after; L. struere, to build.] Ultramicroscopic organisation.
- metasyndesis (mět'ásĭn'desĭs) n. [Gk. meta, after; syndesis, bond.] Telosyndesis, q.v.
- metatarsal (mět'ātâr'săl) a. [Gk. meta, after; L. tarsus, ankle.] In region of metatarsus; appl. arteries, veins, etc.; pert. metatarsal bones.
- metatarsophalangeal (měťátár'söfălăn'jeăl) a. [Gk. meta, after; L. tarsus, ankle; Gk. phalangx, troop.] Appl. articulations between metatarsus and phalanges of foot.
- metatarsus (mět'ātâr'sūs) n. [Gk. meta, after; L. tarsus, ankle.] Part of foot between tarsus and toes; first joint of tarsus in insects; first dactylopodite or basitarsus in spiders.

- metathalamus (mět'áthăl'ámůs) n. [Gk. meta, after; thalamos, chamber.] The geniculate bodies of the thalamencephalon.
- metatherian (mět'áthē'riăn) a. [Gk. meta, beyond; therion, small animal.] Appl. marsupials, with short-term placenta and later development in marsupium.

metathetely (měť áthěť eli) n. [Gk. metathein, to run behind; telos, completion.] Hysterotely.

- metathorax (měťáthō′rāks) n. [Gk. meta, after; thorax, chest.]

 Posterior segment of insect thorax.
- metatracheal (měťátra'kēål) a. [Gk. meta, between; L.L. trachia, windpipe.] Appl. wood, with xylem parenchyma located independently of the vessels.
- metatroch (mět'átrŏk) n. [Gk. meta, after; trochos, wheel.] In a trochophore, a circular band of cilia behind the mouth.
- metatrophic (měťátrof'ík) a. [Gk. meta, change of; trophe, nourishment.] Living on both nitrogenous and carbonaceous organic matter.
- metatympanic,—entotympanic, q.v. metatype (mět'átīp) n. [Gk. meta, after; typos, image.] A topotype of the same species as the holotype or lectotype.
- metaxenia (mět'ăzē'nīă) n [Gk. meta, after; I.. xenia, gıfts.]
 Physiological effect of pollen upon maternal tissue.
- metaxylem (mět'āzī'lěm) n. [Gk. meta, after; xylon, wood.] Secondary xylem with many thick-walled cells.
- metazoaea (mět'ázōē'á) n. [Gk. meta, after; zoe, life.] A larval stage of Crustacea between zoaea and megalopa stages.
- metazonite (měťázōnīt) n. [Gk. meta, after; zone, girdle.] The posterior ring of a diplosomite. Opp. prozonite.
- metazoon (mět'ázō'ŏn) n. [Gk. meta, after; zoon, animal.] A multicellular animal; metazoan.
- metembryo (mětěm'briö) n. [Gk.

- meta, towards; embryon, embryo.] The gastrula.
- metencephalon (mět'ěnkěf'álŏn, -sěf-) n. [Gk. meta, after; en, in; kephale, head.] Part of hind-brain, consisting of cerebellum, pons, and intermediate part of fourth ventricle; or hind-brain.
- metenteron (mětěn'tërŏn) n. [Gk. meta, after; enteron, gut.] Intermesenteric chamber of sea-anemone or other coelenterate.
- metepimeron (měť epimē'rŏn) n. [Gk. meta, after; epi, upon; meros, upper thigh.] Epimeron of insect metathorax.
- metepisternum (mět'épistěr'nům) n. [Gk. meta, after; epi, upon; sternon, breast-bone.] Episternum of insect metathorax; metaepisternum.
- metestrum,—metoestrus, q.v.
- methaemoglobin (mět-hē'möglō'bĭn)

 n. [Gk. meta, after; haima, blood;
 L. globus, globe.] An oxidation
 product of haemoglobin; HbO,
 HbOH.
- metochy (měť'ökĭ) n. [Gk. metoche, sharing.] Relationship between a neutral guest insect and its host.
- metoecious (mětē'sĭūs) a. [Gk. meta, after; oikos, house.] Metoxenous or heteroecious; with two hosts.
- metoestrus (mětē'strūs) n. [Gk. meta, after; oistros, gadfly.] The luteal phase, period when activity subsides after oestrus; metoestrum.
- metope (mět'ōpē) n. [Gk. metopon, forehead.] The middle frontal portion of a crustacean.
- metopic (mětőp'ík) a. [Gk. metopon, forehead.] Pert. forehead; appl. frontal suture.
- metopion (mětô'piŏn) n. [Gk. metopion, forehead.] Point on forehead where mid-sagittal plane intersects line connecting frontal eminences.
- metosteon (mětŏs'těŏn) n. [Gk. meta, after; osteon, bone.] A posterior sternal ossification in birds.
- metovum (mětő'vům) n. [Gk. meta, after; L. ovum, egg.] An egg-cell surrounded by nutritive material.
- metoxenous (mětox'ēnus) a. [Gk.

meta, after; xenos, guest.] Parasitic on different hosts at different stages in life-history.

metraterm (mē'trātērm) n. [Gk. metra, womb; terma, end.] Terminal portion of uterus in trematodes.

metula (mē'tūlă) n. [Dim. of L. meta, end-post.] A spore-bearing branch having flask-shaped outgrowths, as in certain fungi.

M-factor,—a certain antigen in erythrocytes of higher animals.

micella (misel'à) n. [L. dim. of mica, morsel.] Hypothetical unit, q.v.; an orderly aggregate of chain-like molecules.

micraesthetes (mīkrēsthē'tēz) n. plu. [Gk. mikros, small; aisthetes, perceiver.] The smaller sensory organs of Placophora.

micraner (mīkrā'nër) n. [Gk. mikros, small; aner, male.] A dwarf male ant. micrergate (mīkrĕr'gāt) n. [Gk. mikros, small; ergates, worker.] A dwarf worker ant.

micro-aerophiles, — organisms requiring less oxygen than is present in the air.

microbe (mī'krōb) n. [Gk. mikros, small; bios, life.] A bacterium; a micro-organism.

microbiology (mī'kröbīðl'öjǐ) n. [Gk. mikros, small; bios, life; logos, discourse.] Biology of microscopic organisms.

microblast (mī'kröblāst) n. [Gk. mikros, small; blastos, bud.] An erythroblast smaller than normal.

microcaltrops (mī'krokăl'trops) n. [Gk. mikros, small; A.S. coltraeppe, kind of thistle.] A primitive tetraxon, or cuaster with four persistent rays.

microcentrum (mi'krösen'trum) n.
[Gk. mikros, small; kentron, centre.] The dynamic centre of a cell, composed of centrosomes; kinetonucleus.

microcephalic (mī'krökēfāl'īk, -sĕf-)
a. [Gk. mikros, small; kephale, head.] With abnormally small head; having a cranial capacity of under 1350 c.c.; cf. megacephalic, mesocephalic.

microchromosome (mī'krökrō'mösōm) n. [Gk. mikros, small; chroma, colour; soma, body.] A chromosome considerably smaller than the other chromosomes of the same type of nucleus; M-chromosome.

microconidium (mī'krokŏnīd'iŭm) n. [Gk. mikros, small; konis, dust; idion, dim.] A comparatively small conidium.

microconjugant (mī'krökŏn'joogănt)

n. [Gk. mikros, small; L. conjugare, to unite.] A motile ciliated
free-swimming conjugant or gamete
which attaches itself to a macroconjugant and fertilises it.

microcyst (mi'krösist) n. [Gk. mikros, small; kystis, bladder.] A resting - spore stage of slime

fungi.

microcytase (mī'krösī'tās) n. [Gk. mikros, small; kytos, hollow.] The enzyme of microphages or smaller leucocytes.

microcytes (mī'krösīts) n. plu. [Gk. mikros, small; kytos, hollow.] Blood-corpuscles about half the size of erythrocytes, numerous in diseased conditions.

microdont (mī'krodont) a. [Gk. mikros, small; odous, tooth.] With comparatively small teeth.

microevolution (mī'kröēvölū'shūn)

n. [Gk. mikros, small; L. evolvere,
to unroll.] Evolutionary processes
that can be noticed within a relatively brief period, as during a
human life-time; evolution due to
gene mutation and recombination.
Opp. macro-evolution

microfilaria (mī'krofflā'rīā) n. [Gk. mikros, small; L. filum, thread.]
The embryo of a Filaria.

microgamete (mī'krögămēt') n. [Gk. mikros, small; gametes, spouse.]
The smaller of two conjugant gametes, regarded as male.

microgametoblast (mī'krögāmē'töblāst) n. [Gk. mukros, small; gametes, spouse; blastos, bud.] Intermediate stage between microgametocyte and microgamete in certain Sporozoa.

- microgametocyte (mī'krögămē'tösīt)

 n. [Gk. mikros, small; gametes,
 spouse; kytos, hollow.] Cell developed from merozoite in certain
 protozoa, giving rise to microgametes.
- microgametogenesis (mī'krogămē'töjēn'ēsis) n. [Gk. mikros, small; gametes, spouse; genesis, descent.] Development of microgametes or spermatozoa.
- microgamy (mikrŏg'āmi) n. [Gk. mikros, small; gamos, marriage.] Syngamy between smallest individuals produced by fission or gemmation, as in Foraminifera; merogamy.
- microglia (mīkröglī'ā, mī'krōglē'ā) n.

 plu. [Gk. mikros, small; glia,
 glue.] Mesodermal phagocytic
 cells in grey and white nervous
 matter; mesoglia.
- microgonidium (mī'krögŏnīd'iŭm) n. [Gk. mikros, small; gonos, off-spring; idion, dim.] A comparatively small gonidium; a male gamont or gametocyte.
- microgyne (mī'kröjīnē) n. [Gk. mi-kros, small; gyne, female.] Dwarf female ant.
- microhenad (mī'kröhēn'ād) n. [Gk. mikros, small; henas, unit.] A filter-passer.
- microlecithal (mī'krölĕs'ĭthăl) a. [Gk. mikros, small; lekithos, yolk.] Containing little yolk.
- microleucocyte (mī'krölū'kösīt, -loo-)

 n. [Gk. mikros, small; leukos,
 white; kytos, hollow.] A small
 amoebocyte.
- micromere (mī'krömēr) n. [Gk. mikros, small; meros, part.] A cell of upper or animal hemisphere in meroblastic and other eggs.
- micromerozoite (mī'krömer'özōīt) n.
 [Gk. mikros, small; meros, part; zoon, animal.] Cell derived from microschizont and developing into gametocyte in Haemosporidia.
- micromesentery (mī'kröměs'éntërī, -měz-) n. [Gk. mikros, small; mesos, middle; enteron, gut.] A secondary incomplete mesentery in Zoantharia. micromutation (mī'krömūtā'shūn)

n. [Gk. mikros, small; L. mutare, to change.] Mutation at only one gene locus; genovariation, point mutation, transgenation.

- micromyelocyte (mī'krömī'ēlösīt) n. [Gk. mikros, small; myelos, marrow; kytos, hollow.] A small heterophil myelocyte.
- micron (mi'kron) n. [Gk. mikros, small.] Micromillimetre, one-thousandth part of a millimetre; symbol: μ.
- micronephridia (mī'krönĕfrid'ĭă) n.
 plu. [Gk. mikros, small; nephros,
 kidney; idion, dim.] Small
 nephridia; meronephridia.
- micront (mī'krŏnt) n. [Gk. mikros, small; on, being.] A small cell formed by schizogony, itself giving rise to microgametes.
- micronucleocyte (mī'krönū'kleösīt)

 n. [Gk. mikros, small; L. nucleus,
 kernel; Gk. kytos, hollow.] An
 amoebocyte with a relatively small
 nucleus.
- micronucleus (mī'kronū'kleūs) n. [Gk. mikros, small; L. nucleus, kernel.] The smaller, reproductive nucleus of many protozoa, in close proximity to meganucleus; gononucleus.
- micro-nutrients,—substances essential to health of organisms, but required in minute quantity; e.g. trace elements, q.v.
- micro-organism (mī'krö-ôr'gănīzm)

 n. [Gk. mikros, small; organon, instrument.] A microscopic organism; microbe; protist.
- microparasite (mī'kröpăr'āsīt) n. [Gk. mikros, small; para, beside; sitos, food.] A parasite of microscopic size.
- microphages (mī'kröfā'jēz) n. plu. [Gk. mikros, small; phagein, to eat.] Chiefly the polymorphonuclear heterophil leucocytes.
- microphagic (mi'kröfāj'īk) a. [Gk. mikros, small; phagein, to eat.] Feeding on minute organisms or particles, appl. protozoa; microphagous, feeding on small prey, appl. agnathous fishes. Opp. macrophagous.

- microphagocyte (mī'krofāg'osīt) n. [Gk. mikros, small; phagein, to eat; kytos, hollow.] A microphage or small phagocyte of blood.
- microphanerophyte (mī'kröfān'eröfīt) n. [Gk. mikros, small;
 phaneros, manifest; phyton, plant.]
 Tree or shrub from 2 to 8 metres
 in height.
- microphil (mī'kröfil) a. [Gk. mikros, small; philein, to love.] Tolerating only a narrow range of temperature, appl. certain bacteria; microphilic. n. Microphile.
- microphyllous (mī'kröfīl'ŭs) a. [Gk. mikros, small; phyllon, leaf.] With small leaves.
- microphyte (mī'kröfīt) n. [Gk. mikros, small; phyton, plant.] A microscopic plant.
- microphytology (mī'kröfītŏl'ojī) n. [Gk. mikros, small; phyton, plant; logos, discourse.] Science of microphytes; bacteriology.
- microplankton (mi'kroplang'kton)

 n. [Gk. mikros, small; plangktos,
 wandering.] Small organisms drifting with the surrounding water,
 somewhat larger than those of
 nanoplankton, q.v.
- micropodous (mīkrŏp'oduš) a. [Gk. mikros, small; pous, foot.] With rudimentary or small foot or feet.
- micropore (nī'kröpōr) n. [Gk. mikros, small; poros, channel.]
 A small pore in a Chiton shell, containing a sense-organ.
- micropterism (mikrop'terizm) n. [Gk. mikros, small; pteron, wing.] Condition of having unusually small wings, as in some insects.
- micropterous (mikröp'terus) a. [Gk. mikros, small; pteron, wing.] Having small hind wings invisible till tegmina are expanded, as in some insects; with small or rudimentary fins.
- micropyle (mi'kröpil) n. [Gk. mikros, small; pyle, gate.] Aperture for admission of pollen-tube at ovule apex; aperture between hilum and point of radicle; small opening in

- cyst wall of macrogamete, for entry of microgamete; pore of oocyst; aperture in egg-membrane for admission of spermatozoon; pore in spongin-coat of sponges for escape of gemmules.
- micropyle apparatus,—raised processes or porches, sometimes of elaborate structure, developed round micropyle of certain insect eggs.
- micropyrenic (mī'kröpīrē'nīk) a. [Gk. mikros, small; pyren, fruitstone.] With nuclet markedly smaller than average for the species or other group. n. A micropyrenic individual.
- microrhabdus (mī'krörāb'dūs) n. [Gk. mikros, small; rhabdos, rod.] Minute monaxon or rod-like spicule.
- microschizogony (mī'kroskīzog'önī)

 n. [Gk. mikros, small; schizein, to cleave; gonos, birth.] Schizogony resulting in small merozoites.
- microschizont (mī'kroskīzont') n. [Gk. mikros, small; schizein, to cut; onta, beings.] A male schizont of certain protozoa.
- microsclere (mī/krösklēr) n. [Gk. mikros, small; skleros, hard.] One of small spicules found lying scattered in tissues of sponges. Opp. megasclere.
- microseptum (mī'krösēp'tūm) n. [Gk. mikros, small; L. septum, partition.] An incomplete mesentery of Zoantharia.
- microsere (mī'krösēr) n. [Gk. mikros, small; L. serere, to put in a row.] A successional series of plant communities in a microhabitat.
- microsmatic (mī'krŏsmāt'īk) a. [Gk. mukros, small; osme, smell.] With feebly-developed sense of smell.
- microsome (mī'krösōm) n. [Gk. mikros, small; soma, body.] Granule of protoplasm as opposed to ground-substance; a minute particle, centre for enzyme localisation, in cytoplasm.
- microsorus (mī'krösōrūs) n. [Gk. mikros, small; soros, heap.] A sorus containing microsporangia, opp. megasorus.

- microspecies (mī'kröspē'shēz) [Gk. mikros, small; L. species, particular kind.] A small species, or subspecies, with little variability; Iordanon.
- microsphere (mī'krösfēr) n. [Gk. mikros, small; sphaira, globe.] The initial chamber of Foraminifera when very small; centrosphere.

microspheric (mī'krösfĕr'ik) a. [Gk. mikros, small; sphaira, globe.] Appl. foraminifera when initial chamber of shell is small.

microsplanchnic (mī'krösplangk'nik) [Gk. mikros, small; splangchnon, entrail.] Small-bodied and

long-legged.

microsporangium (mī'kröspörăn'jiŭm) n. [Gk. *mikros*, small; sporos, seed; anggeion, vessel.] A sporangium bearing a number of microspores; pollen sac or anther

lobe of phanerogams.

- microspore (mīkröspōr) n. mikros, small; sporos, seed.] The spore developed in a microsporangium of heterosporous plants; the cell from which a pollen grain develops; a pollen grain; androspore; the smaller anisospore of Sarcodina.
- microsporocyte (mī'kröspō'rösīt) n. [Gk. mikros, small; sporos, seed; kytos, hollow.] The pollen mother cell which produces microspores resulting from two meioses.

microsporophore (mī'kröspō'röfōr) [Gk. mikros, small; sporos, seed; pherein, to bear.] A micro-

sporangium.

microsporophyll (mī'kröspō'röfil) n. [Gk. mikros, small; sporos, seed; phyllon, leaf.] A microsporangiumbearing leaf; stamen.

microsporozoite (mī'kröspō'rözō'īt) n. [Gk. mikros, small; sporos, seed; zoon, animal.] A smaller endogenous sporozoite of Sporozoa. microstome (mī'kröstōm) n. [Gk.

mikros, small; stoma, mouth.] A

small opening or orifice.

microstrobilus (mī'kröstrob'ilus) n. [Gk. mikros, small; strobilos, cone.] A small cone, as in cycads.

- microstylospore (mī'kröstī'löspōr) n. [Gk. mikros, small; stylos, pillar; sporos, seed. Λ comparatively small stylospore.
- microstylous (mī'kröstīl'ŭs) a. [Gk. mikros, small; stylos, pillar.] Having short styles; appl. heterostylous flowers.
- microtaxonomy (mī'krötāksŏn'ömĭ) [Gk. mikros, small; taxis, arrangement; nomos, law.] Classification and its principles as applied to subspecies, varieties, or races.

microtherm (mī'krötherm) n. [Gk. mikros, small; therme, heat.] A plant of the cold temperate zone.

- microtrichia (mī'krötrīk'ĭă) n. plu. [Gk. mikros, small; thrix, hair.] Small hairs without basal articulation on insect wings.
- microtype (mī'krötīp) n. mikros, small; L. typus, type.] Normal mesentery arrangement of Anthozoa; cf. macrotype.
- microzoid ($m\bar{i}'kr\bar{o}z\bar{o}'\bar{i}d$) n. [Gk. mikros, small; zoon, animal; idion, Male dim.gamete, as algae.
- microzooid (mī'krözō'oid) n. [Gk. mikros, small; zoon, animal; eidos, form.] A free-swimming motile ciliated bud of Vorticella and other protozoa.
- microzoon (mī'krözō'ŏn) n. [Gk. mikros, small; zoon, animal.] A microscopic animal.
- microzoospore ($m\bar{i}'kr\bar{o}z\bar{o}'\bar{o}sp\bar{o}r$) n. [Gk. mikros, small; zoon, animal; sporos, seed.] Small planogamete; small anisospore of Radiolaria.
- microzyma (mī'krözī'mă) n. mikros, small; zyme, leaven.] A hypothetical ultimate unit, q.v.
- microzyme (mi'krözim) n. mikros, small; zyme, leaven.] micro-organism of fermenting or decomposing liquids.
- micton (mik'ton) n. [Gk. miktos, mixed; on, being.] A species resulting from interspecific hybridisation and of which the individuals are interfertile.
- micturition (mik'tūrish'ŭn) n. mingere, to void water.] Act